



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

EP 1 285 651 A1

(12)

**EUROPEAN PATENT APPLICATION**  
published in accordance with Art. 158(3) EPC

(43) Date of publication:  
26.02.2003 Bulletin 2003/09

(21) Application number: 01925947.2

(22) Date of filing: 26.04.2001

(51) Int Cl.7: **A61K 31/167**, A61K 31/40,  
A61K 31/4453, A61K 31/4709,  
A61K 31/4545, A61K 31/454,  
A61K 31/4375, A61K 31/47,  
A61K 31/4725, C07C 233/80,  
C07C 237/48, C07D 211/16,  
C07D 211/14, C07D 211/18,  
C07D 295/135, C07D 405/12,  
C07D 401/06, C07D 417/06,  
C07D 401/12, C07D 211/22,  
C07D 471/04

(86) International application number:  
PCT/JP01/03614

(87) International publication number:  
WO 01/082925 (08.11.2001 Gazette 2001/45)

(84) Designated Contracting States:  
**AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU**  
**MC NL PT SE TR**  
Designated Extension States:  
**AL LT LV MK RO SI**

(30) Priority: 28.04.2000 JP 2000134295  
13.12.2000 JP 2000384897


(71) Applicant: Takeda Chemical Industries, Ltd.  
Osaka-shi, Osaka 541-0045 (JP)

(72) Inventors:  
• ISHIHARA, Yuji  
Itami-shi, Hyogo 664-0874 (JP)  
• SUZUKI, Nobuhiko  
Minoo-shi, Osaka (JP)  
• TAKEKAWA, Shiro  
Nishinomiya, Hyogo (JP)

(74) Representative: Lewin, John Harvey  
Takeda Patent Office,  
11-12 Charles II Street  
London SW1Y 4QU (GB)

(54) **MELANIN CONCENTRATING HORMONE ANTAGONISTS**

(57) A melanin-concentrating hormone antagonist comprising a compound of the formula (I):



wherein Ar<sup>1</sup> is a cyclic group which may be substituted;

X and Y are the same or different and are a spacer having a main chain of 1 to 6 atoms; Ar is a condensed polycyclic aromatic ring which may be substituted; R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom, Y and Ar, may form a condensed ring; or a salt thereof is useful as an agent for preventing or treating obesity, etc.

## Description

## TECHNICAL FIELD

- 5 [0001] The present invention relates to a melanin-concentrating hormone antagonist which is useful as an agent for preventing or treating obesity, etc.

## BACKGROUND ART


- 10 [0002] Feeding behavior is an essential action for many living beings including humans. Therefore, if irregularities in feeding behavior occur, disorders, often connected to diseases, will occur in normal life-maintaining activities. Accompanying recent changes of our dietary environment, obesity is now becoming a social problem. In addition, not only is obesity a serious risk factor for life-style diseases such as diabetes, hypertension, and arteriosclerosis; it is also widely known that increased body weight places excessive burdens on joints such as knee joints, causing arthritis and pain. The "diet boom," etc. show that there is a potentially great percentage of the population hoping to reduce body weight; on the other hand, many cases of feeding problems such as overeating, occurring due to causes such as hereditary neurosis or neurosis due to stress, have been reported.

[0003] Therefore, research on and development of agents for preventing or treating obesity, or agents for inhibiting eating, have been vigorously done for a long time. The centrally acting anorectic drug, Mazindol, is now being marketed.

- 20 [0004] Many appetite control factors such as leptin, have recently been discovered, and the development of anti-obesity agents or anorectic agents which will regulate the functions of these appetite control factors is progressing. In particular, it is known that melanin-concentrating hormone (hereinafter also abbreviated as "MCH") originates in the hypothalamus and has orexigenic action. In addition, it has been reported that even though the daily behavior of MCH knock-out mice was normal, the amount of feeding by MCH knock-out mice was significantly reduced and their body weights were lighter than those of normal mice [Nature, Vol. 396, p.670, 1998]. This indicates that, if a MCH antagonist was produced, it can be expected to be an excellent anorectic agent or anti-obesity agent; but at present there are no known compound, especially non-peptide type compounds, which possess MCH antagonistic actions.


[0005] On the other hand, the following compounds are known as amine derivatives.

- 30 1) WO98/38156 describes a compound of the formula:




- 40 wherein Ar is an optionally substituted ring assembly aromatic group or an optionally substituted condensed aromatic group; X is a bond, etc.; Y is an optionally substituted bivalent  $C_{1-8}$  aliphatic hydrocarbon group which may have an intervening oxygen atom or sulfur atom;  $R^1$  and  $R^2$  are independently hydrogen atom or an optionally substituted lower alkyl, or  $R^1$  and  $R^2$ , together with the adjacent nitrogen atom, form an optionally substituted nitrogen-containing heterocyclic ring; Ring A is a benzene ring which may have further substituents in addition to the groups of the formula: -X-Ar where each symbol is as defined above; Ring B is a 4- to 8-membered ring which may have further substituents in addition to the group of the formula: -Y-NR<sup>1</sup>R<sup>2</sup> where each symbol is as defined above; with the proviso that, when the condensed ring formed by ring A and ring B is an indole ring, the group of the formula: -X-Ar where each symbol is as defined above is substituted at the 4-, 6-, or 7- position on the indole ring; or its salt, which has an action of inhibiting the production and secretion of  $\beta$ -amyloid protein.

- 50 2) WO95/3296 describes compound of the formula:




wherein A is CONR, in which R is hydrogen or C<sub>1-6</sub> alkyl; Q is an optionally substituted 5- to 7-membered heterocyclic ring containing 1 to 3 hetero atoms selected from oxygen, nitrogen or sulfur; R<sup>1</sup> is hydrogen, halogen, etc.; R<sup>2</sup> and R<sup>3</sup> are independently hydrogen, halogen, etc.; R<sub>4</sub> and R<sub>5</sub> are independently hydrogen or C<sub>1-3</sub> alkyl; R<sup>6</sup> is halogen, hydroxy, etc.; R<sub>7</sub> and R<sub>8</sub> are independently hydrogen, C<sub>1-6</sub> alkyls, etc.; m is 0 to 4; n is 0, 1 or 2; or its salt, which has 5HT<sub>1D</sub> antagonist activity and can be expected to ameliorate anorexia.

3) WO98/15274 describes a compound of the formula:




wherein Ar is phenyl, etc.; X is -O- or -S-; Y is CR<sup>5</sup>R<sup>6</sup>, where R<sup>5</sup> is H and R<sup>6</sup> is -H, etc.; Z is -CH<sub>2</sub>- or -N-; R is H or -C<sub>1</sub>-C<sub>6</sub> alkyl; R<sup>1</sup> and R<sup>2</sup> are independently -C<sub>1</sub>-C<sub>6</sub> alkyl, etc.; R<sup>3</sup> is H etc.; R<sup>4</sup> is hydrogen, etc.; m is an integer of 0 to 2; q is 0 or 1; n is an integer of 0 to 4; p is an integer of 1 to 6; t is an integer of 1 to 4; which has an anti-oxidant activity and can be expected to ameliorate Alzheimer's disease.

4) EP533266 describes a compound of the formula:




wherein R<sup>1</sup> is halogen, etc.; R<sup>2</sup> is phenyl optionally substituted by 1 or 2 substituents selected from halogen, etc.; R<sup>3</sup> is




; R<sup>4</sup> and R<sup>5</sup> are independently hydrogen, halogen, etc.; R<sup>11</sup> is hydrogen or C<sub>1-6</sub> alkyl; which has 5HT<sub>1D</sub> antagonist activity, and can be expected to ameliorate anorexia.

5) DE2502588 describes a compound of the formula:



wherein  $R^1$  is hydrogen, or lower alkyl such as Me, Et, etc.;  $NR^2R^3$  is  $NH_2$ , a primary amine such as  $NHMe$ , etc., a secondary amine such as  $NEt_2$ ,  $NBu_2$ , etc., or a cyclic amine such as pyrrolidinyl, piperidinyl, morpholinyl, etc.;  $R^4$  and  $R^5$  are hydrogen, lower alkyl such as Me, etc., lower alkoxy such as  $OMe$ , etc., or halogen;  $R^6$  is hydrogen, or lower alkyl such as Me, Et, etc.;  $R^7$  is H, lower alkyl such as Me, Et, etc.,  $COR^8$  ( $R^8$  is alkoxy, aryloxy,  $NR^9R^{10}$  ( $NR^9R^{10}$  is  $NH_2$ , an optionally substituted primary amine such as  $NHMe$ , etc., secondary amine such as  $NEt_2$ ,  $NBu_2$ , etc., or a cyclic amine such as pyrrolidinyl, piperidinyl, morpholinyl, etc.));  $A$  is an alkyl chain such as  $-CH_2-$ ,  $-CH_2CH_2-$ , etc., which can be expected to ameliorate hypertrophy of the small intestine.

6) J. Chem. Soc., 4678 (1962) or J. Heterocycl. Chem., 24, 345 (1987) describes a compound of the formula:




wherein  $R^1$  is hydrogen, or alkyl such as Me, Et, etc.;  $R^2$  is hydrogen, halogen or a carboxylic acid ester, which has folic acid antagonistic activity.

[0006] There has been great desire for the development of a melanin-concentrating hormone antagonist which is useful as an agent for preventing or treating obesity, excellent in oral absorberency, and safe.

#### DISCLOSURE OF INVENTION


[0007] As a result of intensive studies of compounds with a MCH antagonistic action, the present inventors found that a derivative which is obtained by introducing a group of the formula:  $Ar^1-X$  where each symbol is as defined hereafter, into a compound of the formula:



wherein each symbol is as defined hereinafter, had an excellent MCH antagonistic actions, to complete the present invention.

[0008] Namely, the present invention relates to:

1) A melanin-concentrating hormone antagonist which comprises a compound of the formula:



wherein Ar<sup>1</sup> is a cyclic group which may be substituted;

X and Y are the same or different and are a spacer having a main chain of 1 to 6 atoms;

Ar is a condensed polycyclic aromatic ring which may be substituted;


R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom, Y and Ar, may form a nitrogen-containing condensed ring which may be substituted; or a salt thereof;

2) The antagonist according to the above 1), wherein R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, form a nitrogen-containing heterocyclic ring which may be substituted;

3) The antagonist according to the above 1) which is an agent for preventing or treating diseases caused by melanin-concentrating hormone;

4) The antagonist according to the above 1) which is an agent for preventing or treating obesity;

5) A compound of the formula:



wherein Ar<sup>1</sup> is a cyclic group which may be substituted;

X<sup>1</sup> is CONR<sup>8</sup>, NR<sup>8</sup>CO (wherein R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl or optionally halogenated C<sub>1-6</sub> alkylsulfonyl), OCO or COO;

Y is a spacer having a main chain of 1 to 6 atoms;

Ar is a condensed polycyclic aromatic ring which may be substituted;

R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom, Y and Ar, may form a nitrogen-containing condensed ring which may be substituted;

provided that, when X<sup>1</sup> is CONR (wherein R is hydrogen atom or C<sub>1-6</sub> alkyl), Ar is not indole or benzoxazole which may have one or two halogen, hydroxy, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> alkoxy; when X<sup>1</sup> is CONH, Ar is not 4-methyl-2-quinolone which may have a substituent selected from the group consisting of alkyl, alkoxy and halogen, or is not 2-benzoylamino-quinazoline; and, when X<sup>1</sup> is COO, Ar<sup>1</sup> is not an aromatic group which may be substituted; or a salt thereof;

6) The compound according to the above 5), wherein X<sup>1</sup> is CONR<sup>8</sup> or NR<sup>8</sup>CO (wherein R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl or optionally halogenated C<sub>1-6</sub> alkylsulfonyl); and R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be

substituted; or R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted;

- 7) The compound according to the above 5), wherein the cyclic group represented by Ar<sup>1</sup> is an aromatic group;
- 8) The compound according to the above 7), wherein the aromatic group is formed by removing an optional one hydrogen atom from an aromatic ring assembly formed by 2 or 3 members selected from C<sub>6-14</sub> monocyclic or condensed polycyclic aromatic hydrocarbon and 5- to 10-membered aromatic heterocyclic ring;
- 9) The compound according to the above 5), wherein Ar<sup>1</sup> is phenyl, biphenyl or phenyl-pyridyl, each of which may be substituted with 1 to 3 substituents selected from halogen, optionally halogenated C<sub>1-6</sub> alkyl and optionally halogenated C<sub>1-6</sub> alkoxy;
- 10) The compound according to the above 5), wherein Ar<sup>1</sup> is piperidinyl which may be substituted with C<sub>6-14</sub> aryl which may be substituted;
- 11) The compound according to the above 5), wherein X<sup>1</sup> is CONH or COO;
- 12) The compound according to the above 5), wherein the condensed polycyclic aromatic ring represented by Ar is a condensed polycyclic aromatic hydrocarbon having 9 to 14 carbon atoms;
- 13) The compound according to the above 4), wherein the condensed polycyclic aromatic ring represented by Ar is a 10-membered condensed polycyclic aromatic heterocyclic ring;
- 14) The compound according to the above 5), wherein the condensed polycyclic aromatic ring represented by Ar is quinoline or naphthalene;
- 15) The compound according to the above 5), wherein X<sup>1</sup> is CONR<sup>8</sup> or NR<sup>8</sup>CO (wherein R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl or optionally halogenated C<sub>1-6</sub> alkyl-sulfonyl), and Ar is quinoline or naphthalene;
- 16) The compound according to the above 4), wherein the "spacer having a main chain of 1 to 6 atoms" represented by Y is a bivalent group consisting of 1 to 3 species selected from -O-, -S-, -CO-, -SO-, -SO<sub>2</sub>-, -NR<sup>8</sup>- (R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl, optionally halogenated C<sub>1-6</sub> alkylsulfonyl), and an optionally halogenated bivalent C<sub>1-6</sub> non-cyclic hydrocarbon group;
- 17) The compound according to the above 5), wherein Y is C<sub>1-3</sub> alkylene;
- 18) The compound according to the above 4), wherein R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, form a nitrogen-containing heterocyclic ring which may be substituted;
- 19) The compound according to the above 18), wherein the nitrogen-containing heterocyclic ring is morpholine, piperidine, piperazine, pyrrolidine, 1,3-thiazolidine, 1H-imidazole, 4,5-dihydro-1H-imidazole, 2,3-dihydroindole, 1,2,3,4-tetrahydroquinoline or 1,2,3,4-tetrahydroisoquinoline;
- 20) A pharmaceutical composition comprising the compound according to the above 5), or a salt thereof.
- 21) A prodrug of the compound according to the above 5).
- 22) The compound according to the above 5) which is:

4'-chloro-N-[6-[(N,N-dimethylamino)methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 4'-fluoro-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4'-fluoro-N-[2-(1-piperidinylmethyl)-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[2-[(2-methyl-4,5-dihydro-1H-imidazol-1-yl)methyl]-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4-(4-chlorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl]-1-piperidinecarboxamide;  
 N-[2-(1-pyrrolidinylmethyl)-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 6-(4-methylphenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl]nicotinamide;  
 6-(4-methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide;  
 6-(4-methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide;  
 6-(4-methylphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide;

or a salt thereof;

23) A process for producing a compound of the formula (I), or a salt thereof, which comprises reacting a compound of the formula:



(XII)

wherein Ar<sup>1</sup> is as defined in the above 5), or a salt thereof with a compound of the formula:

24) The antagonist according to the above 1) which is an anorectic agent:

26) A method for preventing or treating diseases caused by a melanin-concentrating hormone in a mammal in need thereof, which comprises administering to said mammal an effective amount of the compound represented by the formula (I), or a salt thereof:

27) A method for preventing or treating obesity in a mammal in need thereof, which comprises administering to said mammal an effective amount of the compound represented by the formula (I), or a salt thereof:

28) Use of the compound represented by the formula (I), or a salt thereof for the manufacture of a pharmaceutical preparation for preventing or treating diseases caused by a melanin-concentrating hormone;

29) Use of the compound of the formula (I) or a salt thereof for the manufacture of a pharmaceutical preparation for preventing or treating obesity; and the like.

[0010] Here, examples of the "aromatic groups" include monocyclic aromatic groups, condensed aromatic groups, ring assembly aromatic groups and the like.

[0012] Examples of the "5- or 6-membered aromatic heterocyclic ring" include a 5- or 6-membered aromatic heterocyclic ring containing one or more (for example, 1 to 3) hetero atoms selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms, and the like. Specifically, thiophene, furan, pyrrole, imidazole, pyrazole, thiazole, isothiazole, oxazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, 1,2,4-oxadiazole, 1,3,4-oxadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, furan, etc., can be mentioned.

[0014] The "condensed aromatic groups" mean a univalent group that can be formed by removing an optional one hydrogen atom from condensed polycyclic (preferably bicyclic to tetracyclic, more preferably bicyclic or tricyclic) aromatic rings, etc.

[0016] Examples of the "condensed polycyclic aromatic hydrocarbons" include C<sub>9-14</sub> condensed polycyclic (bicyclic or tricyclic) aromatic hydrocarbons (e.g. naphthalene, indene, fluorene, anthracene, etc.), etc.

[0017] Examples of the "condensed polycyclic aromatic heterocyclic rings" include 9- to 14-membered, preferably, 9- or 10-membered, condensed polycyclic aromatic heterocyclic rings containing one or more (for example, 1 to 4) hetero atoms selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms, and the like. The "condensed polycyclic aromatic heterocyclic rings" is preferably 10-membered condensed polycyclic aromatic heterocyclic ring. Specific examples of the "condensed polycyclic aromatic heterocyclic rings" include benzofuran, benzimidazole, benzoxazole, benzothiazole, benzisothiazole, naphtho[2,3-b]thiophene, isoquinoline, quinoline, indole, quinoxaline, phenanthridine, phenothiadine, phenoxazine, phthalazine, naphthyridine, quinoxaline, cinnoline, carbazole,  $\beta$ -carboline, acridine, phenazine, phthalimide, thioxanthene, etc.

[0018] Specific examples of the "condensed aromatic groups" include 1-naphthyl; 2-naphthyl; 2-, 3-, 4-, 5- or 8-quinolyl; 1-, 3-, 4-, 5-, 6-, 7- or 8-isoquinolyl; 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl; 1-, 2-, 4- or 5-isindolyl; 1-, 5- or 6-phthalazinyl;

2-, 3- or 5-quinoxaliny; 2-, 3-, 4-, 5- or 6-benzofuranyl; 2-, 4-, 5- or 6-benzothiazolyl; 1-, 2-, 4-, 5- or 6-benzimidazolyl; etc.  
 [0019] The "ring assembly aromatic group" means a group formed by removing an optional one hydrogen atom from an aromatic ring assembly in which 2 or more (preferably 2 or 3) aromatic rings are directly bonded by single bonds, and in which the number of bonds which directly bond the rings, is less by one than the number of ring systems.

[0020] Examples of the aromatic ring assembly include an aromatic ring assemblies formed by 2 or 3 (preferably 2) species selected from  $C_{6-14}$  monocyclic or condensed polycyclic aromatic hydrocarbons (e.g. benzene and naphthalene) and 5- to 10-membered (preferably 5- or 6-membered) aromatic heterocyclic rings, etc.

[0021] Preferred example of the aromatic ring assemblies include aromatic ring assemblies comprising 2 or 3 aromatic rings selected from benzene, naphthalene, pyridine, pyrimidine, thiophene, furan, thiazole, isothiazole, oxazole, isoxazole, 1,2,4-oxadiazole, 1,3,4-oxadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, quinoline, isoquinoline, indole, benzo-thiophene, benzoxazole, benzothiazole and benzofuran.

[0022] Specific examples of the "ring assembly aromatic groups" include 2-, 3- or 4-biphenyl; 3-(1-naphthyl)-1,2,4-oxadiazol-5-yl; 3-(2-naphthyl)-1,2,4-oxadiazol-5-yl; 3-(2-benzofuranyl)-1,2,4-oxadiazol-5-yl; 3-phenyl-1,2,4-oxadiazol-5-yl; 3-(2-benzoxazolyl)-1,2,4-oxadiazol-5-yl; 3-(3-indolyl)-1,2,4-oxadiazol-5-yl; 4-phenylthiazol-2-yl; 4-(2-benzofuranyl)thiazol-2-yl; 4-phenyl-1,3-oxazol-5-yl; 5-phenyl-isothiazol-4-yl; 5-phenyl-oxazol-2-yl; 4-(2-thienyl)phenyl; 4-(3-thienyl)phenyl; 3-(3-pyridyl)phenyl; 4-(3-pyridyl)phenyl; 6-phenyl-3-pyridyl; 5-phenyl-1,3,4-oxadiazol-2-yl; 4-(2-naphthyl)phenyl; 4-(2-benzofuranyl)phenyl; 4,4'-terphenyl; 5-phenyl-2-pyridyl; 2-phenyl-5-pyrimidinyl; 4-(4-pyridyl)phenyl; 2-phenyl-1,3-oxazol-5-yl; 2,4-diphenyl-1,3-oxazol-5-yl; 3-phenyl-isoxazol-5-yl; 5-phenyl-2-furyl; 4-(2-furyl)phenyl; etc.

[0023] Preferred groups among the above "aromatic groups" are "a group formed by removing an optional one hydrogen atom from an aromatic ring assembly formed by 2 or 3 members selected from a  $C_{6-14}$  monocyclic or condensed polycyclic aromatic hydrocarbon and 5- to 10-membered aromatic heterocyclic ring (preferably, 2-, 3- or 4-biphenyl; 6-phenyl-3-pyridyl, 5-phenyl-2-pyridyl, etc.)."

[0024] Examples of the "non-aromatic cyclic hydrocarbon groups" include  $C_{3-8}$  cycloalkyl,  $C_{3-8}$  cycloalkenyl, etc.

[0025] Here, specific examples of the  $C_{3-8}$  cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, etc.

[0026] Specific examples of the  $C_{3-8}$  cycloalkenyl include cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, etc.

[0027] Among the above "non-aromatic cyclic hydrocarbon groups",  $C_{3-8}$  cycloalkyl is preferred, and cyclohexyl is particularly preferred.

[0028] Examples of "non-aromatic heterocyclic groups" include monocyclic non-aromatic heterocyclic groups, condensed polycyclic non-aromatic heterocyclic groups, and the like.

[0029] Examples of the "monocyclic non-aromatic heterocyclic groups" include univalent groups formed by removing an optional one hydrogen atom from monocyclic non-aromatic heterocyclic ring. Examples of the "monocyclic non-aromatic heterocyclic groups" include 5- to 8-membered monocyclic non-aromatic heterocyclic groups containing one or more (e.g. 1 to 3) hetero atoms selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms. Specifically, tetrahydrothiophene, tetrahydrofuran, pyrrolidine, imidazoline, imidazolidine, pyrazoline, pyrazolidine, tetrahydrothiazole, tetrahydroisothiazole, tetrahydrooxazole, tetrahydroisoxazole, piperidine, tetrahydropyridine, dihydropyridine, piperazine, morpholine, thiomorpholine, tetrahydropyrimidine, tetrahydropyridazine, hexamethylenimine, 1,3-dioxane, 1,4-dioxane, etc. can be mentioned.

[0030] The "condensed polycyclic non-aromatic heterocyclic group" means a univalent group formed by removing an optional one hydrogen atom from a condensed polycyclic (preferably bicyclic to tetracyclic, more preferably bicyclic or tricyclic) non-aromatic heterocyclic ring. Examples of the "condensed polycyclic non-aromatic heterocyclic ring" include 9- to 14-membered, preferably 9- or 10-membered condensed polycyclic non-aromatic heterocyclic rings which contain one or more (e.g. 1 to 4) hetero atoms selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms. Specifically, dihydrobenzofuran, dihydrobenzimidazole, dihydrobenzoxazole, dihydrobenzothiazole, dihydrobenzisoquinoline, dihydrophenanthridine, hexahydrophenanthridine, hexahydrophenoxazine, tetrahydrophenanthridine, tetrahydroquinoline, tetrahydroquinazoline, tetrahydrocinoline, tetrahydrocarbazole, tetrahydro- $\beta$ -carboline, tetrahydroacridine, tetrahydrophenazine, tetrahydrothioxanthene, etc., can be mentioned.

[0031] Among the above "non-aromatic heterocyclic groups", "5- to 8-membered monocyclic non-aromatic heterocyclic groups (preferably piperidinyl (piperidino); piperazinyl; pyrrolidinyl; 1,3-dioxanyl; etc. are preferred.

[0032] The "cyclic group" represented by  $Ar^1$  is preferably monocyclic aromatic groups (preferably phenyl), ring assembly aromatic groups (preferably biphenyl, phenylpyridyl), 5- to 8-membered monocyclic non-aromatic heterocyclic groups (preferably piperidinyl (piperidino), 1,3-dioxane), etc.

[0033] Examples of the "substituent" in the "cyclic group which may be substituted" represented by  $Ar^1$  include oxo, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.),  $C_{1-3}$  alkyleneedioxy (e.g. methyleneedioxy, ethyleneedioxy, etc.), nitro, cyano, optionally halogenated  $C_{1-6}$  alkyl, hydroxy- $C_{1-6}$  alkyl,  $C_{6-14}$  aryloxy- $C_{1-6}$  alkyl (e.g. phenoxyethyl,



etc.), C<sub>1-6</sub> alkyl-C<sub>6-14</sub> aryl-C<sub>2-6</sub> alkenyl (e.g. methylphenylethenyl, etc.), optionally halogenated C<sub>3-6</sub> cycloalkyl, optionally halogenated C<sub>1-6</sub> alkoxy, optionally halogenated C<sub>1-6</sub> alkylthio, C<sub>7-19</sub> aralkyl which may be substituted, hydroxy, C<sub>6-14</sub> aryloxy which may be substituted, C<sub>7-19</sub> aralkyloxy which may be substituted, C<sub>6-14</sub> aryl-carbamoyl which may be substituted, amino, amino-C<sub>1-6</sub> alkyl (e.g. aminomethyl, aminoethyl, aminopropyl, aminobutyl, etc.), mono-C<sub>1-6</sub> alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di-C<sub>1-6</sub> alkylamino (e.g. dimethylamino, diethylamino, dipropylamino, dibutylamino, ethylmethylamino, etc.), mono-C<sub>1-6</sub> alkylamino-C<sub>1-6</sub> alkyl (e.g. methylaminomethyl, ethylaminomethyl, propylaminomethyl, isopropylaminomethyl, butylaminomethyl, etc.), di-C<sub>1-6</sub> alkylamino-C<sub>1-6</sub> alkyl (e.g. dimethylaminomethyl, diethylaminomethyl, dipropylaminomethyl, diisopropylaminomethyl, dibutylaminomethyl, etc.), 5- to 7-membered saturated cyclic amino which may be substituted, 5- to 7-membered non-aromatic heterocyclic groups which may be substituted, acyl, acylamino, acyloxy, etc.

[0034] The "cyclic group" represented by Ar<sup>1</sup> may have 1 to 5, preferably 1 to 3, of the above-mentioned substituents at a substitutable position on the cyclic group. When the number of substituents is 2 or more, each substituents can be the same or different.

[0035] Also, when the "cyclic group" represented by Ar<sup>1</sup> is a non-aromatic cyclic hydrocarbon group or a non-aromatic heterocyclic group, the "cyclic group" may have as its substituent(s), C<sub>6-14</sub> aryl which may be substituted, 5- to 10-membered aromatic heterocyclic groups which may be substituted, etc.

[0036] Here, the groups exemplified as the "substituent" in the "5- to 7-membered saturated cyclic amino which may be substituted" mentioned hereinafter, can be mentioned as "C<sub>6-14</sub> aryl which may be substituted" and "5- to 10-membered aromatic heterocyclic groups which may be substituted". The number of substituents is, for example, 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0037] Specific examples of the above "optionally halogenated C<sub>1-6</sub> alkyl" include C<sub>1-6</sub> alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.). Specific examples include methyl, chloromethyl, difluoromethyl, trichloromethyl, trifluoromethyl, ethyl, 2-bromoethyl, 2,2,2-trifluoroethyl, pentafluoroethyl, propyl, 3,3,3-trifluoropropyl, isopropyl, butyl, 4,4,4-trifluorobutyl, isobutyl, sec-butyl, tert-butyl, pentyl, isopentyl, neopentyl, 5,5,5-trifluoropentyl, hexyl, 6,6,6-trifluorohexyl, etc.

[0038] The C<sub>1-6</sub> alkyl in the above "optionally halogenated C<sub>1-6</sub> alkyl" can be mentioned as the C<sub>1-6</sub> alkyl in the above "hydroxy-C<sub>1-6</sub> alkyl".

[0039] Examples of the above "optionally halogenated C<sub>3-6</sub> cycloalkyl" include C<sub>3-6</sub> cycloalkyl (e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.) which may have 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.). Specific examples include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, 4,4-dichlorocyclohexyl, 2,2,3,3-tetrafluorocyclopentyl, 4-chlorocyclohexyl, etc.

[0040] Examples of the above "optionally halogenated C<sub>1-6</sub> alkoxy" include C<sub>1-6</sub> alkoxy (e.g. methoxy, ethoxy, propoxy, butoxy, pentyloxy, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.). Specific examples include methoxy, difluoromethoxy, trifluoromethoxy, ethoxy, 2,2,2-trifluoroethoxy, propoxy, isopropoxy, butoxy, 4,4,4-trifluorobutoxy, isobutoxy, sec-butoxy, pentyloxy, hexyloxy, etc.

[0041] Examples of the above "optionally halogenated C<sub>1-6</sub> alkylthio" include C<sub>1-6</sub> alkylthio (e.g. methylthio, ethylthio, propylthio, isopropylthio, butylthio, sec-butylthio, tert-butylthio, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.). Specific examples include methylthio, difluoromethylthio, trifluoromethylthio, ethylthio, propylthio, isopropylthio, butylthio, 4,4,4-trifluorobutylthio, pentylthio, hexylthio, etc.

[0042] Examples of the "C<sub>7-19</sub> aralkyl" in the above "C<sub>7-19</sub> aralkyl which may be substituted" include benzyl, phenethyl, diphenylmethyl, triphenylmethyl, 1-naphthylmethyl, 2-naphthylmethyl, 2,2-diphenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, etc. Benzyl is particularly preferred.

[0043] Examples of the "substituent" in the above "C<sub>7-19</sub> aralkyl which may be substituted" include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), C<sub>1-3</sub> alkylenedioxy (e.g. methylenedioxy, ethylenedioxy, etc.), nitro, cyano, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>3-6</sub> cycloalkyl, optionally halogenated C<sub>1-6</sub> alkoxy, optionally halogenated C<sub>1-6</sub> alkylthio, hydroxy, amino, mono-C<sub>1-6</sub> alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di-C<sub>1-6</sub> alkylamino (e.g. dimethylamino, diethylamino, dipropylamino, dibutylamino, ethylmethylamino, etc.), amino-C<sub>1-6</sub> alkyl (e.g. aminomethyl, aminoethyl, aminopropyl, aminobutyl, etc.), mono-C<sub>1-6</sub> alkylamino-C<sub>1-6</sub> alkyl (e.g. methylaminomethyl, ethylaminomethyl, propylaminomethyl, isopropylaminomethyl, butylaminomethyl, etc.), di-C<sub>1-6</sub> alkylamino-C<sub>1-6</sub> alkyl (e.g. dimethylaminomethyl, diethylaminomethyl, dipropylaminomethyl, diisopropylaminomethyl, dibutylaminomethyl, etc.), formyl, carboxy, carbamoyl, thiocarbamoyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl, C<sub>1-6</sub> alkoxy-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), mono-C<sub>1-6</sub> alkyl-carbamoyl (e.g., methylcarbamoyl, ethylcarbamoyl, etc.), di-C<sub>1-6</sub> alkyl-carbamoyl (e.g. dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl, etc.), optionally halogenated C<sub>1-6</sub> alkylsulfonfyl, formylamino, optionally halogenated C<sub>1-6</sub> alkyl-carboxamide, C<sub>1-6</sub> alkoxy-carboxamide (e.g. methoxycarboxamide, ethoxycarboxamide, propoxycarboxamide, butoxycarboxamide, etc.), C<sub>1-6</sub> alkylsulfonfylamino (e.g. methylsulfonfylamino, ethylsulfonfylamino, etc.), C<sub>1-6</sub> alkyl-carboxyloxy (e.g. acetoxycarboxyloxy, etc.), C<sub>1-6</sub> alkoxy-carboxyloxy (e.g. methoxycarboxyloxy, etc.).

onyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, butoxycarbonyloxy, etc.), mono-C<sub>1-6</sub> alkyl-carbamoyloxy (e.g. methylcarbamoyloxy, ethylcarbamoyloxy, etc.), di-C<sub>1-6</sub> alkyl-carbamoyloxy (e.g. dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.), etc. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0044] As the "optionally halogenated C<sub>1-6</sub> alkyl", "optionally halogenated C<sub>3-6</sub> cycloalkyl", "optionally halogenated C<sub>1-6</sub> alkoxy" and "optionally halogenated C<sub>1-6</sub> alkylthio", those exemplified as the "substituents" in the above "cyclic group which may be substituted" can be used, respectively.

[0045] Examples of the above "optionally halogenated C<sub>1-6</sub> alkyl-carbonyl" include C<sub>1-6</sub> alkyl-carbonyl (e.g. acetyl, propanoyl, butanoyl, pentanoyl, hexanoyl, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.), etc. Specific examples include acetyl, monochloroacetyl, trifluoroacetyl, trichloroacetyl, propanoyl, butanoyl, pentanoyl, hexanoyl, etc.

[0046] Examples of the above "optionally halogenated C<sub>1-6</sub> alkylsulfonyle" include C<sub>1-6</sub> alkylsulfonyle (e.g. methylsulfonyle, ethylsulfonyle, propylsulfonyle, isopropylsulfonyle, butylsulfonyle, sec-butylsulfonyle, tert-butylsulfonyle, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g., fluorine, chlorine, bromine, iodine, etc.), etc. Specific examples include methylsulfonyle, difluoromethylsulfonyle, trifluoromethylsulfonyle, ethylsulfonyle, propylsulfonyle, isopropylsulfonyle, butylsulfonyle, 4,4,4-trifluorobutylsulfonyle, pentylsulfonyle, hexylsulfonyle, etc.

[0047] Examples of the above "optionally halogenated C<sub>1-6</sub> alkyl-carboxamide" include C<sub>1-6</sub> alkyl-carboxamide (e.g. acetamide, propanamide, butanamide, etc.) which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.), etc. Specific examples include acetamide, trifluoroacetamide, propanamide, butanamide, etc.

[0048] Examples of the "C<sub>6-14</sub> aryloxy" in the above "C<sub>6-14</sub> aryloxy which may be substituted" include phenyloxy, 1-naphthoxy, 2-naphthoxy, etc.

[0049] Examples of the "C<sub>7-19</sub> aralkyloxy" in the above "C<sub>7-19</sub> aralkyloxy which may be substituted" include benzyloxy, phenethyloxy, diphenylmethyloxy, triphenylmethyloxy, 1-naphthylmethyloxy, 2-naphthylmethyloxy, 2,2-diphenylethyloxy, 3-phenylpropyloxy, 4-phenylbutyloxy, 5-phenylpentyloxy, etc.

[0050] Examples of the "C<sub>6-14</sub> aryl-carbamoyl" in the above "C<sub>6-14</sub> aryl-carbamoyl which may be substituted" include phenylcarbamoyl, 1-naphthylcarbamoyl, 2-naphthylcarbamoyl, etc.

[0051] As the "substituents" in the "C<sub>6-14</sub> aryloxy which may be substituted", "C<sub>7-19</sub> aralkyloxy which may be substituted" and "C<sub>6-14</sub> aryl-carbamoyl which may be substituted", those exemplified for the "substituents" in the above "C<sub>7-19</sub> aralkyl which may be substituted" can be used. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0052] Examples of the "5- to 7-membered saturated cyclic amino" in the above "5- to 7-membered saturated cyclic amino which may be substituted" include morpholino, thiomorpholino, piperazin-1-yl, piperidino, pyrrolidin-1-yl, etc. The "5- to 7-membered saturated cyclic amino" can be condensed with a benzene ring.

[0053] Examples of the "substituent" in the "5- to 7-membered saturated cyclic amino which may be substituted" include oxo, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl, optionally halogenated C<sub>1-6</sub> alkylsulfonyle, C<sub>6-14</sub> aryl which may be substituted, C<sub>7-19</sub> aralkyl which may be substituted, C<sub>6-14</sub> aryl-carbonyl which may be substituted, 5- to 10-membered aromatic heterocyclic group which may be substituted, C<sub>1-6</sub> alkoxy-C<sub>1-6</sub> alkyl (e.g., methoxymethyl, ethoxymethyl), C<sub>1-6</sub> alkoxy-carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl), etc. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0054] Here, as the "optionally halogenated C<sub>1-6</sub> alkyl" and "C<sub>7-19</sub> aralkyl which may be substituted", those exemplified as the "substituents" in the above "cyclic group which may be substituted" can be used, respectively.

[0055] As the "optionally halogenated C<sub>1-6</sub> alkyl-carbonyl" and "optionally halogenated C<sub>1-6</sub> alkylsulfonyle", those exemplified for the "substituents" in the above "C<sub>7-19</sub> aralkyl which may be substituted" can be used.

[0056] Examples of the "C<sub>6-14</sub> aryl" in the "C<sub>6-14</sub> aryl which may be substituted" include phenyl, 1-naphthyl, 2-naphthyl, 2-indenyl, 2-anthryl, etc. Phenyl is especially preferable.

[0057] As the "substituents" in the "C<sub>6-14</sub> aryl which may be substituted", those exemplified as the "substituents" in the above "C<sub>7-19</sub> aralkyl which may be substituted" can be used. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0058] Examples of the "C<sub>6-14</sub> aryl-carbonyl" in the "C<sub>6-14</sub> aryl-carbonyl which may be substituted" include benzoyl, 1-naphthoyl, 2-naphthoyl, etc.

[0059] As the "substituents" in the "C<sub>6-14</sub> aryl-carbonyl which may be substituted", those exemplified as "substituents" in the above "C<sub>7-19</sub> aralkyl which may be substituted" can be used. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0060] Examples of the "5- to 10-membered aromatic heterocyclic groups" in the "5- to 10-membered aromatic heterocyclic groups which may be substituted" include 5- to 10-membered (monocyclic or bicyclic) aromatic heterocyclic groups containing 1 or 2 kinds of, preferably 1 to 4 hetero atoms selected from nitrogen, sulfur and oxygen atom in

addition to carbon atoms. Specific examples include 2- or 3-thienyl; 2-, 3- or 4-pyridyl; 2- or 3-furyl; 2-, 4- or 5-thiazolyl; 2-, 4- or 5-oxazolyl; 1-, 3- or 4-pyrazolyl; 2-pyrazinyl; 2-, 4- or 5-pyrimidinyl; 1-, 2- or 3-pyrrolyl; 1-, 2- or 4-imidazolyl; 3- or 4-pyridazinyl; 3-isothiazolyl; 3-isoxazolyl; 1,2,4-oxadiazol-5-yl, 1,2,4-oxadiazol-3-yl; 2-, 3-, 4-, 5- or 8-quinolyl; 1-, 3-, 4-, 5-, 6-, 7- or 8-isouinolyl; 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl; 1-, 2-, 4- or 5-isindolyl; 1-, 5- or 6-phthalazinyl; 2-, 3- or 5-quinolalinalyl; 2-, 3-, 4-, 5- or 6-benzofuranyl; 2-, 4-, 5- or 6-benzothiazolyl; 1-, 2-, 4-, 5- or 6-benzimidazolyl, etc.

[0061] Examples of the "substituents" in the "5- to 10-membered aromatic heterocyclic groups which may be substituted" include halogen atom (e.g. fluorine, chlorine, bromine and iodine, etc.), C<sub>1-3</sub> alkyleneedioxy (e.g. methyleneedioxy, ethyleneedioxy, etc.), nitro, cyano, optionally halogenated C<sub>1-6</sub> alkyl, C<sub>6-14</sub> arylalkoxy-C<sub>1-6</sub> alkyl (e.g. phenoxymethyl, etc.), C<sub>1-6</sub> alkyl-C<sub>6-14</sub> aryl-C<sub>2-6</sub> alkenyl (e.g. methylphenylethenyl, etc.), optionally halogenated C<sub>3-6</sub> cycloalkyl, optionally halogenated C<sub>1-6</sub> alkoxy, optionally halogenated C<sub>1-6</sub> alkylthio, C<sub>7-19</sub> aralkyl which may be substituted, hydroxy, C<sub>6-14</sub> arylalkoxy which may be substituted, C<sub>7-19</sub> aralkyloxy which may be substituted, amino, amino-C<sub>1-6</sub> alkyl (e.g. aminomethyl, aminoethyl, aminopropyl, aminobutyl, etc.), mono-C<sub>1-6</sub> alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di-C<sub>1-6</sub> alkylamino (e.g. dimethylamino, diethylamino, dipropylamino, dibutylamino, ethylmethylamino, etc.), mono-C<sub>1-6</sub> alkylamino-C<sub>1-6</sub> alkyl (e.g. methylaminomethyl, ethylaminomethyl, propylaminomethyl, isopropylaminomethyl, butylaminomethyl, etc.), di-C<sub>1-6</sub> alkylamino-C<sub>1-6</sub> alkyl (e.g. dimethylaminomethyl, diethylaminomethyl, dipropylaminomethyl, diisopropylaminomethyl, dibutylaminomethyl, etc.), 5- to 7-membered saturated cyclic amino, acyl, acylamino, acyloxy, etc. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0062] Here, as the "optionally halogenated C<sub>1-6</sub> alkyl", "optionally halogenated C<sub>3-6</sub> cycloalkyl", "optionally halogenated C<sub>1-6</sub> alkoxy", "optionally halogenated C<sub>1-6</sub> alkylthio", "C<sub>7-19</sub> aralkyl which may be substituted", "C<sub>6-14</sub> arylalkoxy which may be substituted", "C<sub>7-19</sub> aralkyloxy which may be substituted", those exemplified as the "substituents" in the above "cyclic group which may be substituted" can be used, respectively.

[0063] As the "5- to 7-membered saturated cyclic amino", those exemplified as "5- to 7-membered saturated cyclic amino" regarding "5- to 7-membered saturated cyclic amino which may be substituted" which is the "substituent" in the above "cyclic group which may be substituted" can be used.

[0064] Examples of the above "acyl" include acyl of the formulas: -CO-R<sup>3</sup>, -CO-OR<sup>3</sup>, -CO-NR<sup>3</sup>R<sup>4</sup>, -CS-NR<sup>3</sup>R<sup>4</sup>, -SO<sub>2</sub>-R<sup>3a</sup>, -PO<sub>2</sub>(OR<sup>3</sup>)-OR<sup>4</sup> or -PO<sub>2</sub>-R<sup>3a</sup> wherein R<sup>3</sup> is (i) hydrogen atom, (ii) a hydrocarbon group which may be substituted, or (iii) a heterocyclic group which may be substituted; R<sup>3a</sup> is (i) a hydrocarbon group which may be substituted, or (ii) a heterocyclic group which may be substituted; R<sup>4</sup> is hydrogen atom or C<sub>1-6</sub> alkyl; R<sup>3</sup> and R<sup>4</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted, and the like.

[0065] Examples of the "hydrocarbon group" in "hydrocarbon group which may be substituted" represented by R<sup>3</sup> or R<sup>3a</sup> include straight-chain or cyclic hydrocarbon groups (e.g. alkyl, alkenyl, alkynyl, cycloalkyl, aryl, aralkyl, etc.), etc. Among these, C<sub>1-19</sub> straight-chain or cyclic hydrocarbon groups as shown below are preferred.

- a) C<sub>1-6</sub> alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.);
- b) C<sub>2-6</sub> alkenyl (e.g., vinyl, allyl, isopropenyl, 2-butenyl, etc.);
- c) C<sub>2-6</sub> alkynyl (e.g. ethynyl, propargyl, 2-butenyl, etc.);
- d) C<sub>3-6</sub> cycloalkyl (e.g. cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, etc.); the C<sub>3-6</sub> cycloalkyl may be condensed with one benzene ring;
- e) C<sub>6-14</sub> aryl (e.g. phenyl, 1-naphthyl, 2-naphthyl, 2-indenyl, 2-anthryl, etc.), preferably phenyl;
- f) C<sub>7-19</sub> aralkyl (e.g. benzyl, phenethyl, diphenylmethyl, triphenylmethyl, 1-naphthylmethyl, 2-naphthylmethyl, 2,2-diphenylethyl, 3-phenylpropyl, 4-phenylbutyl, 5-phenylpentyl, etc.), preferably benzyl.

[0066] The "hydrocarbon groups" are preferably C<sub>1-6</sub> alkyl, C<sub>6-14</sub> aryl, C<sub>7-19</sub> aralkyl, etc.

[0067] Examples of the "substituent" in the "hydrocarbon groups which may be substituted" include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), C<sub>1-3</sub> alkyleneedioxy (e.g. methyleneedioxy, ethyleneedioxy, etc.), nitro, cyano, optionally halogenated C<sub>1-6</sub> alkoxy, optionally halogenated C<sub>1-6</sub> alkylthio, hydroxy, amino, mono-C<sub>1-6</sub> alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di-C<sub>1-6</sub> alkylamino (e.g. dimethylamino, diethylamino, dipropylamino, dibutylamino, ethylmethylamino, etc.), formyl, carboxy, carbamoyl, thiocarbamoyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl, C<sub>1-6</sub> alkoxy-carbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), 5- to 10-membered aromatic heterocyclic groups which may be substituted, C<sub>6-14</sub> aryl-carbonyl which may be substituted, C<sub>6-14</sub> arylalkoxy-carbonyl which may be substituted, C<sub>7-19</sub> aralkyloxy-carbonyl which may be substituted, 5- to 6-membered heterocyclic ring-carbonyl which may be substituted, mono-C<sub>1-6</sub> alkyl-carbamoyl (e.g. methylcarbamoyl, ethylcarbamoyl, etc.), di-C<sub>1-6</sub> alkyl-carbamoyl (e.g. dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl, etc.), C<sub>6-14</sub> aryl-carbamoyl which may be substituted, 5- to 6-membered heterocyclic ring-carbamoyl which may be substituted, optionally halogenated C<sub>1-6</sub> alkylsulfonyl, C<sub>6-14</sub> arylsulfonyl which may be substituted, formylamino, C<sub>1-6</sub> alkyl-carboxyloxy (e.g. acetoxy, propanoyloxy, etc.), C<sub>6-14</sub> aryl-carboxyloxy which may be substituted, C<sub>1-6</sub> alkoxy-carboxyloxy (e.g. methoxycarboxyloxy, ethoxycarboxyloxy, propoxycarboxyloxy, butoxycarboxyloxy,

etc.), mono- $C_{1-6}$  alkyl-carbamoyloxy (e.g. methylcarbamoyloxy, ethylcarbamoyloxy, etc.), di- $C_{1-6}$  alkyl-carbamoyloxy (e.g. dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.),  $C_{6-14}$  aryl-carbamoyloxy which may be substituted, nicotinoyloxy, etc. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0068] Here, as the "optionally halogenated  $C_{1-6}$  alkoxy", "optionally halogenated  $C_{1-6}$  alkylthio" and " $C_{6-14}$  aryl-carbamoyl which may be substituted", those exemplified as the "substituent" in the above "cyclic group which may be substituted" can be used, respectively.

[0069] As the "optionally halogenated  $C_{1-6}$  alkyl-carbonyl" and "optionally halogenated  $C_{1-6}$  alkylsulfonyle", those exemplified as the "substituent" in the above " $C_{7-19}$  aralkyl which may be substituted" can be used, respectively.

[0070] As the above "5- to 10-membered aromatic heterocyclic groups which may be substituted" and " $C_{6-14}$  aryl-carbonyl which may be substituted", those exemplified as the "substituent" in the above "5- to 7-membered saturated cyclic amino group which may be substituted" can be used, respectively.

[0071] Examples of the " $C_{6-14}$  aryloxy-carbonyl" in the " $C_{6-14}$  aryloxy-carbonyl which may be substituted" include phenyloxy-carbonyl, 1-naphthoxy-carbonyl, 2-naphthoxy-carbonyl, etc.

[0072] Examples of the " $C_{7-19}$  aralkyloxy-carbonyl" in the " $C_{7-19}$  aralkyloxy-carbonyl which may be substituted" include benzyloxy-carbonyl, phenethyloxy-carbonyl, diphenylmethyloxy-carbonyl, triphenylmethyloxy-carbonyl, 1-naphthylmethyloxy-carbonyl, 2-naphthylmethyloxy-carbonyl, 2,2-diphenylethyloxy-carbonyl, 3-phenylpropyloxy-carbonyl, 4-phenylbutyloxy-carbonyl, 5-phenylpentyloxy-carbonyl, etc.

[0073] Examples of the "5- to 6-membered heterocyclic ring-carbonyl" in the above "5- to 6-membered heterocyclic ring-carbonyl which may be substituted" include nicotinoyl, isonicotinoyl, 2-thienoyl, 3-thienoyl, 2-furyl, 3-furyl, morpholinocarbonyl, piperidinocarbonyl, pyrrolidin-1-ylcarbonyl, etc.

[0074] Examples of the "5- to 6-membered heterocyclic ring-carbamoyl" in the above "5- to 6-membered heterocyclic ring-carbamoyl which may be substituted" include morpholinocarbamoyl, piperidinocarbamoyl, 2-pyridylcarbamoyl, 3-pyridylcarbamoyl, 4-pyridylcarbamoyl, 2-thienylcarbamoyl, 3-thienylcarbamoyl, etc.

[0075] Examples of the " $C_{6-14}$  arylsulfonyl" in the above " $C_{6-14}$  arylsulfonyl which may be substituted" include phenylsulfonyl, 1-naphthylsulfonyl, 2-naphthylsulfonyl, etc.

[0076] Examples of the " $C_{6-14}$  aryl-carbonyloxy" in the above " $C_{6-14}$  aryl-carbonyloxy which may be substituted" include benzyloxy, 1-naphthoxy, 2-naphthoxy, etc.

[0077] Examples of the " $C_{6-14}$  aryl-carbamoyloxy" in the above " $C_{6-14}$  aryl-carbamoyloxy which may be substituted" include phenylcarbamoyloxy, naphthylcarbamoyloxy, etc.

[0078] As the "substituents" in the above " $C_{6-14}$  aryloxy-carbonyl which may be substituted", " $C_{7-19}$  aralkyloxy-carbonyl which may be substituted", "5- to 6-membered heterocyclic ring-carbonyl which may be substituted", "5- to 6-membered heterocyclic ring-carbamoyl which may be substituted", " $C_{6-14}$  arylsulfonyl which may be substituted", " $C_{6-14}$  aryl-carbonyloxy which may be substituted" and " $C_{6-14}$  aryl-carbamoyloxy which may be substituted", those exemplified as the "substituents" in the above " $C_{7-19}$  aralkyl which may be substituted" can be mentioned. The number of the substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0079] Examples of the "heterocyclic groups" in the "heterocyclic groups which may be substituted" represented by  $R^3$  or  $R^{3a}$  include univalent groups formed by removing an optional one hydrogen atom from a 5- to 14-membered (monocyclic, bicyclic or tricyclic) heterocyclic ring containing 1 or 2 kinds of, 1 to 4 hetero atoms selected from nitrogen, sulfur and oxygen atom in addition to carbon atoms, preferably, (i) an aromatic heterocyclic ring, (ii) a 5- to 10-membered non-aromatic heterocyclic ring, or (iii) a 7- to 10-membered heterocyclic-bridge ring.

[0080] Here, examples of the "aromatic heterocyclic ring" include a 5- to 14-membered, preferably 5- to 10-membered, aromatic heterocyclic ring containing one or more hetero atoms (e.g. 1 to 4) selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms. Specific examples include aromatic heterocyclic rings such as thiophene, furan, pyrrole, imidazole, pyrazole, thiazole, isothiazole, oxazole, isoxazole, pyridine, pyrazine, pyrimidine, pyridazine, 1,2,4-oxadiazole, 1,3,4-oxadiazole, 1,2,4-thiadiazole, 1,3,4-thiadiazole, furazan, benzothiofene, benzofuran, benzimidazole, benzoxazole, benzothiazole, benzisothiazole, naphtho[2,3-b]thiophene, phenoxathiin, indole, isoindole, 1H-indazole, purine, 4H-quinoline, isouquinoline, quinoline, phenalazine, naphthyridine, quinoxaline, quinoxaline, cinnoline, carbazole,  $\beta$ -carboline, phenanthridine, acridine, phenazine, phenothiazine, phenoxazine, phthalimide, etc.; or a ring formed by condensing these rings (preferably monocyclic rings) with one to multiple (preferably 1 or 2) aromatic rings (e.g. benzene ring, etc.), etc.

[0081] Examples of "5- to 10-membered non-aromatic heterocyclic rings" include 2- or 3-pyrroline, pyrrolidine, 2- or 3-imidazoline, 2-oxazoline, oxazolidine, 2- or 3-pyrazoline, pyrazolidine, 2-thiazoline, piperidine, piperazine, hexamethylenimine, morpholine, thiomorpholine, etc.

[0082] Examples of "7- to 10-membered heterocyclic-bridge rings" include quinuclidine, 7-azabicyclo[2.2.1]heptane, etc.

[0083] The "heterocyclic groups" are preferably 5- to 10-membered (monocyclic or bicyclic) heterocyclic groups con-

taining 1 or 2 kinds of, preferably 1 to 4, hetero atoms selected from nitrogen, sulfur and oxygen atoms in addition to carbon atoms. Specific examples include aromatic heterocyclic groups such as 2- or 3-thienyl; 2-, 3- or 4-pyridyl; 2- or 3-furyl; 2-, 4- or 5-thiazolyl; 2-, 4- or 5-oxazolyl; 1-, 3- or 4-pyrazolyl; 2-pyrazinyl; 2-, 4- or 5-pyrimidinyl; 1-, 2- or 3-pyrrolyl; 1-, 2- or 4-imidazolyl; 3- or 4-pyridazinyl; 3-isothiazolyl; 3-isoxazolyl; 1,2,4-oxadiazol-5-yl; 1,2,4-oxadiazol-3-yl; 2-, 3-, 4-, 5- or 8-quinolyl; 1-, 3-, 4-, 5-, 6-, 7- or 8-isquinolyl; 1-, 2-, 3-, 4-, 5-, 6- or 7-indolyl; 1-, 2-, 4- or 5-isolindolyl; 1-, 5- or 6-phthalazinyl; 2-, 3- or 5-quinoxalyl; 2-, 3-, 4-, 5- or 6-benzofuranyl; 2-, 3-, 4-, 5- or 6-benzothienyl; 2-, 4-, 5- or 6-benzothiazolyl; 1-, 2-, 4-, 5- or 6-benzimidazolyl; etc.; and non-aromatic heterocyclic groups such as 1-, 2- or 3-pyrrolidinyl; 1-, 2-, 4- or 5-imidazolidinyl; 2- or 4-imidazolyl; 2-, 3- or 4-pyrazolidinyl; piperidino; 2-, 3- or 4-piperidyl; 1- or 2-piperazinyl; morpholino; etc.

[0084] As the "substituents" in the "heterocyclic groups which may be substituted", those exemplified as the "substituents" in the above "5- to 10-membered aromatic heterocyclic groups which may be substituted" can be used. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0085] Examples of the "C<sub>1-6</sub> alkyl" represented by R<sup>4</sup> include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc.

[0086] Examples of the "nitrogen-containing heterocyclic ring" in the "nitrogen-containing heterocyclic ring which may be substituted" formed by R<sup>3</sup> and R<sup>4</sup> together with the adjacent nitrogen atom include a 5- to 7-membered nitrogen-containing heterocyclic ring which contains at least one nitrogen atom in addition to carbon atoms and may contain 1 to 3 hetero atoms selected from nitrogen, sulfur and oxygen atoms. The "nitrogen-containing heterocyclic rings" are preferably piperidine, morpholine, thiomorpholine, piperazine, pyrrolidine, etc.

[0087] As the "substituents" in the "nitrogen-containing heterocyclic ring which may be substituted", those exemplified as the "substituents" in the above "5- to 10-membered aromatic heterocyclic groups which may be substituted" can be used. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0088] The "acyl" is preferably formyl, carboxy, carbamoyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl (e.g. acetyl, etc.), C<sub>1-6</sub> alkoxy-carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), C<sub>6-14</sub> aryl-carbonyl which may be substituted (e.g. benzoyl, 1-naphthoyl, 2-naphthoyl, etc.), C<sub>6-14</sub> aryloxy-carbonyl which may be substituted (e.g. phenyloxycarbonyl, 1-naphthyloxycarbonyl, 2-naphthyloxycarbonyl, etc.), C<sub>7-19</sub> aralkyloxy-carbonyl which may be substituted (e.g. benzyloxycarbonyl, phenethyloxycarbonyl, etc.), a 5- to 6-membered heterocyclic ring-carbonyl which may be substituted (e.g. nicotinyl, etc.), mono-C<sub>1-6</sub> alkyl-carbamoyl (e.g. methylcarbamoyl, ethylcarbamoyl, etc.), di-C<sub>1-6</sub> alkyl-carbamoyl (e.g. dimethylcarbamoyl, diethylcarbamoyl, ethyldimethylcarbamoyl, etc.), C<sub>6-14</sub> aryl-carbamoyl which may be substituted (e.g. phenylcarbamoyl, 4-methoxyphenylcarbamoyl, 3,4-dimethoxyphenylcarbamoyl, etc.), aromatic heterocyclic ring-carbamoyl which may be substituted (e.g. 2-pyridinylcarbamoyl, 2-quinolinylcarbamoyl, etc.), optionally halogenated C<sub>1-6</sub> alkylsulfonyl (e.g. methylsulfonyl, etc.), C<sub>6-14</sub> arylsulfonyl which may be substituted (e.g. phenylsulfonyl, etc.), etc.

[0089] Here, as the "optionally halogenated C<sub>1-6</sub> alkyl-carbonyl" and "optionally halogenated C<sub>1-6</sub> alkylsulfonyl", those exemplified as the "substituents" in the above "C<sub>7-19</sub> aralkyl which may be substituted" can be used, respectively.

[0090] As the "C<sub>6-14</sub> aryl-carbonyl which may be substituted", those exemplified as the "substituents" in the above "5- to 7-membered saturated cyclic amino which may be substituted" can be used.

[0091] As the "C<sub>6-14</sub> aryloxy-carbonyl which may be substituted", "C<sub>7-19</sub> aralkyloxy-carbonyl which may be substituted", "5- to 6-membered heterocyclic ring-carbonyl which may be substituted", "aromatic heterocyclic ring-carbamoyl which may be substituted" and "C<sub>6-14</sub> arylsulfonyl which may be substituted", those exemplified as the "substituents" in the above "hydrocarbon groups which may be substituted" can be used, respectively.

[0092] As the "C<sub>6-14</sub> aryl-carbamoyl which may be substituted", those exemplified as the "substituents" in the above "cyclic group which may be substituted" can be used.

[0093] Examples of the above "acylamino" include amino which is substituted by 1 or 2 of the above "acyl". Preferably, acylamino of the formulas: -NR<sup>5</sup>-COR<sup>6</sup>, -NR<sup>5</sup>-COOR<sup>6a</sup>, -NR<sup>5</sup>-SO<sub>2</sub>-R<sup>6a</sup>, -NR<sup>5</sup>-CONR<sup>6a</sup>-R<sup>6b</sup>, -PO(-OR<sup>5</sup>)-OR<sup>6</sup> or -PO<sub>2</sub>-R<sup>6</sup> wherein R<sup>5</sup> is hydrogen atom or C<sub>1-6</sub> alkyl; R<sup>6</sup> has the same meaning as the above R<sup>3</sup>; R<sup>6a</sup> has the same meaning as the above R<sup>3a</sup>; and R<sup>6b</sup> has the same meaning as R<sup>4</sup>, etc., can be mentioned.

[0094] As the "C<sub>1-6</sub> alkyl" represented by R<sup>5</sup>, the same one as the "C<sub>1-6</sub> alkyl" for the above R<sup>4</sup> can be mentioned.

[0095] The "acylamino" is preferably formylamino, optionally halogenated C<sub>1-6</sub> alkyl-carboxamide (e.g. methylcarboxamide, trifluoromethylcarboxamide, etc.), C<sub>6-14</sub> aryl-carboxamide which may be substituted (e.g. phenylcarboxamide, 2-methoxyphenylcarboxamide, 4-methoxyphenylcarboxamide, etc.), N-(C<sub>6-14</sub> aryl-carbonyl which may be substituted)-N-C<sub>1-6</sub> alkylamino (e.g. N-4-methoxybenzoyl-N-methylamino, etc.), C<sub>7-19</sub> aralkyl-carboxamide which may be substituted (e.g. benzyloxycarbonyl, etc.), aromatic heterocyclic ring-carboxamide which may be substituted (e.g. benzothienophen-2-ylcarboxamide, etc.), optionally halogenated C<sub>1-6</sub> alkoxy-carboxamide (e.g. methoxycarboxamide, ethoxycarboxamide, propoxycarboxamide, butoxycarboxamide, etc.), C<sub>6-14</sub> arylaminocarboxylamino which may be substituted (e.g. phenylaminocarboxylamino, etc.), optionally halogenated C<sub>1-6</sub> alkylsulfonylamino (e.g. methylsulfo-

nylamino, trifluoromethylsulfonylamino, ethylsulfonylamino, etc.),  $C_{6-14}$  arylsulfonylamino which may be substituted (e.g. 4-methoxyphenylsulfonylamino, etc.), etc.

[0096] Here, as the "substituents" in the " $C_{6-14}$  aryl-carboxamide which may be substituted", " $N-(C_{6-14}$  aryl-carbonyl which may be substituted)- $N-C_{1-6}$  arylkylamino", " $C_{7-19}$  aralkyl-carboxamide which may be substituted", "aromatic heterocyclic ring-carboxamide which may be substituted", " $C_{6-14}$  arylaminocarbonylamino which may be substituted" and " $C_{6-14}$  arylsulfonylamino which may be substituted", those exemplified as the "substituents" in the above " $C_{7-19}$  aralkyl which may be substituted" can be mentioned. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0097] Examples of the above "acyloxy" include oxy substituted by one of the above "acyl". Preferably, acyloxy of the formulas:  $-O-COR^7$ ,  $-O-COOR^7$ ,  $-O-CONHR^7$ ,  $-PO(OH)-OR^7$  or  $-PO_2-R^7$  wherein  $R^7$  has the same meaning as the above  $R^5$ , etc., can be mentioned.

[0098] The "acyloxy" is preferably optionally halogenated  $C_{1-6}$  alkyl-carbonyloxy (e.g. acetoxy, propanoyloxy, etc.),  $C_{6-14}$  aryl-carbonyloxy which may be substituted (e.g. benzyloxy, 4-methoxybenzyloxy, etc.), optionally halogenated  $C_{1-6}$  alkoxy-carbonyloxy (e.g. methoxycarbonyloxy, trifluoromethoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, butoxycarbonyloxy, etc.), mono- $C_{1-6}$  alkyl-carbamoyloxy (e.g. methylcarbamoyloxy, ethylcarbamoyloxy, etc.), di- $C_{1-6}$  alkyl-carbamoyloxy (e.g. dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.),  $C_{6-14}$  aryl-carbamoyloxy which may be substituted (e.g. phenylcarbamoyloxy, naphthylcarbamoyloxy, etc.), nicotinoyloxy, etc.

[0099] As the "substituents" in " $C_{6-14}$  aryl-carbonyloxy which may be substituted" and " $C_{6-14}$  aryl-carbamoyloxy which may be substituted", those exemplified as the "substituents" in the above " $C_{7-19}$  aralkyl which may be substituted" can be mentioned. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0100] Examples of the "5- to 7-membered non-aromatic heterocyclic groups which may be substituted", which is the "substituents" in "cyclic group which may be substituted" represented by  $Ar^1$ , include 4,5-dihydro-1,3-oxazol-2-yl, 4,5-dihydro-1,3-thiazol-2-yl, 4,5-dihydro-1H-2-imidazolyl, etc. As the "substituents" in the "5- to 7-membered non-aromatic heterocyclic groups which may be substituted", those exemplified as the "substituents" in the above "5- to 7-membered saturated cyclic amino group which may be substituted" can be used.

[0101] As the "acyl", "acyloxy" and "acylamino", which are the "substituents" in the "cyclic group which may be substituted" represented by  $Ar^1$ , those exemplified as the "substituents" in the above "5- to 10-membered aromatic heterocyclic groups which may be substituted" can be used.

[0102] The "substituents" in the "cyclic group which may be substituted" for  $Ar^1$  are preferably halogen atom (preferably fluorine, chlorine and bromine, etc.); nitro;  $C_{1-3}$  alkylendioxy (preferably methylenedioxy, etc.); optionally halogenated  $C_{1-6}$  alkyl (preferably, methyl, ethyl, propyl, trifluoromethyl, etc.); hydroxy- $C_{1-6}$  alkyl (preferably hydroxymethyl, etc.); optionally halogenated  $C_{3-6}$  cycloalkyl (preferably cyclohexyl, etc.); optionally halogenated  $C_{1-6}$  alkoxy (preferably methoxy, ethoxy, etc.); optionally halogenated  $C_{1-6}$  alkylthio (preferably methylthio, etc.); hydroxy- $C_{7-19}$  aralkyloxy which may be substituted (preferably benzyloxy, 4-methoxybenzyloxy, 3-methoxybenzyloxy, 4-fluorobenzyloxy, 4-methylthiobenzyloxy, 4-ethylbenzyloxy, etc.);  $C_{6-14}$  aryloxy which may be substituted (preferably phenyloxy, etc.); amino; mono- $C_{1-6}$  alkylamino (preferably methylamino, etc.); di- $C_{1-6}$  alkylamino (preferably dimethylamino, etc.); 5- to 7-membered saturated cyclic amino which may be substituted and may be condensed with a benzene ring (preferably 1,3-dioxo-1,3-dihydro-2H-isoindol-2-yl, methylpiperidino, oxopiperidino, etc.); 5- to 7-membered non-aromatic heterocyclic groups which may be substituted (preferably 4,5-dihydro-1,3-oxazol-2-yl, etc.); formyl; carboxy;  $C_{6-14}$  aryl-carbonyl which may be substituted (preferably benzoyl, etc.);  $C_{6-14}$  aryl-carbamoyl which may be substituted (preferably, phenylcarbamoyl, 4-methoxyphenylcarbamoyl, 3,4-dimethoxyphenylcarbamoyl, etc.); aromatic heterocyclic ring-carbamoyl which may be substituted (preferably 2-pyridinylcarbamoyl, 2-quinolinylcarbamoyl, etc.);  $C_{1-6}$  alkoxy-carbonyl (preferably methoxycarbonyl, ethoxycarbonyl, etc.); optionally halogenated  $C_{1-6}$  alkyl-carboxamide (preferably methylcarboxamide, trifluoromethylcarboxamide, etc.);  $C_{6-14}$  aryl-carboxamide which may be substituted (preferably phenylcarboxamide, 2-methoxyphenylcarboxamide, 4-methoxyphenylcarboxamide, etc.);  $C_{7-19}$  aralkyl-carboxamide which may be substituted (preferably benzylcarboxamide, etc.); aromatic heterocyclic ring-carboxamide which may be substituted (preferably benzothienophen-2-ylcarboxamide, etc.);  $N-(C_{6-14}$  aryl-carbonyl which may be substituted)- $N-C_{1-6}$  alkylamino (preferably  $N-4$ -methoxybenzoyl- $N$ -methylamino, etc.);  $C_{6-14}$  arylamino-carbonylamino which may be substituted (preferably phenylaminocarbonylamino, etc.);  $C_{6-14}$  arylsulfonylamino which may be substituted (preferably 4-methoxyphenylsulfonylamino, etc.);  $C_{6-14}$  aryl-carbonyloxy which may be substituted (preferably 4-methoxybenzyloxy, etc.); oxo; etc.

[0103] When the "cyclic group" in the "cyclic group which may be substituted" represented by  $Ar^1$  is a non-aromatic cyclic hydrocarbon group or a non-aromatic heterocyclic group,  $C_{6-14}$  aryl which may be substituted (preferably phenyl, 4-fluorophenyl, chlorophenyl, methylphenyl, methoxyphenyl, etc.) can be used as a preferred substituent.

[0104]  $Ar^1$  is preferably phenyl, biphenyl (preferably 4-biphenyl), phenyl-pyridyl (preferably 6-phenyl-3-pyridyl, 5-phenyl-2-pyridyl), phenyl-furyl (preferably 5-phenyl-2-furyl), phenyl-isoxazolyl (preferably 3-phenyl-isoxazol-5-yl), diphenyl-oxazolyl (preferably 2,4-diphenyl-1,3-oxazol-5-yl), pyridyl-phenyl (preferably 4-(4-pyridyl)phenyl), phenyl-py-

rimidinyl (preferably 2-phenyl-5-pyrimidinyl), benzofuranyl-phenyl (preferably 4-(2-benzofuranyl)phenyl), or furyl-phenyl (preferably 4-(2-furyl)phenyl); each of which may have 1 to 3 (preferably 1 or 2) substituents selected from the group consisting of halogen atom (preferably fluorine, chlorine, bromine, etc.); nitro; C<sub>1-6</sub> alkylenedioxy (preferably methylenedioxy, etc.); optionally halogenated C<sub>1-6</sub> alkyl (preferably methyl, ethyl, propyl, trifluoromethyl, etc.); hydroxy-C<sub>1-6</sub> alkyl (preferably hydroxymethyl, etc.); optionally halogenated C<sub>3-6</sub> cycloalkyl (preferably cyclohexyl, etc.); optionally halogenated C<sub>1-6</sub> alkoxy (preferably methoxy, ethoxy, etc.); optionally halogenated C<sub>1-6</sub> alkythio (preferably methylthio, etc.); hydroxy; C<sub>7-19</sub> aralkyloxy which may be substituted (preferably benzoyloxy, 4-methoxybenzoyloxy, 3-methoxybenzoyloxy, 4-fluorobenzyloxy, 4-methylthiobenzyloxy, 4-ethylbenzyloxy, etc.); C<sub>6-14</sub> aryloxy which may be substituted (preferably phenyloxy, etc.); amino; mono-C<sub>1-6</sub> alkylamino (preferably methylamino, etc.); di-C<sub>1-6</sub> alkylamino (preferably dimethylamino, etc.); 5- to 7-membered saturated cyclic amino which may be substituted and may be condensed with a benzene ring (preferably 1,3-dioxo-1,3-dihydro-2H-isindol-2-yl, methylpiperidino, oxopiperidino, etc.); 5- to 7-membered non-aromatic heterocyclic groups which may be substituted (preferably 4,5-dihydro-1,3-oxazol-2-yl, etc.); formyl; carboxy; C<sub>6-14</sub> aryl-carbonyl which may be substituted (preferably benzoyl, etc.); C<sub>6-14</sub> aryl-carbamoyl which may be substituted (preferably phenylcarbamoyl, 4-methoxyphenylcarbamoyl, 3,4-dimethoxyphenylcarbamoyl, etc.); aromatic heterocyclic ring-carbamoyl which may be substituted (e.g. 2-pyridinylcarbamoyl, 2-quinolinylcarbamoyl, etc.); C<sub>1-6</sub> alkoxy-carbonyl (preferably methoxycarbonyl, ethoxycarbonyl, etc.); optionally halogenated C<sub>1-6</sub> alkyl-carboxamide (preferably, methylcarboxamide, trifluoromethylcarboxamide, etc.); C<sub>6-14</sub> aryl-carboxamide which may be substituted (preferably phenylcarboxamide, 2-methoxyphenylcarboxamide, 4-methoxyphenylcarboxamide, etc.); C<sub>7-19</sub> aralkyl-carboxamide which may be substituted (preferably benzylcarboxamide, etc.); aromatic heterocyclic ring-carboxamide which may be substituted (preferably benzothioiphen-2-ylcarboxamide, etc.); N-(C<sub>6-14</sub> aryl)-carbonyl which may be substituted (preferably N-C<sub>1-6</sub> alkylamino (preferably N-4-methoxybenzoyl-N-methylamino, etc.); C<sub>6-14</sub> arylamino-carbonylamino which may be substituted (preferably phenylaminocarbonylamino, etc.); C<sub>6-14</sub> arylsulfonylamino which may be substituted (preferably 4-methoxyphenylsulfonylamino, etc.); C<sub>6-14</sub> aryl-carbonyloxy which may be substituted (preferably 4-methoxybenzyloxy, etc.); oxo; etc.

[0105] Further, preferred examples of Ar<sup>1</sup> include piperidinyl (piperidino), piperazinyl, pyrrolidinyl, 1,3-dioxanyl, etc.; each of which may have 1 or 2 substituents selected from the group consisting of oxo and C<sub>6-14</sub> aryl which may be substituted (preferably phenyl, fluorophenyl, chlorophenyl, methylphenyl, methoxyphenyl).

[0106] Ar<sup>1</sup> is more preferably,

(1) phenyl, biphenyl (preferably 4-biphenyl) or phenyl-pyridyl (preferably 6-phenyl-3-pyridyl, 5-phenyl-2-pyridyl); each of which may have 1 to 3 substituents selected from the group consisting of halogen atom (preferably fluorine, chlorine, bromine, etc.); optionally halogenated C<sub>1-6</sub> alkyl (preferably methyl, ethyl, propyl, trifluoromethyl, etc.); and optionally halogenated C<sub>1-6</sub> alkoxy (preferably methoxy, ethoxy, etc.); or

(2) piperidinyl (piperidino) which may have 1 or 2 substituents selected from C<sub>6-14</sub> aryl (preferably phenyl, fluorophenyl, chlorophenyl, methylphenyl, methoxyphenyl) which may be substituted (preferably by 1 to 3 substituents selected from the group consisting of halogen atom (preferably fluorine, chlorine, bromine, etc.), optionally halogenated C<sub>1-6</sub> alkyl (preferably methyl, etc.) and optionally halogenated C<sub>1-6</sub> alkoxy (preferably methoxy, etc.)).

[0107] The "spacer having a main chain of 1 to 6 atoms" represented by X and Y means a space in which 1 to 6 atoms are linked. Here, the "number of atoms in the main chain" is counted so that the number of atoms in the main chain is minimum. For example, the number of atoms of 1,2-cyclopentylene is counted as 2, and the number of atoms of 1,3-cyclopentylene is counted as 3.

[0108] Examples of the "spacer having a main chain of 1 to 6 atoms" include a bivalent group consisting of 1 to 3 species selected from -O-, -S-, -CO-, -SO-, -SO<sub>2</sub>-, -NR<sup>9</sup> (R<sup>9</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl, optionally halogenated C<sub>1-6</sub> alkylsulfonyl), optionally halogenated bivalent C<sub>1-6</sub> non-cyclic hydrocarbon groups, and bivalent C<sub>6-8</sub> monocyclic non-aromatic hydrocarbon groups, and the like.

[0109] Here, as the "optionally halogenated C<sub>1-6</sub> alkyl", those exemplified as the "substituents" in the above "cyclic group which may be substituted" can be used.

[0110] As the "optionally halogenated C<sub>1-6</sub> alkyl-carbonyl" and "optionally halogenated C<sub>1-6</sub> alkylsulfonyl", those exemplified as the "substituents" in the above "C<sub>7-19</sub> aralkyl which may be substituted" can be used, respectively.

[0111] Examples of the "bivalent C<sub>1-6</sub> non-cyclic hydrocarbon groups" in the "optionally halogenated bivalent C<sub>1-6</sub> non-cyclic hydrocarbon groups" include

(1) C<sub>1-6</sub> alkylene (e.g. -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -(CH<sub>2</sub>)<sub>5</sub>-, -(CH<sub>2</sub>)<sub>6</sub>-, -CH(CH<sub>3</sub>)-, -C(CH<sub>3</sub>)<sub>2</sub>-, -CH(CF<sub>3</sub>)-, -(CH(CH<sub>3</sub>))<sub>2</sub>-, -(CF<sub>3</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>-, -(CH<sub>3</sub>)<sub>3</sub>C(CH<sub>3</sub>)<sub>2</sub>-, etc.);

(2) C<sub>2-6</sub> alkenylene (e.g. -CH=CH-, -CH<sub>2</sub>CH=CH-, -CH<sub>2</sub>CF=CH-, -C(CH<sub>3</sub>)<sub>2</sub>CH=CH-, -CH<sub>2</sub>CH=CH-CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>CH=CH-, -CH=CH-CH=CH-, -CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, etc.);

(3) C<sub>2-6</sub> alkynylene (e.g. -C≡C-, -CH<sub>2</sub>C≡C-, -CH<sub>2</sub>C≡C-CH<sub>2</sub>-CH<sub>2</sub>-, etc.), etc.,

each of which may have 1 to 5, preferably 1 to 3, halogen atoms (e.g. fluorine, chlorine, bromine, iodine, etc.).

[0112] As the "bivalent  $C_{5-8}$  monocyclic non-aromatic hydrocarbon groups", for example, bivalent groups formed by removing optional two hydrogen atoms from  $C_{5-8}$  cycloalkane or  $C_{5-8}$  cycloalkene, can be mentioned. Specific examples include 1,2-cyclopentylene; 1,3-cyclopentylene; 1,2-cyclohexylene; 1,3-cyclohexylene; 1,4-cyclohexylene; 1,2-cycloheptylene; 1,3-cycloheptylene; 1,4-cycloheptylene; 3-cyclohexen-1,4-ylene; 3-cyclohexen-1,2-ylene; 2,5-cyclohexadien-1,4-ylene, etc. Especially,  $C_{5-8}$  cycloalkylene is preferable.

[0113] The "spacer having a main chain of 1 to 6 atoms" represented by X and Y is preferably a bivalent group consisting of 1 to 3 species selected from -O-, -S-, -CO-, -SO-, -SO<sub>2</sub>-, -NR<sup>R</sup> (R<sup>R</sup> is as defined above) and optionally halogenated bivalent  $C_{1-6}$  non-cyclic hydrocarbon groups.

[0114] Preferred examples of the "spacer having a main chain of 1 to 6 atoms" include

- (1)  $C_{1-6}$  alkylene (e.g. -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, -(CH<sub>2</sub>)<sub>4</sub>-, -(CH<sub>2</sub>)<sub>5</sub>-, -(CH<sub>2</sub>)<sub>6</sub>-, -CHCH<sub>3</sub>-, -(CH(CH<sub>3</sub>))<sub>2</sub>-, -(CF<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>C(CH<sub>3</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>C(CH<sub>3</sub>)<sub>2</sub>-, etc.);
- (2)  $C_{2-6}$  alkenylene (e.g. -CH=CH-, -CH<sub>2</sub>-CH=CH-, -C(CH<sub>3</sub>)<sub>2</sub>-CH=CH-, -CH<sub>2</sub>-CH=CH-CH<sub>2</sub>-, -CH<sub>2</sub>-CH<sub>2</sub>-CH=CH-, -CH=CH-CH=CH-, -CH=CH-CH<sub>2</sub>-CH<sub>2</sub>-CH<sub>2</sub>-, etc.);
- (3)  $C_{2-6}$  alkynylene (e.g. -C≡C-, -CH<sub>2</sub>-C≡C-, -CH<sub>2</sub>-C≡C-CH<sub>2</sub>-CH<sub>2</sub>-, etc.);
- (4) -(CH<sub>2</sub>)<sub>w1</sub>-(O(CH<sub>2</sub>)<sub>w2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>w1</sub>-S(CH<sub>2</sub>)<sub>w2</sub>-, -(CH<sub>2</sub>)<sub>w1</sub>-CO(CH<sub>2</sub>)<sub>w2</sub>-, -(CH<sub>2</sub>)<sub>w1</sub>-SO(CH<sub>2</sub>)<sub>w2</sub>-, -(CH<sub>2</sub>)<sub>w1</sub>-SO<sub>2</sub>(CH<sub>2</sub>)<sub>w2</sub>-, -(CH<sub>2</sub>)<sub>w1</sub>NR<sup>R</sup>(CH<sub>2</sub>)<sub>w2</sub>-,
- (5) -(CH<sub>2</sub>)<sub>w3</sub>CONR<sup>R</sup>(CH<sub>2</sub>)<sub>w4</sub>-, -(CH<sub>2</sub>)<sub>w3</sub>NR<sup>R</sup>CO(CH<sub>2</sub>)<sub>w4</sub>-, -(CH<sub>2</sub>)<sub>w3</sub>SO<sub>2</sub>NR<sup>R</sup>(CH<sub>2</sub>)<sub>w4</sub>-, -(CH<sub>2</sub>)<sub>w3</sub>NR<sup>R</sup>SO<sub>2</sub>(CH<sub>2</sub>)<sub>w4</sub>-,
- (6) -(CH<sub>2</sub>)<sub>w5</sub>NR<sup>R</sup>CONR<sup>R</sup>(CH<sub>2</sub>)<sub>w6</sub>-,

wherein R<sup>R</sup> is as defined above; R<sup>Bb</sup> has the same meaning as R<sup>B</sup>; w1 and w2 is an integer of 0 to 5, and w1 + w2 is 0 to 5; w3 and w4 is an integer of 0 to 4, and w3 + w4 is 0 to 4; w5 and w6 is an integer of 0 to 3, and w5 + w6 is 0 to 3, etc.

[0115] More preferably, the "spacer having a main chain of 1 to 6 atoms" represented by X is -(CH<sub>2</sub>)<sub>w1</sub>-(O(CH<sub>2</sub>)<sub>w2</sub>)<sub>2</sub>-, -CONR<sup>R</sup>-, -NR<sup>R</sup>CO- (wherein the symbols are as defined above), -OCO-, -COO-, etc. Among these, -CONH-, -NHCO-, -COO-, etc. are preferred. In particular, -CONH- or -COO- are preferred.

[0116] More preferably, the "spacer having a main chain of 1 to 6 atoms" represented by Y is  $C_{1-3}$  alkylene (e.g., -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, etc.), -(CH<sub>2</sub>)<sub>w3</sub>CONH(CH<sub>2</sub>)<sub>w4</sub>-, -(CH<sub>2</sub>)<sub>w3</sub>COO(CH<sub>2</sub>)<sub>w4</sub> (wherein the symbols are as defined above), etc. In particular,  $C_{1-3}$  alkylene (e.g. -CH<sub>2</sub>-, -(CH<sub>2</sub>)<sub>2</sub>-, -(CH<sub>2</sub>)<sub>3</sub>-, etc.), etc., are preferred.

[0117] As the "condensed polycyclic aromatic rings" in "condensed polycyclic aromatic rings which may be substituted" represented by Ar, those exemplified as the "cyclic group" in the "cyclic group which may be substituted" represented by the above Ar<sup>1</sup> can be used.

[0118] The "condensed polycyclic aromatic rings" are preferably  $C_{9-14}$  condensed polycyclic (bicyclic or tricyclic) aromatic hydrocarbons, or 10-membered condensed polycyclic aromatic heterocyclic rings.

[0119] More preferably, the "condensed polycyclic aromatic rings" are naphthalene, isoquinoline, quinoline, quinoxaline, phthalazine, naphthyridine, quinoxaline, cinnoline, indole, etc. In particular, naphthalene, quinoline, etc. are preferred.

[0120] As the "substituents" in the "condensed polycyclic aromatic rings which may be substituted" represented by Ar, those exemplified as the "substituents" in the "cyclic group which may be substituted" represented by the above Ar<sup>1</sup> can be used.

[0121] The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0122] Ar is especially preferably quinoline or naphthalene.

[0123] As the "hydrocarbon groups which may be substituted" represented by R<sup>1</sup> and R<sup>2</sup>, those exemplified as the above R<sup>3</sup> can be used.

[0124] The "hydrocarbon groups which may be substituted" are preferably  $C_{1-6}$  alkyl which may be substituted" or phenyl.

[0125] Here, examples of the " $C_{1-6}$  alkyl" in the " $C_{1-6}$  alkyl which may be substituted" include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, etc. Especially, methyl, ethyl, propyl, etc. are preferred.

[0126] Examples of the "substituents" in the " $C_{1-6}$  alkyl which may be substituted" include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.),  $C_{1-3}$  alkylenedioxy (e.g. methylenedioxy, ethylenedioxy etc.), nitro, cyano, optionally halogenated  $C_{3-6}$  cycloalkyl, optionally halogenated  $C_{1-6}$  alkoxy, optionally halogenated  $C_{1-6}$  alkylthio, hydroxy, amino, mono- $C_{1-6}$  alkylamino (e.g. methylamino, ethylamino, propylamino, isopropylamino, butylamino, etc.), di- $C_{1-6}$  alkylamino (e.g. dimethylamino, diethylamino, dipropylamino, dibutylamino, ethylmethylamino, etc.), formyl, carboxy, carbamoyl, thiocarbamoyl, optionally halogenated  $C_{1-6}$  alkyl-carbonyl,  $C_{1-6}$  alkoxy-carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, tert-butoxycarbonyl, etc.), mono- $C_{1-6}$  alkyl-carbamoyl (e.g. methylcarbamoyl, ethylcarbamoyl, etc.), di- $C_{1-6}$  alkyl-carbamoyl (e.g. dimethylcarbamoyl, diethylcarbamoyl, ethylmethylcarbamoyl etc.), option-



ally halogenated  $C_{1-6}$  alkylsulfonfyl, formylamino, optionally halogenated  $C_{1-6}$  alkyl-carboxamide,  $C_{1-6}$  alkoxy-carboxamide (e.g. methoxycarboxamide, ethoxycarboxamide, propoxycarboxamide, butoxycarboxamide, etc.),  $C_{1-6}$  alkylsulfonfylamino (e.g. methylsulfonfylamino, ethylsulfonfylamino, etc.),  $C_{1-6}$  alkyl-carbonyloxy (e.g. acetoxy, propanoyloxy, etc.),  $C_{1-6}$  alkoxy-carbonyloxy (e.g. methoxycarbonyloxy, ethoxycarbonyloxy, propoxycarbonyloxy, butoxycarbonyloxy, etc.), mono- $C_{1-6}$  alkyl-carbamoyloxy (e.g. methylcarbamoyloxy, ethylcarbamoyloxy, etc.), di- $C_{1-6}$  alkyl-carbamoyloxy (e.g. dimethylcarbamoyloxy, diethylcarbamoyloxy, etc.), aromatic groups which may be substituted, etc. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0127] Here, as the "optionally halogenated  $C_{3-6}$  cycloalkyl", "optionally halogenated  $C_{1-6}$  alkoxy" and "optionally halogenated  $C_{1-6}$  alkylthio", those exemplified as the "substituents" in the above "cyclic group which may be substituted" can be used.

[0128] As the "optionally halogenated  $C_{1-6}$  alkyl-carbonyl", "optionally halogenated  $C_{1-6}$  alkylsulfonfyl" and "optionally halogenated  $C_{1-6}$  alkyl-carboxamide", those exemplified as the "substituents" in the above " $C_{7-19}$  aralkyl which may be substituted" can be used.

[0129] As the "substituents" and "aromatic groups" in the "aromatic groups which may be substituted", those exemplified as the "substituents" and "aromatic groups" in the "cyclic group which may be substituted" represented by the above  $Ar^1$  can be used. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0130] More preferably, "hydrocarbon group which may be substituted" represented by  $R^1$  and  $R^2$  is  $C_{1-8}$  alkyl. In particular, methyl, ethyl, isopropyl, etc., are preferred.

[0131] Examples of the "nitrogen-containing heterocyclic rings" in the "nitrogen-containing heterocyclic rings which may be substituted" formed by  $R^1$  and  $R^2$  together with the adjacent nitrogen atom include 3- to 10-membered (preferably 3- to 8-membered) nitrogen-containing heterocyclic rings which contain at least one nitrogen atom in addition to carbon atoms, and which may further contain 1 to 3 hetero atoms selected from nitrogen, sulfur and oxygen atoms. Specific examples include aziridine, azetidine, morpholine, thiomorpholine, piperidine, piperazine, pyrrolidine, hexamethylenimine, heptamethylenimine, hexahydropyrimidine, 1,4-diazepan, thiazolidine, imidazolidine, heptahydroindole, decahydroquinoline, decahydroisoquinoline, and their unsaturated cyclic amines (e.g. 1,2,5,6-tetrahydroindole, 1H-imidazole, 4,5-dihydro-1H-imidazole, 2,3-dihydroindole, 1,2,3,4-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, etc.), etc. can be mentioned. Especially, morpholine, piperidine, piperazine, pyrrolidine, 1,3-thiazolidine, 1H-imidazole, 4,5-dihydro-1H-imidazole, 2,3-dihydroindole, 1,2,3,4-tetrahydroquinoline, 1,2,3,4-tetrahydroisoquinoline, etc., are preferred. In particular, morpholine, piperidine, piperazine, pyrrolidine, etc. are preferred.

[0132] As the "substituents" in the "nitrogen-containing heterocyclic rings which may be substituted", for example, those exemplified as the "substituents" in the above "5- to 7-membered saturated cyclic amino group which may be substituted" can be used. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituents can be the same or different.

[0133] The substituents are preferably optionally halogenated  $C_{1-6}$  alkyl (preferably methyl); 5- to 10-membered aromatic heterocyclic groups (preferably pyridyl);  $C_{6-14}$  aryl which may be substituted (preferably with  $C_{1-6}$  alkyl) (preferably phenyl, methylphenyl);  $C_{7-19}$  aralkyl (preferably benzyl);  $C_{6-14}$  aryl-carbonyl which may be substituted (preferably with halogen atom) (preferably fluoro-benzoyl, chlorobenzoyl);  $C_{1-6}$  alkoxy- $C_{1-6}$  alkyl (preferably methoxymethyl);  $C_{1-6}$  alkoxy-carbonyl (preferably tert-butoxycarbonyl), etc.

[0134] Preferably,  $R^1$  and  $R^2$ , together with the adjacent nitrogen atom, form a nitrogen-containing heterocyclic ring which may be substituted.

[0135] In particular,  $R^1$  and  $R^2$ , together with the adjacent nitrogen atom, form piperidino, pyrrolidin-1-yl, etc.

[0136] As the "nitrogen-containing heterocyclic rings which may be substituted" formed by  $R^2$  together with the adjacent nitrogen atom and Y, those exemplified as the "nitrogen-containing heterocyclic rings which may be substituted" formed by the above  $R^1$  and  $R^2$  together with the adjacent nitrogen atom, can be mentioned.

[0137] As the "nitrogen-containing condensed heterocyclic rings" in the "nitrogen-containing condensed heterocyclic rings which may be substituted" formed by  $R^2$  together with the adjacent nitrogen atom, Y and Ar, for example, 11- to 20-membered, preferably 11- or 18-membered condensed polycyclic (preferably tricyclic or tetracyclic) heterocyclic rings which contain at least one nitrogen atom and further contain one or more (for example, 1 to 4) hetero atoms selected from nitrogen, sulfur and oxygen atoms, in addition to carbon atoms, etc., can be mentioned. Specific examples include tetrahydrobenzo[g]isoquinoline, tetrahydrobenzo[b][1,6]naphthyridine, tetrahydrobenzo[b][1,7]naphthyridine, tetrahydropyrido[4,3-g]quinoline, tetrahydropyrido[3,4-g]quinoline, tetrahydropyrido[3,4-g]isoquinoline, tetrahydropyrido[3,4-g]quinoxaline, tetrahydropyrido[3,4-b]quinoxaline, tetrahydropyrido[4,3-b][1,5]naphthyridine, tetrahydropyrido[3,4-b][1,5]naphthyridine, tetrahydropyrido[3,4-g]quinoxaline, tetrahydropyrido[3,4-g]phthalazine, tetrahydrodronaphtho[2,3-d]piperazine, tetrahydroazepino[4,5-g]isoquinoline, tetrahydroazepino[4,5-b]quinoline, tetrahydroazepino[4,5-b]quinoxaline, tetrahydroazepino[4,5-g]quinoxaline, tetrahydroazepino[4,5-b][1,5]naphthyridine, tetrahydroazepino[4,5-g]phthalazine, hexahydrodronaphtho[4,5-d]azocine, hexahydroazocino[4,5-b]quinoline, tetrahydro- $\beta$ -carboline, tetrahydro-

pyrido[4,3-b]indole, tetrahydropyrrolo[3,2-g]isoquinoline, tetrahydropyrrolo[2,3-g]isoquinoline, tetrahydropyrrolo[3,4-b]acridine, tetrahydropyrrolo[3,4-b]phenazine, etc. In particular, tetrahydrobenzo[b][1,6]naphthyridine, etc. are preferred.

[0138] As the "substituents" in the "nitrogen-containing condensed heterocyclic rings which may be substituted", for example, those exemplified as the "substituents" in the above "5- to 7-membered saturated cyclic amino" which may be substituted can be used. The number of substituents is, for example, 1 to 5, preferably 1 to 3. When the number of substituents is 2 or more, each substituent can be the same or different.

[0139] The substituent is preferably optionally halogenated C<sub>1-6</sub> alkyl (preferably methyl).

[0140] Among the compounds represented by the formula (I), the compounds wherein X is X<sup>1</sup> [wherein X<sup>2</sup> is CONR<sup>8</sup>, NR<sup>8</sup>CO (wherein R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl or optionally halogenated C<sub>1-6</sub> alkylsulfonyl), OCO or COO] (provided that, when X<sup>1</sup> is CONR (wherein R is hydrogen atom or C<sub>1-6</sub> alkyl), Ar is not indole or benzoxazole which may have one or two halogen, hydroxy, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> alkoxy; when X<sup>1</sup> is CONH, Ar is not 4-methyl-2-quinolone which may have a substituent selected from the group consisting of alkyl, alkoxy and halogen, or is not 2-benzoylamino-quinazoline; and, when X<sup>1</sup> is COO, Ar<sup>1</sup> is not an aromatic group which may be substituted), that is, the compounds represented by the formula (I') are novel compounds.

[0141] Among the compounds represented by the formula (I'), the compounds wherein X<sup>1</sup> is CONR<sup>8</sup> or NR<sup>8</sup> (wherein R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl or optionally halogenated C<sub>1-6</sub> alkylsulfonyl) and Ar is quinoline or naphthalene, and the like are preferred.

[0142] Suitable examples of the compounds represented by the formula (I') include the following compounds:

4'-chloro-N-[6-[(N,N-dimethylamino)methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[6-(1-piperidinylmethyl)-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 N-(4-bromophenyl)-6-[(dimethylamino)methyl]-2-naphthamide;  
 N-(4'-chloro[1,1'-biphenyl]-4-yl)-6-[(N,N-dimethylamino)-methyl]-2-naphthamide;  
 4'-chloro-N-[2-[(N,N-dimethylamino)methyl]-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4'-fluoro-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4'-fluoro-N-[2-(1-piperidinylmethyl)-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[2-(1-piperidinylmethyl)-6-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 6-(4-fluorophenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide;  
 6-(4-chlorophenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide;  
 6-(4-methylphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide;  
 6-(4-fluorophenyl)-N-[6-[(dimethylamino)methyl]-2-naphthyl]nicotinamide;  
 6-(4-chlorophenyl)-N-[6-[(dimethylamino)methyl]-2-naphthyl]nicotinamide;  
 6-(4-methylphenyl)-N-[6-[(dimethylamino)methyl]-2-naphthyl]nicotinamide;  
 6-(4-fluorophenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-methylphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-methylphenyl)-N-[6-[(dimethylamino)methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 6-(4-fluorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl]nicotinamide;  
 6-(4-chlorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl]nicotinamide;  
 6-(4-fluorophenyl)-N-[2-[(dimethylamino)methyl]-6-quinolyl]nicotinamide;  
 6-(4-chlorophenyl)-N-[2-[(dimethylamino)methyl]-6-quinolyl]nicotinamide;  
 4-(4-fluorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl]-1-piperidinecarboxamide;  
 4-(4-chlorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolyl]-1-piperidinecarboxamide;  
 4-(4-fluorophenyl)-N-[2-[(dimethylamino)methyl]-6-quinolyl]-1-piperidinecarboxamide;  
 4-(4-chlorophenyl)-N-[2-[(dimethylamino)methyl]-6-quinolyl]-1-piperidinecarboxamide;  
 6-(4-fluorophenyl)-N-[7-(1-pyrrolidinylmethyl)-3-quinolyl]nicotinamide;  
 6-(4-chlorophenyl)-N-[7-[(dimethylamino)methyl]-3-quinolyl]nicotinamide;  
 4-(4-fluorophenyl)-N-[7-(1-pyrrolidinylmethyl)-3-quinolyl]-1-piperidinecarboxamide;  
 4-(4-chlorophenyl)-N-[7-[(dimethylamino)methyl]-3-quinolyl]-1-piperidinecarboxamide;  
 4'-chloro-N-[7-[(dimethylamino)methyl]-3-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 4'-fluoro-N-[7-(1-pyrrolidinylmethyl)-3-quinolyl][1,1'-biphenyl]-4-carboxamide;  
 5-(4-chlorophenyl)-N-[6-[(dimethylamino)methyl]-2-naphthyl]-2-pyridinecarboxamide;  
 5-(4-fluorophenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-2-pyridinecarboxamide;  
 4'-fluoro-N-[6-[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[6-[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 4'-methoxy-N-[6-[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;

N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]-4'-methyl[1,1'-biphenyl]-4-carboxamide;  
 4'-fluoro-N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4'-methoxy[1,1'-biphenyl]-4-carboxamide;  
 5 N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4'-methyl[1,1'-biphenyl]-4-carboxamide;  
 4'-fluoro-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 4'-chloro-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]-4'-methoxy[1,1'-biphenyl]-4-carboxamide;  
 4'-methyl-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 10 N-[6-[[4-(4-hydroxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]-4'-methoxy[1,1'-biphenyl]-4-carboxamide;  
 N-[6-[[4-(4-aminophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4'-methoxy[1,1'-biphenyl]-4-carboxamide;  
 4'-methoxy-N-[6-[[4-(4-nitrophenyl)-1-piperidinyl]methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide;  
 N-[6-[[4-(4-chlorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4'-methoxy[1,1'-biphenyl]-4-carboxamide;  
 6-(4-fluorophenyl)-N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 15 6-(4-chlorophenyl)-N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 6-(4-methoxyphenyl)-N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]-6-(4-methylphenyl)nicotinamide;  
 6-(4-fluorophenyl)-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 6-(4-chlorophenyl)-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 20 6-(4-methoxyphenyl)-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 6-(4-chlorophenyl)-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 6-(4-fluorophenyl)-N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 6-(4-chlorophenyl)-N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]nicotinamide;  
 N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-6-(4-methoxyphenyl)nicotinamide;  
 25 N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-6-(4-methylphenyl)nicotinamide;  
 4-(4-fluorophenyl)-N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-chlorophenyl)-N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-methoxyphenyl)-N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 N-[6-[[4-(4-methoxyphenyl)-1-piperidinyl]methyl]-2-naphthyl]-4-(4-methylphenyl)-1-piperidinecarboxamide;  
 30 4-(4-fluorophenyl)-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-chlorophenyl)-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-methoxyphenyl)-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-methylphenyl)-N-[6-[[4-(4-methylphenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 35 4-(4-fluorophenyl)-N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 4-(4-chlorophenyl)-N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4-(4-methoxyphenyl)-1-piperidinecarboxamide;  
 N-[6-[[4-(4-fluorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4-(4-methylphenyl)-1-piperidinecarboxamide;  
 N-[6-[[4-(4-chlorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4-(4-fluorophenyl)-1-piperidinecarboxamide;  
 4-(4-chlorophenyl)-N-[6-[[4-(4-chlorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-1-piperidinecarboxamide;  
 40 N-[6-[[4-(4-chlorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4-(4-methoxyphenyl)-1-piperidinecarboxamide;  
 N-[6-[[4-(4-chlorophenyl)-1-piperidinyl]methyl]-2-naphthyl]-4-(4-methylphenyl)-1-piperidinecarboxamide.

[0143] Examples of salts of compound (I) or (I') include salts with inorganic bases, ammonium salts, salts with organic bases, salts with inorganic acids, salts with organic acids, salts with basic or acidic amino acids, and the like.

45 [0144] Preferred examples of salts with inorganic bases include alkali metal salts such as sodium salts, potassium salts, etc.; alkaline earth metal salts such as calcium salts, magnesium salts, barium salts, etc.; aluminum salts, etc.

[0145] Preferred examples of salts with organic bases include salts with trimethylamine, triethylamine, pyridine, picoline, ethanolamine, diethanolamine, triethanolamine, dicyclohexylamine, N,N-dibenzylethylenediamine, etc.

50 [0146] Preferred examples of salts with inorganic acids include salts with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, etc.

[0147] Preferred examples of salts with organic acids include salts with formic acid, acetic acid, trifluoroacetic acid, fumaric acid, oxalic acid, tartaric acid, maleic acid, citric acid, succinic acid, malic acid, methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid, etc.

55 [0148] Preferred examples of salts with basic amino acids include salts with arginine, lysine, ornithine, etc. Preferred examples of salts with acidic amino acids include salts with aspartic acid, glutamic acid, etc.

[0149] Among these salts, pharmaceutically acceptable salts are preferred. For example, when compound (I) or (I') possesses an acidic functional group, it can form an inorganic salt such as an alkali metal salt (e.g., sodium salt, potassium salt, etc.), an alkaline earth metal salt (e.g., calcium salt, magnesium salt, barium salt, etc.), etc., an ammo-

nium salt, etc. When compound (I) or (I') possesses a basic functional group, it can form an inorganic salt such as hydrochloride, sulfate, phosphate, hydrobromate, etc.; or an organic salt such as acetate, maleate, fumarate, succinate, methanesulfonate, p-toluenesulfonate, citrate, tartarate, etc.

[0150] Compounds (I) and (I') (hereinafter also abbreviated as the compound of the present invention) can be either anhydrides or hydrates. A hydrate may have 0.5 to 3 water molecules.

[0151] In addition, the compounds of the present invention can be labeled using isotopes (e.g.  $^3\text{H}$ ,  $^{14}\text{C}$ , and  $^{35}\text{S}$ , etc.).

[0152] When the compound of the present invention contain optical isomers, stereoisomers, regio isomers, rotational isomers, these are also included as the compound of the present invention, and each of them can be obtained as a single substance by per se known synthesis methods and separation methods. For example, when optical isomers exist in the compound of the present invention, the optical isomers resolved from the compound are included in the compound of the present invention.

[0153] The optical isomers can be produced using per se known methods. Specifically, the optical isomer can be obtained by using an optically active synthetic intermediate, or subjecting a racemic mixture of the final product to optical resolution in accordance with common method.

[0154] Examples of optical resolution methods include per se known methods such as the fractional recrystallization method, chiral column method, diastereomer method, etc., which are described in detail below.

#### 1) Fractional recrystallization method

[0155] The method which comprises allowing a racemate to form a salt with an optically active compound (e.g. (+)-mandelic acid, (-)-mandelic acid, (+)-tartaric acid, (-)-tartaric acid, (+)-1-phenethylamine, (-)-1-phenethylamine, cinchonine, (-)-cinchonidine, brucine, etc.), separating the salt using a fractional recrystallization method, followed by, if desired, neutralizing process to obtain a free optical isomer.

#### 2) Chiral column method

[0156] This method comprises subjecting a racemate or its salt to a column for separating an optical isomer (chiral column) for separation. For example, in the case of liquid chromatography, an optical isomer mixture is added to the chiral column such as ENANTIO-OVM [produced by Tosoh] or CHIRAL series [produced by Daicel], which is developed using water, various buffer solutions (e.g. phosphate buffer), organic solvents (e.g. ethanol, methanol, isopropanol, acetonitrile, trifluoroacetic acid, diethylamine, etc.) as single or mixed solutions, and the optical isomers are separated. Also, in the case of gas chromatography, for example, separation is conducted using a chiral column such as CP-Chirasil-DeX (produced by G.L.Science Co.).

#### 3) Diastereomer method

[0157] In this method, a racemic mixture is subjected to a chemical reaction with an optically active reagent to give a diastereomer mixture, which is separated into a single substance by an ordinary separation means (e.g. fractional recrystallization, chromatography method, etc.). This single substance is subjected to removal of the optically active reagent part using chemical processing such as a hydrolysis reaction. For example, when a compound of the present invention possesses hydroxy or primary or secondary amino in its molecule, this compound is subjected to a condensation reaction with an optically active organic acid (e.g. MTPA [ $\alpha$ -methoxy- $\alpha$ -(trifluoromethyl)phenylacetic acid], (-)-menthoxyacetic acid, etc.), to give the diastereomer in an ester form or an amide form, respectively. On the other hand, when a compound of the present invention possesses carboxylic acid group, this compound is subjected to a condensation reaction with an optically active amine or alcohol reagent, to give the diastereomer in an amide form or an ester form, respectively. The separated diastereomer can be converted to an optical isomer of the original compound, by applying acidic hydrolysis or basic hydrolysis.

[0158] A prodrug of compound (I) is a compound which is converted to compound (I') by reactions involving enzymes and gastric acid, etc. under physiological conditions in the living body; in other words, a compound that is changed into compound (I') by enzymatically-caused oxidation, reduction and hydrolysis, and a compound that is changed into compound (I') by hydrolysis caused by gastric acid. Examples of the prodrugs of compound (I) include compounds in which amino groups of compound (I) have been acylated, alkylated, or phosphorylated [e.g. compounds in which amino groups of compound (I) have been eicosanoylated, alanylated, pentyaminocarbonylated, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methoxycarbonylated, tetrahydrofuranylated, pyrrolidylmethylated, pivaloyloxymethylated, tert-butylated, etc.]; compounds in which hydroxyl groups of compound (I) have been acetylated, alkylated, phosphorylated, borated [e.g. compounds in which hydroxyl groups of compound (I) have been acetylated, palmitoylated, propanoylated, pivaloylated, succinylated, fumarilated, alanylated, dimethylaminomethylcarbonylated, etc.]; compounds in which carboxyl groups of compound (I) have been esterified or amidated [e.g. compounds in which carboxyl groups of com-

pond (I') have been ethylesterified, phenylesterified, carboxymethylesterified, dimethylaminomethylesterified, pivaloyloxymethylesterified, ethoxycarbonyloxylesterified, phthalidylesterified, (5-methyl-2-oxo-1,3-dioxolen-4-yl)methylesterified, cyclohexyloxycarbonylethylesterified, or methylamidated, etc.]. These compounds can be produced from compound (I') using per se known methods.

[0159] Also, a produg of compound (I') can be a compound which is changed to compound (I') by physiological conditions, as described in pages 163 to 198 of Molecular Design, volume 7, "Development of Drugs", published in 1990 by Hirokawa Shoten.

[0160] The compound of the present invention can be produced by [Production method 1] to [Production method 7] which are described in detail below, or analogous methods thereto.


[0161] Compounds (II) to (XIII), compound (IV'), compound (IIIa), compound (IIIb), compound (IIIc), compound (IIId), compound (IIIf), compound (IIHg) and compound (IIIf), used as raw materials, can be used in the form of salts, respectively. As such salts, those exemplified as salts of the above compound (I) or (I') can be used.

[0162] In the following [Production method 1] to [Production method 7], when an alkylation reaction, a hydrolysis reaction, an amination reaction, an esterification reaction, an amidation reaction, an esterification reaction, an etherification reaction, an oxidation reaction, a reduction reaction, etc. are carried out, these reactions are carried out in accordance with per se known methods. Examples of such methods include the methods described in Organic Functional Group Preparations, Second Edition, Academic Press, Inc., published in 1989; Comprehensive Organic Transformations, VCH Publishers Inc., published in 1989, etc.

[Production method 1]

[0163] Compound (Ia) having  $-(CH_2)_{w3}CONR^{8a}(CH_2)_{w4}$  for X in the formula (I) is produced, for example, by the following amidation reaction.

(Amidation reaction)



wherein  $R^{8a}$  is hydrogen atom or an optionally halogenated  $C_{1-6}$  alkyl; the other symbols are as defined above.

[0164] As the "optionally halogenated  $C_{1-6}$  alkyl", those exemplified as "substituents" in the above "cyclic group" which may be substituted" can be used.

[0165] The "amidation reaction" includes the following "method using a dehydration and condensation agent" and "method using a reactive derivative of carboxylic acid".

i) Method using a dehydration and condensation agent

[0166] Compound (III), 1 to 5 equivalents of compound (II), and 1 to 2 equivalents of a dehydration and condensation agent are reacted in an inert solvent. If necessary, the reaction can be carried out with the coexistence of 1 to 1.5 equivalents of 1-hydroxybenzotriazole (HOBT) and/or catalytic quantity to 5 equivalents of a base.

[0167] Examples of the "dehydrating and condensation agent" include dicyclohexylcarbodiimide (DCC), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (WSC). WSC is particularly preferable.

[0168] Examples of the "inert solvent" include nitrile solvents (preferably acetonitrile), amide solvents (preferably DMF), halogenated hydrocarbon solvents (preferably dichloromethane), ether solvents (preferably THF). Two or more

kinds of these can be mixed in an appropriate ratio for use.

[0169] Examples of the "base" include

- 1) strong bases exemplified by hydrides of alkali metals or alkaline earth metals (e.g. lithium hydride, sodium hydride, potassium hydride, calcium hydride, etc.), amides of alkali metals or alkaline earth metals (e.g. lithium amide, sodium amide, lithium diisopropylamide, lithium dicyclohexylamide, lithium hexamethyldisilazide, sodium hexamethyldisilazide, potassium hexamethyldisilazide, etc.), lower alkoxides of alkali metals or alkaline earth metals (e.g. sodium methoxide, sodium ethoxide, potassium tert-butoxide, etc.), etc.;
- 2) inorganic bases exemplified by hydroxides of alkali metals or alkaline earth metals (e.g. sodium hydroxide, potassium hydroxide, lithium hydroxide, barium hydroxide, etc.), carbonates of alkali metals or alkaline earth metals (e.g. sodium carbonate, potassium carbonate, cesium carbonate, etc.) and hydrogencarbonates of alkali metals or alkaline earth metals (e.g. sodium hydrogencarbonate, potassium hydrogencarbonate, etc.), etc.; and
- 3) organic bases exemplified by amines such as triethylamine, diisopropylethylamine, N-methylmorpholine, dimethylaminopyridine, DBU (1,8-diazabicyclo[5.4.0]undec-7-en), DBN (1,5-diazabicyclo[4.3.0]non-5-en), etc.; basic heterocyclic compounds such as pyridine, imidazole, 2,6-lutidine, etc.; and the like.

[0170] Among the above bases, triethylamine, 4-dimethylaminopyridine, etc., are preferable.

[0171] Reaction temperature is usually room temperature (0°C to 30°C, hereafter the same). Reaction time is, for example, 10 to 24 hours.

ii) Method using a reactive derivative of carboxylic acid

[0172] A reactive derivative of compound (II) and 1 to 5 equivalents (preferably 1 to 3 equivalents) of compound (III) are reacted in an inert solvent. If necessary, the reaction can be carried out with the coexistence of 1 to 10 equivalents, preferably 1 to 3 equivalents of a base.

Examples of the "reactive derivative" of compound (II) include acid halides (e.g., acid chloride, acid bromide, etc.), mixed acid anhydrides (e.g. acid anhydrides with C<sub>1-6</sub> alkyl-carboxylic acid, C<sub>6-10</sub> aryl-carboxylic acid or C<sub>1-6</sub> alkylcarbonate), active esters (e.g. esters with phenol which may be substituted, 1-hydroxybenzotriazole or N-hydroxysuccinimide, etc.), etc.

[0173] Examples of the "substituents" in the "phenol which may be substituted" include halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), nitro, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkoxy. The number of substituents is, for example, 1 to 5.

[0174] As the "optionally halogenated C<sub>1-6</sub> alkyl" and "optionally halogenated C<sub>1-6</sub> alkoxy", those exemplified as "substituents" in the above "cyclic group which may be substituted" can be used.

[0175] Specific examples of "phenol which may be substituted" include phenol, pentachlorophenol, pentafluorophenol, p-nitrophenol, etc. The reactive derivative is, preferably, an acid halide.


[0176] Examples of the "inert solvent" include ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents, and water. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, THF, dichloromethane, chloroform, etc. are preferable.

[0177] As the "base", the same as above are used. The base is preferably sodium hydride, potassium carbonate, sodium carbonate, sodium hydroxide, potassium hydroxide, sodium hydrogencarbonate, potassium hydrogencarbonate, triethylamine, pyridine, etc.

[0178] Reaction temperature is usually -20°C to 50°C, preferably room temperature. Reaction time is usually 5 minutes to 40 hours, preferably 1 to 18 hours.

[0179] Compound (II) described above can be produced by per se known methods.

[0180] Compound (III) can be produced by subjecting the compound of the formula:



wherein W is a protecting group for amino; and the other symbols are as defined above, to a deprotection reaction to

remove W.


[0181] Examples of the protecting group for amino represented by W include formyl, C<sub>1-6</sub> alkyl-carbonyl (e.g. acetyl, propionyl, etc.), C<sub>1-6</sub> alkoxy-carbonyl (e.g. methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl, etc.), benzoyl, C<sub>7-10</sub> aralkyl-carbonyl (e.g. benzylcarbonyl, etc.), C<sub>7-14</sub> aralkyloxy-carbonyl (e.g. benzyloxycarbonyl, 9-fluorenylmethoxycarbonyl, etc.), trityl, phthaloyl, N,N-dimethylaminomethylene, silyl (e.g. trimethylsilyl, triethylsilyl, dimethylphenylsilyl, tert-butylidimethylsilyl, tert-butylidiethylsilyl, etc.), C<sub>2-6</sub> alkenyl (e.g. 1-allyl, etc.), etc. These groups may be substituted by 1 to 3 of halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.), C<sub>1-6</sub> alkoxy (e.g. methoxy, ethoxy, propoxy, etc.) or nitro, etc.

[0182] The deprotection reaction is carried out, for example, by maintaining compound (IIIa), preferably at 20°C to 140°C, in an aqueous solution of an acid such as a mineral acid (e.g., hydrochloric acid, sulfuric acid, hydrobromic acid, iodic acid, periodic acid, etc.) etc., or a base such as an alkali metal hydroxide (e.g., sodium hydroxide, potassium hydroxide, lithium hydroxide, etc.) etc. The acid or base is usually used in an amount of 1 to 100 equivalents, preferably 1 to 40 equivalents based on compound (IIIa). Strength of the acid or base is usually 0.1 N to 18 N, preferably 1 N to 12 N. Reaction time is usually 0.5 hour to 48 hours, preferably 1 hour to 24 hours.


[0183] Further, when W is t-butoxycarbonyl group, etc., the deprotection reaction can also be carried out by dissolving compound (IIIa) in an organic acid (e.g., trifluoroacetic acid, formic acid, acetic acid, methanesulfonic acid, benzenesulfonic acid, trifluoromethanesulfonic acid, etc.) and maintaining the solution usually at -20°C to 200°C, preferably 0°C to 100°C. The organic acid is used in an amount of 1 to 100 equivalents, preferably 1 to 40 equivalents based on compound (IIIa).

[0184] The deprotection reaction can also be carried out by subjecting compound (IIIa) to catalytic reduction in an alcoholic solvent, for example, ethanol, etc., or a solvent such as acetic acid, etc., with a catalyst such as palladium, palladium-carbon, Raney nickel, Raney cobalt, platinum oxide, etc. at normal pressure or, if necessary, under pressure.

[0185] Compound (IIIa) can be produced by reacting a compound of the formula:



wherein L is a leaving group and the other symbols are as defined above, with a compound of the formula:



wherein the symbols are as defined above.

[0186] Examples of the "leaving group" represented by L include halogen atom (e.g. chlorine, bromine, iodine, etc.), optionally halogenated C<sub>1-6</sub> arylsulfonyloxy (e.g. methanesulfonyloxy, ethanesulfonyloxy, trifluoromethanesulfonyloxy, etc.), C<sub>6-10</sub> arylsulfonyloxy which may be substituted, hydroxy, etc.

[0187] Examples of the "substituents" in the "C<sub>6-10</sub> arylsulfonyloxy which may be substituted" include halogen atom (e.g. chlorine, bromine, iodine, etc.), optionally halogenated C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy, etc. The number of substituents is, for example, 1 to 3. Specific examples of the "C<sub>6-10</sub> arylsulfonyloxy which may be substituted" include benzenesulfonyloxy, p-toluenesulfonyloxy, 1-naphthalenesulfonyloxy, 2-naphthalenesulfonyloxy, etc.

[0188] The "leaving group" is preferably halogen atom (e.g. chlorine, bromine, iodine, etc.), methanesulfonyloxy, trifluoromethanesulfonyloxy, p-toluenesulfonyloxy, etc.

[0189] This reaction is usually carried out in an inert solvent.

[0190] Examples of the "inert solvent" include alcohol solvents, ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents, water, etc. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, N,N-dimethylformamide (DMF), acetone, ethanol, pyridine, etc., are preferred.


[0191] Compound (IIIi) is used in an amount of 1 equivalent to 100 equivalents based on compound (IIIb). Further,

compound (IIIi) can be used in an amount corresponding to a reaction solvent.

[0192] Reaction temperature is about -20°C to 200°C, preferably room temperature to 100°C. Reaction time is, for example, 0.5 hour to 1 day.

[0193] This reaction can be carried out in the presence of a base. The base is preferably sodium hydride, potassium carbonate, sodium carbonate, sodium hydroxide, potassium hydroxide, sodium hydrogencarbonate, potassium hydrogencarbonate, triethylamine, pyridine, etc. The amount of the base is 0.1 to 100 equivalents, preferably 1 to 10 equivalents based on compound (IIIb).

[0194] Compound (IIIb) can be produced, for example, from the compound of the formula:



wherein the symbols are as defined above.

[0195] In compound (IIIb), the compound wherein L is optionally halogenated C<sub>1-5</sub> alkylsulfonyloxy, or C<sub>6-10</sub> arylsulfonyloxy which may be substituted can be produced by subjecting compound (IIIh) to a known acylation reaction. This reaction is carried out, for example, by reacting compound (IIIh) with 1 to 5 equivalents of a corresponding sulfonyl halide in an inert solvent in the presence of a base. The base is preferably potassium carbonate, sodium hydrogen carbonate, triethylamine, N-methylmorpholine, pyridine, etc. The base is preferably used in an amount of 1 to 10 equivalents.

[0196] Examples of the "inert solvent" include ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents, etc.

[0197] Reaction temperature is -20°C to 200°C, preferably 0°C to 100°C. Reaction time is 0.1 hour to 48 hours, preferably 1 hour to 24 hours.

[0198] In compound (IIIb), the compound wherein L is a halogen atom can be produced by subjecting compound (IIIh) to a known halogenation reaction.

[0199] This reaction is carried out by using a halogenating agent. Examples of the halogenating agent include an inorganic acid chloride such as thionyl chloride, thionyl bromide, phosphorus trichloride, phosphorus pentachloride, phosphorus oxychloride, etc.; a hydrogen halide acid such as hydrogen chloride acid, hydrobromic acid, etc., and the like.


[0200] Further, in compound (IIIb), the compound wherein L is a halogen atom can be produced by subjecting compound (IIIi) described hereinafter to the reaction described in Journal of Medicinal Chemistry, 35, 2761 (1992), etc. or a modification thereof. In this reaction, a halogenating agent such as bromine, N-bromosuccinimide, etc. and an additive such as benzoylperoxide, 2,2'-azobis(isobutyronitrile), etc. are used.

[0201] Compound (IIIh) can be produced by reducing an ester compound (IIIc) described hereinafter by a known reduction method. As a reduction method, for example, there are a method using a reducing agent (e.g., a boron hydride reagent such as sodium borohydride, etc.; an aluminum hydride reagent such as lithium aluminum hydride, etc.), and the like.

[0202] Further, compound (IIIh) can be produced from the compound wherein Ar is 2-methylquinolines by N-oxide transfer according to a method described in a literature (e.g., Journal of Medicinal Chemistry 34, 3212 (1991); Journal of Medicinal Chemistry 35, 2761 (1992); etc.).

[0203] The above compound (IIIi) can be produced by a per se known method.

[0204] Compound (IIIa) can also be produced by reacting a compound of the formula:



wherein R<sup>8a</sup> is a bond or an optionally halogenated bivalent C<sub>1-5</sub> non-cyclic hydrocarbon group, R<sup>8b</sup> is hydrogen atom or an optionally halogenated C<sub>1-5</sub> alkyl group, and the other symbols are as defined above, with the above compound (IIIi).



[0205] Here, examples of the "optionally halogenated bivalent  $C_{1-6}$  non-cyclic hydrocarbon group" include, among the "optionally halogenated bivalent  $C_{1-6}$  non-cyclic hydrocarbon group" exemplified with respect to the above X and Y, the group having 1 to 5 carbon atoms.

[0206] Further, examples of the "optionally halogenated  $C_{1-5}$  alkyl group" include, among the "optionally halogenated  $C_{1-5}$  alkyl group" exemplified as the substituents of the above "cyclic group which may be substituted", the group having 1 to 5 carbon atoms.

[0207] This reaction can be carried out by reacting compound (IIIc) and, usually, 1 to 20 equivalents, preferably 1 to 5 equivalents of compound (IIIi) with a reducing agent in an inert solvent.

[0208] Examples of the "inert solvent" include alcohol solvents, ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, organic acid solvents, etc. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, methanol, ethanol, acetic acid, etc., are preferred.

[0209] Examples of the reducing agent include sodium borohydride, sodium triacetoxylborohydride, sodium cyanoborohydride, etc. The reducing agent is usually used in an amount of 1 to 20 equivalents, preferably 1 to 5 equivalents.


[0210] Reaction temperature is usually  $-20^{\circ}\text{C}$  to  $150^{\circ}\text{C}$ , preferably  $20$  to  $100^{\circ}\text{C}$ . Reaction time is usually 5 minutes to 40 hours, preferably 1 to 24 hours.

[0211] This reaction can also be carried out in the presence of an acid. Examples of the acid to be used include organic acids such as acetic acid, methanesulfonic acid, etc.; inorganic acid such as hydrochloric acid, sulfuric acid, etc.; and the like. The acid is used in an amount of 0.01 equivalent to 0.1 equivalent in case of an inorganic acid, and 0.01 equivalent to 100 equivalents or an amount corresponding to a solvent in case of an organic acid.

[0212] Compound (IIIc) can be produced by subjecting the above compound (IIIh) to a known oxidation reaction. The oxidation reaction can be carried out, for example, by using an oxidizing agent. As the oxidizing agent, there can be used, for example, manganese dioxide, chromic acid, lead tetraacetate, silver oxide, copper oxide, halogen acid, oxidation using dimethylsulfoxide (Swern oxidation), organic peracids, oxygen, electrode oxidation, etc.


[0213] Further, compound (IIIc) can also be produced from an ester compound (IIId) described hereinafter by a known method with an organic metal reagent such as Grignard reagent, lithium dialkylcopper, etc.

[0214] Compound (IIIa) can also be produced by subjecting a compound of the formula:



wherein the symbols are as defined above, and compound (IIIi) to a per se known condensation reaction (for example, a method using the above dehydration condensation agent, a method using a reactive derivative of carboxy), and subjecting the resultant amide compound to a known reduction reaction. The reduction reaction is usually carried out by using a reducing agent. As the reducing agent, there can be used, for example, a borohydride reagent such as diborane, sodium borohydride, etc., an aluminum hydride reagent such as lithium aluminum hydride, etc., and the like.

[0215] Further, compound (IIIa) can also be produced by converting a compound of the formula:



wherein the symbols are as defined above, into an enamine compound by a known method (for example, the method described in Heterocycles, 22, 195 (1984), etc.) and subjecting the resultant enamine compound to a known reduction reaction.

[0216] Here, the enamine compound is produced by using, for example, N,N-dimethylformamide dialkylacetal, etc.

[0217] The reduction reaction is usually carried out by using a reducing agent. As the reducing agent, there can be used sodium borohydride, sodium triacetoxylborohydride, sodium cyanoborohydride, or catalytic hydrogenation, etc.

[0218] This production method is also applicable to an active methylene group at an adjacent position of  $-\text{CO}-$  in case that, for example, Y is a divalent group consisting of  $-\text{CO}-$  and an alkylene in compound (IIIa).

[0219] The above compound (IIId) can be produced by a per se known method. For example, methyl 6-amino-2-naphthalene carboxylate and methyl 5-amino-2-naphthalene carboxylate can be produced according to the method described in WO98/43953, etc.

[0220] These aminonaphthalene carboxylic acids can also be produced by hydrolyzing corresponding naphthalene dicarboxylates according to, for example, the method described in JP 06107599, etc. to form monoesters and the resultant carboxylic acids are subjected to the reactions described in Journal of Organic Chemistry, 60, 4412 (1995); Chemical Pharmaceutical Bulletin, 35, 2698 (1987); etc.


[0221] The above compound (IIId) can be produced by a per se known method. For example, 6-amino-2-methylquinoline can be produced by the methods described in Polymer Bulletin, 42, 175 (1999), Journal of Organic Chemistry, 28, 1753 (1963), Journal of Chemical Society C, 829 (1970), etc. or a modification thereof.

[0222] The above "method using a reactive derivative of carboxy" is also applicable to the production of the corresponding sulfonamide derivative and sulfinamide derivative from a sulfonic acid represented by the formula:  $\text{Ar}^1\text{-(CH}_2\text{)}_3\text{-SO}_2\text{OH}$  (wherein the symbols are as defined above) and a sulfinic acid of the formula:  $\text{Ar}^1\text{-(CH}_2\text{)}_3\text{-SOOH}$  (wherein the symbols are as defined above), respectively.

[Production method 2]

[0223] Compound (Ib) having  $\text{-(CH}_2\text{)}_w\text{3-COO(CH}_2\text{)}_w\text{4-}$  for X in the formula (I), can be produced by the following esterification reaction.

(Esterification reaction)



wherein the symbols are as defined above.

[0224] A reactive derivative of compound (II) and 1 to 5 equivalents (preferably 1 to 3 equivalents) of compound (IV) is reacted in an inert solvent. Usually, this reaction is carried out with the coexistence of 1 to 10 equivalents, preferably 1 to 3 equivalents of a base.

[0225] As the reactive derivative of compound (II), the same as above is used. Especially, an acid halide is preferable.

[0226] Examples of the "inert solvent" include ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, dichloromethane, chloroform, etc. are preferable.

[0227] As the "base", the same one as above can be used. The base is preferably sodium hydride, potassium carbonate, sodium carbonate, sodium hydroxide, potassium hydroxide, sodium hydrogencarbonate, potassium hydrogen carbonate, triethylamine, pyridine, etc.


[0228] Reaction temperature is usually -20°C to 50°C, preferably room temperature. Reaction time is usually 5 minutes to 40 hours, preferably 1 to 18 hours.

[0229] Compound (IV) can be produced by a per se known method, for example the method described in WO9838156 or modification thereof.

[Production method 3]

[0230] Compound (Ic) having  $-(CH_2)_{w1}O(CH_2)_{w2}-$  for X in the formula (I), can be produced by, for example, the following etherification reaction.

(Etherification reaction)



wherein the symbols are as defined above.

[0231] Compound (IV) and about 1 to 5 equivalents (preferably 1 to 2 equivalents) of compound (V) are reacted in inert solvent, with the coexistence of base.

[0232] As the "base", the same one as above can be used. The base is preferably potassium carbonate, sodium hydrogencarbonate, triethylamine, N-methylmorpholine, pyridine, etc. The amount of the base used is usually about 1 to 5 equivalents relative to compound (V).

[0233] Examples of the "inert solvent" include alcohol solvents, ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents, water. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, N,N-dimethylformamide (DMF), acetone, ethanol, pyridine, etc., are preferable.

[0234] Reaction temperature is about  $-20^\circ\text{C}$  to  $100^\circ\text{C}$ , preferably room temperature to  $80^\circ\text{C}$ . Reaction time is, for example, about 0.5 hour to 1 day.

[0235] In the above production method, when the leaving group is hydroxy, Mitsunobu reaction can usually be used. In the Mitsunobu reaction, compound (V) and 0.5 to 5 equivalents (preferably 1 to 1.5 equivalents) of compound (IV') are reacted in inert solvent with the coexistence of 0.5 to 5 equivalents (preferably 1 to 1.5 equivalents) of ethyl acetyl-dicarboxylate.

[0236] Examples of the inert solvent include ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, dichloromethane, chloroform, etc. are preferable.

[0237] Reaction temperature is usually  $-20^\circ\text{C}$  to  $50^\circ\text{C}$ , preferably room temperature. Reaction time is usually 5 minutes to 40 hours, preferably 1 to 18 hours.


[0238] Compound (V) can be produced by a per se known method.

[0239] Compound (IV') can be produced by a per se known method, for example, the method described in WO9838156 or modification thereof.

[Production method 4]

[0240] Compound (Id) having  $-(CH_2)_{w3}NR^{6a}CO(CH_2)_{w4}-$  for X in the formula (I), can be produced, for example, by the following amidation reaction.

## (Amidation reaction)



wherein the symbols are as defined above.

[0241] This production method is carried out in accordance with the above Production method 1.


[0242] Compound (VI) can be produced by a per se known method.

[0243] Compound (VII) can be produced by a per se known method, for example the method described in WO9838156 or modification thereof.

[Production method 5]

[0244] Compound (Ie) having  $-(\text{CH}_2)_{w5}\text{NHCONR}^{\text{a}}(\text{CH}_2)_{w6}-$  for X in the formula (I), can be produced, for example, by the following urea reaction.

## (Urea reaction)



wherein the symbols are as defined above.

[0245] Compound (IX) and 1 to 5 equivalents (preferably 1 to 1.5 equivalents) of compound (VIII) is reacted in an inert solvent with the coexistence of a base.

[0246] As the "base", the same one as above can be used. The base is preferably potassium carbonate, sodium

carbonate, sodium hydroxide, potassium hydroxide, sodium hydrogencarbonate, potassium hydrogencarbonate, triethylamine, pyridine, etc.

[0247] Examples of the "inert solvent" include alcohol solvents, ether solvents, halogenated hydrocarbon solvents, aromatic solvents, nitrile solvents, amide solvents, ketone solvents, sulfoxide solvents, water. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, acetonitrile, DMF, acetone, ethanol, pyridine, etc. are preferable.


[0248] Reaction temperature is usually -20°C to 100°C, preferably room temperature to 80°C. Reaction time is, for example, 0.5 hour to 1 day.

[0249] Compound (VIII) and compound (IX) can be produced by a per se known method.

[Production method 6]

[0250] Compound (II) having, for Ar<sup>1</sup>, a ring assembly aromatic group (Ar<sup>2</sup>-Ar<sup>3</sup>) which may be substituted in the formula (I), can be produced by, for example, the following aryl-coupling reaction.

### (Aryl-coupling reaction)



wherein Ar<sup>2</sup> and Ar<sup>3</sup> are monocyclic aromatic groups or condensed aromatic groups, each of which may be substituted; L<sup>1</sup> is hydroxy or C<sub>1-6</sub> alkyl; L<sup>2</sup> is halogen (preferably chlorine, bromine) or trifluoromethanesulfonyloxy; the other symbols are as defined above.

[0251] As "substituents", "monocyclic aromatic groups" and "condensed aromatic groups" in the "monocyclic aromatic groups or condensed aromatic groups, each of which may be substituted" for Ar<sup>2</sup> and Ar<sup>3</sup>, those exemplified as the above Ar<sup>1</sup> can be used. Especially, it is preferable that both of Ar<sup>2</sup> and Ar<sup>3</sup> are phenyl groups which may be substituted, and Ar<sup>2</sup>-Ar<sup>3</sup> is biphenyl which may be substituted.

[0252] The aryl-coupling reaction can be carried out in accordance with per se known methods such as the method described in Acta. Chemica Scandinavica, pp. 221-230, 1993, or methods analogous thereto.

[0253] Compound (X) and 1 to 3 equivalents (preferably 1 to 1.5 equivalents) of compound (XI) are reacted in an inert solvent in the presence of a base and a transition metal catalyst.

[0254] As the base, the same one as above can be used. The base is preferably sodium carbonate, sodium hydrogencarbonate, etc.

[0255] The amount of the "base" used is, for example, about 1 to 10 equivalents relative to compound (XI).

[0256] Examples of the "transition metal catalyst" include palladium catalyst, nickel catalyst. Examples of the "palladium catalyst" include tetrakis(triphenylphosphine)palladium (O), palladium acetate, bis (triphenylphosphine) palladium (II) chloride, palladium-carbon. Examples of the "nickel catalyst" include tetrakis(triphenylphosphine) nickel (O).

[0257] The amount of the "transition metal catalyst" used is about 0.01 to 1 equivalent, preferably about 0.01 to 0.5 equivalent, relative to compound (XI).

[0258] Reaction temperature is room temperature to 150°C, preferably about 80°C to 150°C. Reaction time is, for example, about 1 to 48 hours.

[0259] Examples of the "inert solvent" include water, alcohol solvents, aromatic solvents. Two or more kinds of these can be mixed in an appropriate ratio for use. Especially, a single solvent such as water, ethanol and toluene; or a mixed solvent of two or more kinds of these is preferred.

[0260] Compound (X) and compound (XI) can be produced by a per se known method.


[0261] Among compound (I), compound (I') can also be produced by the following [Production method 7].

[Production method 7]

[0262] Compound (I') can also be produced by reacting a compound of the formula:



wherein  $\text{Ar}^1$  is as defined above, or a salt thereof with a compound of the formula:



wherein the symbols are as defined above, or a salt thereof.

[0263] This production method can be carried out according to the above Production method 1.

[0264] Compound (XII) and compound (XIII) can be produced by a per se known method.

[0265] Examples of the above "alcohol solvents" include methanol, ethanol, isopropanol, tert-butanol, etc.

[0266] Examples of the above "ether solvents" include diethylether, tetrahydrofuran (THF), 1,4-dioxane, 1,2-dimethoxyethane, etc.

[0267] Examples of the above "halogenated hydrocarbon solvents" include dichloromethane, chloroform, 1,2-dichloroethane, carbon tetrachloride, etc.

[0268] Examples of the above "aromatic solvents" include benzene, toluene, xylene, pyridine, etc.

[0269] Examples of the above "hydrocarbon solvents" include hexane, pentane, cyclohexane, etc.

[0270] Examples of the above "amide solvents" include N,N-dimethylformamide (DMF), N,N-dimethylacetamide, N-methylpyrrolidone, etc.

[0271] Examples of the above "ketone solvent" include acetone, methyl ethyl ketone, etc.

[0272] Examples of the above "sulfoxide solvents" include dimethylsulfoxide (DMSO), etc.

[0273] Examples of the above "nitrile solvents" include acetonitrile, propionitrile, etc.

[0274] In a compound of the present invention thus obtained, the intramolecular functional group can be converted to a desired functional group by combining per se known chemical reactions. Examples of the chemical reactions include oxidation reaction, reduction reaction, alkylation reaction, hydrolysis reaction, amination reaction, esterification reaction, aryl-coupling reaction, deprotection reaction.

[0275] In each of the above reactions, when the starting material compounds possess amino, carboxy, hydroxy, and/or carbonyl as substituents, protecting groups which are generally used in peptide chemicals, etc., can be introduced into these groups, and the desired compound can be obtained by removing the protecting groups after the reaction if necessary.

[0276] Examples of the protecting group for amino include those exemplified with respect to the above W.

[0277] Examples of the protecting group for carboxy include  $\text{C}_{1-6}$  alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, etc.),  $\text{C}_{7-11}$  aralkyl (e.g. benzyl, etc.), phenyl, trityl, silyl (e.g. trimethylsilyl, triethylsilyl, dimethylphenylsilyl, tert-butyl dimethylsilyl, tert-butyl diethylsilyl, etc.),  $\text{C}_{2-6}$  alkenyl (e.g. 1-allyl, etc.). These groups may be substituted by 1 to 3 of halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.),  $\text{C}_{1-6}$  alkoxy (e.g. methoxy, ethoxy, propoxy, etc.) or nitro.

[0278] Examples of the protecting group for hydroxy include  $\text{C}_{1-6}$  alkyl (e.g. methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, etc.), phenyl, trityl,  $\text{C}_{7-10}$  aralkyl (e.g. benzyl, etc.), formyl,  $\text{C}_{1-6}$  alkyl-carbonyl (e.g. acetyl, propionyl, etc.), benzoyl,  $\text{C}_{7-10}$  aralkyl-carbonyl (e.g. benzylcarbonyl, etc.), 2-tetrahydropyranylyl, 2-tetrahydrofuranylyl, silyl (e.g. trimethylsilyl, triethylsilyl, dimethylphenylsilyl, tert-butyl dimethylsilyl, tert-butyl diethylsilyl, etc.),  $\text{C}_{2-6}$  alkenyl (e.g. 1-allyl, etc.). These groups may be substituted by 1 to 3 of halogen atom (e.g. fluorine, chlorine, bromine, iodine, etc.),  $\text{C}_{1-6}$  alkyl (e.g. methyl, ethyl, n-propyl, etc.),  $\text{C}_{1-6}$  alkoxy (e.g. methoxy, ethoxy, propoxy, etc.) or nitro, etc. can be substituted for

these groups.

[0279] Examples of the protecting group for carbonyl include cyclic acetal (e.g. 1,3-dioxane, etc.), and non-cyclic acetal (e.g. di-C<sub>1-6</sub> alkylacetal, etc.).

[0280] Removal of the above protecting groups can be carried out in accordance with per se known methods such as those described in Protective Groups in Organic Synthesis, published by John Wiley and Sons (1980). For instance, the methods using acid, base, ultraviolet light, hydrazine, phenylhydrazine, sodium N-methyldithiocarbamate, tetrabutylammonium fluoride, palladium acetate, trialkylsilyl halide (e.g. trimethylsilyl iodide, trimethylsilyl bromide, etc.), and a reduction method, etc. can be used.

[0281] The compound of the present invention can be isolated and purified by per se known methods such as solvent extraction, changing of liquid properties, transdissolution, crystallization, recrystallization, chromatography, etc. It is also possible to isolate and purify the starting material compounds of the compound of the present invention, or their salts using the same known methods as above, but they can also be used as raw materials in the next process as a reaction mixture without being isolated.

[0282] The compound of the present invention possesses an excellent MCH receptor antagonistic action, therefore, it is useful as an agent for preventing or treating diseases caused by MCH. Also, the compound of the present invention is low in toxicity, and is excellent in oral absorbency and intracerebral transitivity.

[0283] Therefore, a melanin-concentrating hormone antagonist (hereafter, also abbreviated as "MCH antagonist") comprising a compound of the present invention can be safely administered to mammals (e.g. rats, mice, guinea pigs, rabbits, sheep, horses, swine, cattle, monkeys, humans, etc.) as an agent for preventing or treating diseases caused by MCH.

[0284] Here, examples of the diseases caused by MCH include obesity (e.g. malignant mastocytosis, exogenous obesity, hyperinsular obesity, hyperplastic obesity, hypophyseal adiposity, hypoplastic obesity, hypothyroid obesity, hypothalamic obesity, symptomatic obesity, infantile obesity, upper body obesity, alimentary obesity, hypogonadal obesity, systemic mastocytosis, simple obesity, central obesity, etc.), hyperphagia, emotional disorders, reproductive function disorders, etc.

[0285] The compound of the present invention is also useful as an agent for preventing or treating lifestyle diseases such as diabetes, diabetic complications (e.g. diabetic retinopathy, diabetic neuropathy, diabetic nephropathy, etc.), arteriosclerosis, gonitis, etc.

[0286] Further, the compound of the present invention is useful as an anorectic agent.

[0287] The MCH antagonist and the pharmaceutical composition of the present invention can be used in combination with an alimentary therapy (e.g., alimentary therapy for diabetes) and exercise.

[0288] The MCH antagonist and the pharmaceutical composition of the present invention can be produced by subjecting compound (I) or compound (I') respectively, as it is, or together with a pharmacologically acceptable carrier, to pharmaceutical manufacturing process in accordance with a per se known means.

[0289] Here, examples of the pharmacologically acceptable carriers include various organic or inorganic carrier substances which are commonly used as materials for pharmaceutical preparations, such as excipients, lubricants, binders, and disintegrators in solid preparations; solvents, solubilizing agents, suspending agents, isotonicizing agents, buffering agents, soothing agents, in liquid preparations; and the like. Also, in the pharmaceutical manufacturing process, additives such as antiseptics, antioxidants, coloring agents, sweeteners, absorbents, moistening agents, etc., can be used, if necessary.

[0290] Examples of the excipients include lactose, sucrose, D-mannitol, starch, cornstarch, crystalline cellulose, light anhydrous silicic acid, etc.

[0291] Examples of the lubricants include magnesium stearate, calcium stearate, talc, colloidal silica, etc.

[0292] Examples of the binders include crystalline cellulose, sucrose, D-mannitol, dextrin, hydroxypropylcellulose, hydroxypropylmethylcellulose, polyvinylpyrrolidone, starch, saccharose, gelatin, methylcellulose, carboxymethylcellulose sodium, etc.

[0293] Examples of the disintegrators include starch, carboxymethylcellulose, carboxymethylcellulose calcium, croscarmellose sodium, carboxymethylstarch sodium, low-substituted hydroxypropylcellulose (L-HPC), etc.

[0294] Examples of the solvents include distilled water for injection, alcohol, propylene glycol, macrogol, sesame oil, corn oil, etc.

[0295] Examples of the solubilizing agents include polyethylene glycol, propylene glycol, D-mannitol, benzyl benzoate, ethanol, trisaminomethane, cholesterol, triethanolamine, sodium carbonate, sodium citrate, etc.

[0296] Examples of the suspending agents include surfactants such as stearyltriethanolamine, sodium lauryl sulfate, lauryl amino propionic acid, lecithin, benzalkonium chloride, benzethonium chloride, glyceryl monostearate; or hydrophilic polymers such as polyvinyl alcohol, polyvinylpyrrolidone, carboxymethylcellulose sodium, methylcellulose, hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, etc.

[0297] Examples of the isotonicizing agents include glucose, D-sorbitol, sodium chloride, glycerin, D-mannitol, etc.

[0298] Examples of the buffering agents include buffer solutions of phosphate, acetate, carbonate, citrate, etc.

[0299] Examples of the soothing agents include benzyl alcohol, etc.

[0300] Examples of the antiseptics include paraoxybenzoates, chlorobutanol, benzyl alcohol, phenethylalcohol, dehydroacetic acid, sorbic acid, etc.

[0301] Examples of the antioxidants include sulfite, ascorbic acid, etc.

[0302] The MCH antagonist and the pharmaceutical composition of the present invention can be safely administered orally or parenterally (e.g. by local, rectal and intravenous administration) in various dosage forms, for example, as oral drugs such as tablets (including sugar-coated tablets and film-coated tablets), powders, granules, capsules (including soft capsules), solutions; and parenteral preparations such as injectable preparations (e.g. preparations for subcutaneous injections, intravenous injections, intramuscular injections, intraperitoneal injections, etc.), external preparations (e.g. nasal preparations, percutaneous preparations, ointments, etc.), suppositories (e.g. rectal suppositories, vaginal suppositories, etc.), sustained-release preparations (e.g. sustained-release microcapsules, etc.), pellets, drip infusions, etc.

[0303] The content of compound (I) in the MCH antagonist of the present invention and the content of compound (I') in the pharmaceutical composition of the present invention are, for example, about 0.1 to 100% by weight based on the total weight of the MCH antagonist or pharmaceutical composition, respectively.

[0304] The dose of the MCH antagonist and the pharmaceutical composition of the present invention can be appropriately selected depending on the subject of administration, route of administration, disease, etc.

[0305] For example, the dose per day when the MCH antagonist or the pharmaceutical composition of the present invention is orally administered to an adult obesity patient (body weight: about 60 kg), is about 0.1 to about 500 mg, preferably about 1 to about 100 mg, more preferably about 5 to about 100 mg, in terms of compound (I) or compound (I'), each of which is an active ingredient. These amounts can be divided into one to several doses per day for administration.

[0306] The MCH antagonist and pharmaceutical composition of the present invention can be used in combination with other concomitant drugs which do not interfere with the MCH antagonist and pharmaceutical composition of the present invention, for the purpose of "strengthening of therapeutic effect against obesity", "reduction of dose of MCH antagonist", etc. Examples of the concomitant drugs include a "agents for treating diabetes", "agents for treating diabetic complications", "agents for treating obesity other than MCH antagonists", "agents for treating hypertension", "agents for treating hyperlipidemia (agents for treating arteriosclerosis)", "agents for treating arthritis", "anxiety agents", "antidepressant", etc. Two or more kinds of these concomitant drugs can be combined in an appropriate ratio for use.

[0307] Examples of the above "agents for treating diabetes" include insulin sensitizers, insulin secretion enhancers, biguanides, insulins,  $\alpha$ -glucosidase inhibitors,  $\beta$ 3 adrenaline receptor agonists, etc.

[0308] Examples of the insulin sensitizers include pioglitazone or its salt (preferably hydrochloride), troglitazone, rosiglitazone or its salt (preferably maleate), JTT-501, GI-262570, MCC-555, YM-440, DRF-2593, BM-13-1258, KRP-297, R-119702, etc.

[0309] Examples of the insulin secretion enhancers include sulfonylureas. Specific examples of the sulfonylureas include tolbutamide, chlorpropamide, trazamide, acetohexamide, glycopyramide and its ammonium salt, glibenclamide, gliclazide, glimepiride, etc.

[0310] Other than the above, examples of insulin secretion enhancers include repaglinide, nateglinide, mitiglinide (KAD-1229), JTT-608, etc.

[0311] Examples of biguanides include metformin, buformin, phenformin, etc.

[0312] Examples of insulins include animal insulins extracted from bovine or porcine pancreas; semi-synthetic human insulin which is enzymatically synthesized from insulin extracted from porcine pancreas; human insulin synthesized by genetic engineering, using *Escherichia coli* and yeast; etc. As insulin, also employed are insulin-zinc containing 0.45 to 0.9 (w/w)% of zinc; protamine-insulin-zinc produced from zinc chloride, protamine sulfate and insulin; etc. In addition, insulin can be an insulin fragment or derivative (e.g. INS-1, etc.).

[0313] Insulin can also include various types such as ultra immediate action type, immediate action type, two-phase type, intermediate type, prolonged action type, etc., and these can be selected depending on the pathological conditions of patients.

[0314] Examples of  $\alpha$ -glucosidase inhibitors include acarbose, voglibose, miglitol, emiglitate, etc.

[0315] Examples of  $\beta$ 3 adrenaline receptor agonists include AJ-9677, BMS-196085, SB-226552, AZ40140, etc.

[0316] Other than the above, examples of the "agents for treating diabetes" include ergoset, pramlintide, leptin, BAY-27-9955, etc.

[0317] Examples of the above "agents for treating diabetic complications" include aldose reductase inhibitors, glycation inhibitors, protein kinase C inhibitors, etc.

[0318] Examples of aldose reductase inhibitors include torulestat; eparestat; imirestat; zenarestat; SNK-860; zopolrestat; ARI-509; AS-3201, etc.

[0319] Examples of glycation inhibitors include pimagedine. Examples of protein kinase C inhibitors include NGF, LY-333531, etc.



[0320] Other than the above, examples of "agents for treating diabetic complications" include alprostadil, thiapride hydrochloride, clobazam, mexiletine hydrochloride, ethyl eicosapentate, memantine, pimegeline (ALT-711), etc.

[0321] Examples of the above "agents for treating obesity other than MCH antagonists" include lipase inhibitors and anorectics, etc.

[0322] Examples of lipase inhibitors include orlistat, etc.

[0323] Examples of anorectics include mazindol, dexfenfluramine, fluoxetine, sibutramine, baiamine, etc.

[0324] Other than the above, examples of "agents for treating obesity other than MCH antagonists" include lipstatin, etc.

[0325] Examples of the above "agents for treating hypertension" include angiotensin converting enzyme inhibitors, calcium antagonists, potassium channel openers, angiotensin II antagonists, etc.

[0326] Examples of angiotensin converting enzyme inhibitors include captopril, enalapril, alacepril, delapril (hydrochloride), lisinopril, imidapril, benazepril, cilazapril, temocapril, trandolapril, manidipine (hydrochloride), etc.

[0327] Examples of calcium antagonists include nifedipine, amlodipine, felodipine, nicardipine, etc.

[0328] Examples of potassium channel openers include levromakalim, L-27152, AL0671, NIP-121, etc.

[0329] Examples of angiotensin II antagonists include losartan, candesartan cilexetil, valsartan, irbesartan, CS-866, E4177, etc.

[0330] Examples of the above "agents for treating hyperlipidemia (agents for treating arteriosclerosis)" include HMG-CoA reductase inhibitors, fibrate compounds, etc.

[0331] Examples of HMG-CoA reductase inhibitors include pravastatin, simvastatin, lovastatin, atorvastatin, fluvastatin, lipantil, cerivastatin, itavastatin, ZD-4522, or their salts (e.g. sodium salts, etc.), etc.

[0332] Examples of fibrate compounds include bezafibrate, ciprofibrate, clofibrate, simfibrate, etc.

[0333] Examples of the above "agents for treating arthritis" include ibuprofen, etc.

[0334] Examples of the above "anxiety agents" include chlordiazepoxide, diazepam, oxazolam, medazepam, clonazepam, bromazepam, lorazepam, alprazolam, fludiazepam, etc.

[0335] Examples of the above "antidepressants" include fluoxetine, fluvoxamine, imipramine, paroxetine, sertraline, etc.

[0336] The timing of administration of the above concomitant drugs is not limited. The MCH antagonist or pharmaceutical composition and the concomitant drugs can be administered to the subject simultaneously or at staggered times.

[0337] The dosages of the concomitant drugs can be determined in accordance with clinically used dosages, and can be appropriately selected according to the subject of administration, route of administration, diseases and combinations of drugs, etc.

[0338] The administration forms for the concomitant drugs are not particularly limited as long as the MCH antagonist or the pharmaceutical composition are used in combination with a concomitant drugs at the time of administration.

Examples of such administration forms includes 1) administration of a single preparation obtained by simultaneous preparation of MCH antagonist or pharmaceutical composition together with concomitant drugs, 2) simultaneous administration of two kinds of preparations obtained by separate preparation of MCH antagonist or pharmaceutical composition, and concomitant drugs, through the same route of administration, 3) staggered administration of two kinds of preparations obtained by separate preparation of MCH antagonist or pharmaceutical composition, and concomitant drugs, through the same route of administration, 4) simultaneous administration of two kinds of preparations obtained by separate preparation of MCH antagonist or pharmaceutical composition, and concomitant drugs, through different routes of administration, 5) staggered administration of two kinds of preparations obtained by separate preparation of MCH antagonist or pharmaceutical composition, and concomitant drugs, through different routes of administration (for example, administration of MCH antagonist or pharmaceutical composition; and concomitant drugs in this order; or administration in reverse order).

[0339] The ratio of combination of MCH antagonist or pharmaceutical composition with concomitant drugs can be appropriately selected in accordance with the subject of administration, route of administration and diseases, etc.

#### BEST MODE FOR CARRYING OUT THE INVENTION

[0340] The present invention will be explained further in detail by the following Reference Examples, Examples, Preparation Examples, and Experimental Examples. However, these do not limit the present invention, and they can be changed within the scope that does not deviate from the scope of the present invention.

[0341] In the following Reference Examples and Examples, "room temperature" means 0 to 30°C. Anhydrous magnesium sulfate or anhydrous sodium sulfate was used to dry the organic layer. "%" means percent by weight, unless otherwise specified.

[0342] Infrared absorption spectra were determined by the diffuse reflectance method, using fourier transform type infrared spectrophotometer.

[0343] FABMS (pos) is mass spectrum determined by the (+) method, in Fast Atom Bombardment Mass Spectrometry.

[0344] Other symbols used in the description have the following meanings.

5	s :	singlet
	d :	doublet
	t :	triplet
	q :	quartet
	m :	multiplet
10	br :	broad
	J :	coupling constant
	Hz :	Hertz
	CDCl <sub>3</sub> :	Deuterated chloroform
	DMSO-d <sub>6</sub> :	Deuterated dimethylsulfoxide
15	THF :	tetrahydrofuran
	DMF :	N,N-dimethylformamide
	DMSO :	dimethylsulfoxide
	WSCD :	1-ethyl-3-(3-dimethylaminopropyl) carbodimide
	WSC :	1-ethyl-3-(3-dimethylaminopropyl) carbodimide hydrochloride
20	<sup>1</sup> H-NMR :	proton nuclear resonance (Free substances were usually measured in CDCl <sub>3</sub> .)
	IR :	infrared absorption spectrum
	Me :	methyl
	Et :	ethyl
25	HOBT :	1-hydroxy-1H-benzotriazole
	IPE :	diisopropyl ether
	DMAP :	4-dimethylaminopyridine

[0345] In this specification and drawings, when bases and amino acids are shown by codes, these codes are based on those by the IUPAC-IUB Commission on Biochemical Nomenclature or common codes in the concerned fields. Examples of these codes are shown below. Also, where some optical isomers of amino acids can exist, the L form is shown unless otherwise specified.

	DNA :	deoxyribonucleic acid
35	cDNA :	complementary deoxyribonucleic acid
	A :	adenine
	T :	thymine
	G :	guanine
	C :	cytosine
40	RNA :	ribonucleic acid
	mRNA :	messenger ribonucleic acid
	dATP :	deoxyadenosine triphosphate
	dTTP :	deoxythymidine triphosphate
	dGTP :	deoxyguanosine triphosphate
45	dCTP :	deoxycytidine triphosphate
	ATP :	adenosine triphosphate
	EDTA :	ethylenediamine tetraacetic acid
	SDS :	sodium dodecyl sulfate
	EIA :	enzyme immunoassay
50	Gly :	glycine
	Ala :	alanine
	Val :	valine
	Leu :	leucine
	Ile :	isoleucine
55	Ser :	serine
	Thr :	threonine
	Cys :	cysteine
	Met :	methionine

Glu : glutamic acid  
 Asp : aspartic acid  
 Lys : lysine  
 Arg : arginine  
 5 His : histidine  
 Phe : phenylalanine  
 Tyr : tyrosine  
 Trp : tryptophan  
 Pro : proline  
 10 Asn : asparagine  
 Gln : glutamine  
 pGlu : pyroglutamine  
 Me : methyl group  
 Et : ethyl group  
 15 Bu : butyl group  
 Ph : phenyl group  
 TC : thiazolidine-4(R)-carboxamide group

[0346] Substituents, protecting groups and reagents frequently used in this specification, are shown by the following symbols.

Tos : p-toluenesulfonyl  
 CHO : formyl  
 Bzl : benzyl  
 25 Cl<sub>2</sub>Bzl : 2,6-dichlorobenzyl  
 Bom : benzyloxymethyl  
 Z : benzyloxycarbonyl  
 Cl-Z : 2-chlorobenzyloxycarbonyl  
 Br-Z : 2-bromobenzyloxycarbonyl  
 30 Boc : t-butoxycarbonyl  
 DNP : dinitrophenol  
 Trt : trityl  
 Bum : t-butoxymethyl  
 Fmoc : N-9-fluorenylmethoxycarbonyl  
 35 HOBT : 1-hydroxybenzotriazole  
 HOObt : 3,4-dihydro-3-hydroxy-4-oxo-1,2,3-benzotriazine  
 HONB : 1-hydroxy-5-norbornene-2,3-dicarboxylic diimide  
 DCC : N,N'-dicyclohexylcarbodiimide

SEQ ID NO in the SEQUENCE LISTING in the specification of the present application shows the following sequences.

[SEQ ID NO: 1] shows a synthetic DNA used for screening of cDNA coding rat SLC-1.

[SEQ ID NO: 2] shows a synthetic DNA used for screening of cDNA coding rat SLC-1.

[SEQ ID NO: 3] shows an entire amino acid sequence of rat SLC-1.

[SEQ ID NO: 4] shows an entire base sequence of rat SLC-1cDNA wherein Sal I recognition sequence was added to the 5' side, and Spe I recognition sequence was added to the 3' side.

[SEQ ID NO: 5] shows riboprobe used to determine the quantity of SLC-1mRNA expressed in each clone of rat SLC-1 expression CHO cells.

[SEQ ID NO: 6] shows a synthetic DNA used to obtain cDNA for coding of human SLC-1.

[SEQ ID NO: 7] shows a primer used to make double-strand cDNA for coding human SLC-1.

[SEQ ID NO: 8] shows an entire base sequence of cDNA for coding human SLC-1.

[SEQ ID NO: 9] shows an entire amino acid sequence of human SLC-1.

[SEQ ID NO: 10] shows a synthetic DNA used for screening of cDNA for coding human SLC-1(S).

[SEQ ID NO: 11] shows a synthetic DNA used for screening of cDNA for coding human SLC-1(S).

[SEQ ID NO: 12] shows a synthetic DNA used for screening of cDNA for coding human SLC-1(L).

[SEQ ID NO: 13] shows a synthetic DNA used for screening of cDNA for coding human SLC-1(L).

[SEQ ID NO: 14] shows an entire base sequence of human SLC-1(S) cDNA wherein Sal I recognition sequence was added to the 5' side, and Spe I recognition sequence was added to the 3' side.

[SEQ ID NO: 15] shows an entire base sequence of human SLC-1(L) cDNA wherein Sal I recognition sequence was added to the 5' side, and Spe I recognition sequence was added to the 3' side.


[SEQ ID NO: 16] shows riboprobe used to determine the quantity of SLC-1mRNA expressed in each clone of human SLC-1(S) expression CHO cells and SLC-1(L) expression CHO cells.

Transformant *Escherichia coli* DH10B/pSLC1L8 transformed by plasmid containing DNA which codes the base sequence shown by SEQ ID NO: 9, obtained in Reference Example 1 - 6, has been deposited with National Institute of Bioscience and Human-Technology (NIBH), Agency of Industrial Science and Technology, Ministry of International Trade and Industry, under accession of number FERM BP-6632 since February 1, 1999, and with the Institute for Fermentation, Osaka, Japan (IFO), under accession number of IFO 16254 since January 21, 1999.

#### Examples

#### Reference Example 1

#### [0347] Tert-butyl 6-[(N,N-dimethylamino)methyl]-2-naphthylcarbamate



1) 2,6-naphthalenedicarboxylic acid dimethyl ester (26.0 g, 106 mmol) was dissolved in N,N-dimethylformamide (500 ml), and a 1N aqueous sodium hydroxide solution (106 ml) was added dropwise thereto at 100°C over 30 minutes. After stirred for 3 hours, the solvent was distilled off under reduced pressure, water was added to the residue and the insolubles were filtered off. Concentrated hydrochloric acid (9 ml) was added to the filtrate, the precipitated crude product was filtered, washed with water, and recrystallized from hot methanol to obtain 6-(methoxycarbonyl)-2-naphthalenecarboxylic acid (14.6 g) as a white powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 3.94 (3H, s), 8.06 (2H, m), 8.24 (2H, m), 8.69 (2H, s).

2) The 6-(methoxycarbonyl)-2-naphthalenecarboxylic acid (5.00 g, 21.7 mmol) obtained in the above 1) and triethylamine (3.93 ml, 28.2 mmol) were dissolved in tert-butylalcohol (55 ml), diphenylphosphoryl azide (5.62 ml, 26.1 mmol) was added thereto, and the mixture was stirred at room temperature for 30 minutes and at 100°C for 6 hours. To the reaction solution were added ethyl acetate and an aqueous saturated sodium bicarbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with a 10% aqueous citric acid solution and an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting crude carbamate was dissolved in tetrahydrofuran (50 ml), lithium aluminum hydride (728 mg, 19.2 mmol) was added under ice-cooling, and the mixture was stirred at room temperature for 3 hours. To the reaction solution were added ethyl acetate and a 10% aqueous citric acid solution, the mixture was extracted, the organic layer was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, concentrated under reduced pressure, the residue was purified by silica gel column chromatography (developing solvent: toluene: ethyl acetate=10:1), and the eluent was crystallized from n-hexane to obtain tert-butyl 6-(hydroxymethyl)-2-naphthylcarbamate (2.26 g) as a white powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.51 (9H, s), 4.61 (2H, d, J = 5.7 Hz), 5.24 (1H, t, J = 5.7 Hz), 7.40 (1H, d, J = 8.4 Hz), 7.49 (1H, m), 7.70-7.78 (3H, m), 8.07 (1H, s), 9.52 (1H, s).

Elemental analysis for C <sub>16</sub> H <sub>19</sub> NO <sub>3</sub>			
Calcd	C, 70.31;	H, 7.01;	N, 5.12
Found	C, 70.36;	H, 6.89;	N, 5.14

3) The tert-butyl 6-(hydroxymethyl)-2-naphthylcarbamate (1.00 g, 3.66 mmol) obtained in the above 2) was dissolved in dichloromethane (18 ml), manganese dioxide (1.59 g, 18.3 mmol) was added thereto, and the mixture was stirred at room temperature for 4 hours. The insolubles were filtered, the filtrate was concentrated under reduced pressure, and crystallized from diisopropyl ether to obtain tert-butyl 6-formyl-2-naphthylcarbamate (889 mg) as a white powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.57 (9H, s), 6.76 (1H, br), 7.44 (1H, m), 7.83 (1H, d, J = 8.4 Hz), 7.92 (2H, m), 8.09 (1H, s),


8.25 (1H, s), 10.10 (1H, s).

4) The tert-butyl 6-formyl-2-naphthylcarbamate (300 mg, 1.11 mmol) obtained in the above 3) and dimethylamine hydrochloride (270 mg, 3.32 mmol) were dissolved in a mixed solution of methanol (2 ml) and tetrahydrofuran (2 ml), sodium cyanotrihydroborate (210 mg, 3.32 mmol) was added thereto, and the mixture was stirred at room temperature for 2 hours. To the reaction solution was added an aqueous potassium carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by silica gel column chromatography (developing solvent: ethyl acetate:methanol=10:1) to obtain the titled compound (242 mg) as a colorless oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.55 (9H, s), 2.27 (6H, s), 3.54 (2H, s), 6.75 (1H, s), 7.32 (1H, dd, J = 3.1, 8.7 Hz), 7.42 (1H, d, J = 8.4 Hz), 7.64 (1H, s), 7.71 (2H, d-like, J = 8.7 Hz), 7.96 (1H, s).

#### Reference Example 2

[0348] Tert-butyl 6-(1-pyrrolidinylmethyl)-2-naphthylcarbamate




[0349] The tert-butyl 6-(hydroxymethyl)-2-naphthylcarbamate (500 mg, 1.83 mmol) obtained in 2) in Reference Example 1 and triethylamine (0.254 ml, 1.83 mmol) were dissolved in tetrahydrofuran (9 ml), methanesulfonyl chloride (0.142 ml, 1.83 mmol) was added under ice-cooling, and the mixture was stirred at room temperature for 30 minutes. The insolubles were filtered, and the filtrate was concentrated under reduced pressure to obtain a mesylate. The resulting mesylate was dissolved in acetonitrile (9 ml), potassium carbonate (758 mg, 5.49 mmol) and pyrrolidine (0.153 ml, 1.83 mmol) were added thereto, and the mixture was stirred at 60°C for 3 hours. To the reaction solution were added ethyl acetate and water, the mixture was extracted, the organic layer was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by alumina column chromatography (developing solvent: ethyl acetate) to obtain tert-butyl 6-(1-pyrrolidinylmethyl)-2-naphthylcarbamate (388 mg) as a colorless oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.55 (9H, s), 1.80 (4H, m), 2.55 (4H, m), 3.74 (2H, s), 6.62 (1H, s), 7.30 (1H, m), 7.45 (1H, m), 7.69 (3H, m), 7.96 (1H, s).

#### Reference Example 3

[0350] Tert-butyl 6-(1-piperidinylmethyl)-2-naphthylcarbamate




[0351] Using the tert-butyl 6-formyl-2-naphthylcarbamate (500 mg, 1.83 mmol) obtained in 3) of Reference Example 1 and piperidine (0.181 ml, 1.83 mmol), the same procedures as those of 4) of Reference Example 1 were conducted to obtain the titled compound (305 mg) as a colorless oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.46 (2H, m), 1.55 (9H, s), 1.58 (4H, m), 2.42 (4H, m), 3.59 (2H, s), 6.60 (1H, s), 7.31 (1H, m), 7.43 (1H, m), 7.64 (1H, m), 7.71 (2H, m), 7.95 (1H, s).

#### Reference Example 4

[0352] N-(4-Bromophenyl)-6-(hydroxymethyl)-2-naphthamide



1) The 6-(methoxycarbonyl)-2-naphthalenecarboxylic acid (1.00 g, 4.34 mmol) obtained in 1) of Reference Example 1, 4-bromoaniline (747 mg, 4.34 mmol) and dimethylaminopyridine (531 mg, 4.34 mmol) were dissolved in N, N-dimethylformamide (10 ml), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (833 mg, 4.34 mmol) was added thereto under ice-cooling, and the mixture was stirred at room temperature for 1 hour and at 50°C for 2 hours. To the reaction solution was added 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed with a 1N aqueous sodium hydroxide solution and an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and the eluent was crystallized from ethyl acetate to obtain methyl 6-((4-bromoanilino)carbonyl)-2-naphthoate (954 mg) as a white powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 3.95 (3H, s), 7.58 (2H, d, J = 8.6 Hz), 7.82 (2H, d, J = 8.6 Hz), 8.07 (2H, d, J = 8.6 Hz), 8.22 (1H, d, J = 8.6 Hz), 8.29 (1H, d, J = 8.6 Hz), 8.63 (1H, s), 8.72 (1H, s), 10.64 (1H, s).

2) Methyl 6-((4-bromoanilino)carbonyl)-2-naphthoate (900 mg, 2.34 mmol) obtained in the above 1) was dissolved in tetrahydrofuran (10 ml), lithium aluminum hydride (178 mg, 4.68 mmol) was added thereto under ice-cooling, and the mixture was stirred at room temperature for 2 hours. To the reaction solution were added ethyl acetate and 1N hydrochloric acid, and the mixture was extracted with ethyl acetate. The extract was washed with an aqueous saturated sodium bicarbonate solution and an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. Chloroform was added to the residue to effect crystallization to obtain the titled compound (349 mg) as a white powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 4.71 (2H, d, J = 5.6 Hz), 5.41 (1H, t, J = 5.6 Hz), 7.56 (3H, m), 7.82 (2H, d, J = 8.4 Hz), 7.92 (1H, s), 8.02 (3H, m), 8.54 (1H, s), 10.53 (1H, s).

#### Reference Example 5

##### [0353] N-[2-(Hydroxymethyl)-6-quinolinyl]acetamide



1) 6-amino-2-methylquinoline (1.02 g, 6.45 mmol) was dissolved in pyridine (30 ml), acetic anhydride (0.913 ml, 9.67 mmol) was added thereto, and the mixture was stirred at room temperature for 3 hours. The solvent was distilled off under reduced pressure, and diisopropyl ether was added thereto for crystallization, to obtain N-(2-methyl-6-quinolinyl)acetamide (1.20 g) as white powders.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.22 (3H, s), 2.71 (3H, s), 7.25 (1H, m), 7.52 (1H, m), 7.95 (2H, m), 8.10 (1H, s), 8.30 (1H, s).

2) The N-(2-methyl-6-quinolinyl)acetamide (1.20 g, 5.99 mmol) obtained in the above 1) was dissolved in chloroform (30 ml), m-chloroperbenzoic acid (2.48 g, 7.19 mmol) was added thereto, and the mixture was stirred at room temperature for 3 hours. The solvent was distilled off under reduced pressure, ethyl acetate was added to the residue, and the precipitated powders were filtered to obtain N-(2-methyl-1-oxide-6-quinolinyl)acetamide (1.06 g) as a white powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.12 (3H, s), 2.53 (3H, s), 7.51 (1H, d, J = 8.4 Hz), 7.76 (2H, m), 8.40 (1H, s), 8.48 (1H, d, J = 9.3 Hz), 10.36 (1H, s).


3) The N-(2-methyl-1-oxide-6-quinolinyl)acetamide (4.64 g, 21.5 mmol) obtained in the above 2) was dissolved in acetic anhydride (110 ml), and the solution was stirred at 80°C for 4 hours. The solvent was distilled off under reduced pressure, and the residue was purified by alumina column chromatography (developing solvent: ethyl acetate) to obtain an oil. The resulting oil was dissolved in methanol (110 ml), a 1N aqueous sodium hydroxide solution (21.5 ml) was added thereto under ice-cooling, and the mixture was stirred at room temperature for 1 hour. The solvent was distilled off under reduced pressure. The resulting residue was purified by alumina column chro-

matography (developing solvent: ethyl acetate: methanol=5:1), and the eluate was crystallized from ethyl acetate-isopropyl ether (1:3) to obtain the titled compound (2.65 g) as white powders.

<sup>1</sup>H-NMR (CD<sub>3</sub>OD) δ: 2.23 (3H, s), 4.89 (2H, s), 7.68 (1H, d, J = 8.7 Hz), 7.78 (1H, d, J = 8.7 Hz), 7.95 (1H, d, J = 8.7 Hz), 8.27 (1H, d, J = 8.7 Hz), 8.33 (1H, s).

#### Reference Example 6

[0354] N-[2-(Chloromethyl)-6-quinolinyl]acetamide hydrochloride




[0355] To the N-[2-(hydroxymethyl)-6-quinolinyl]acetamide (1.00 g, 4.62 mmol) obtained in Reference Example 5 was added thionyl chloride (23 ml) under ice-cooling, and the mixture was stirred for 2 hours. The reaction solution was concentrated under reduced pressure, and the resulting residue was washed with ethyl acetate to obtain the titled compound (900 mg) as a powder.

<sup>1</sup>H-NMR (CD<sub>3</sub>OD) δ: 2.23 (3H, s), 5.12 (2H, s), 8.08 (1H, d, J = 8.4 Hz), 8.17 (2H, s-like), 8.74 (1H, s), 9.01 (1H, d, J = 8.4 Hz).

#### Reference Example 7

[0356] 2-(Chloromethyl)-6-quinolinylamine dihydrochloride




[0357] To the N-[2-(chloromethyl)-6-quinolinyl]acetamide hydrochloride (900 mg, 3.32 mmol) obtained in Reference Example 6 was added 5N hydrochloric acid (17 ml), and the mixture was stirred at 100°C for 2 hours. The reaction solution was concentrated under reduced pressure, and the resulting residue was washed with tetrahydrofuran to obtain the titled compound (849 mg) as powders.

<sup>1</sup>H-NMR (CD<sub>3</sub>OD) δ: 5.07 (2H, s), 7.42 (1H, d, J = 2.4 Hz), 7.71 (1H, dd, J = 2.4, 9.0 Hz), 7.95 (1H, d, J = 8.4 Hz), 8.06 (1H, d, J = 9.0 Hz), 8.78 (1H, d, J = 8.4 Hz).

#### Reference Example 8

[0358] 4'-Chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0359] Using the 2-(chloromethyl)-6-quinolinylamine dihydrochloride obtained in Reference Example 7 and 4'-chloro [1,1'-biphenyl]-4-carboxylic acid, the same procedures as those for an amidation reaction using WSC of Example 1 were conducted to obtain the titled compound as powders.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 4.95 (2H, s), 7.58 (2H, d, J = 8.4 Hz), 7.66 (1H, d, J = 8.4 Hz), 7.82 (2H, d, J = 8.4 Hz), 7.89

(2H, d, J = 7.8 Hz), 8.01 (1H, d, J = 9.0 Hz), 8.09 (1H, m), 8.13 (2H, d, J = 7.8 Hz), 8.41 (1H, d, J = 8.4 Hz), 8.61 (1H, s), 10.67 (1H, s).

# Reference Example 9

## [0360] 2-[(1-Pyrrolidinylmethyl)-6-quinolinyl]amine




[0361] To the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide (5.53 g, 20.5 mmol) obtained in 1) of Example 7 was added concentrated hydrochloric acid (100 ml), the mixture was stirred at 110°C for 1 hour, and the solvent was distilled off under reduced pressure. To the resulting residue was added ethyl acetate, the mixture was washed with an aqueous potassium carbonate solution and an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and the solvent was distilled off under reduced pressure to obtain the titled compound (4.56 g) as a powder from ethyl acetate-hexane.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.83 (4H, m), 2.62 (4H, m), 3.90 (2H, s), 3.92 (2H, br), 6.92 (1H, d, J = 2.8 Hz), 7.15 (1H, dd, J = 2.8, 8.6 Hz), 7.48 (1H, d, J = 8.6 Hz), 7.88 (1H, d, J = 2.8 Hz), 7.92 (1H, d, J = 8.6 Hz).  
m.p.: 102-104°C

# Reference Example 10

## [0362] N-[2-[(Diisopropylamino)methyl]-6-quinolinyl]acetamide




[0363] Using the N-[2-(hydroxymethyl)-6-quinolinyl]acetamide obtained in Reference Example 5, the same procedures as those of 1) of Example 6 were conducted to obtain the titled compound as a powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (12H, d, J = 6.3 Hz), 2.25 (3H, s), 2.95-3.16 (2H, m), 3.93 (2H, s), 7.40-7.64 (2H, m), 7.80 (1H, d, J = 8.4 Hz), 7.95 (1H, d, J = 8.4 Hz), 8.05 (1H, d, J = 8.1 Hz), 8.28 (1H, br).  
m.p.: 147-148°C (crystallization solvent: diethyl ether-hexane)

# Reference Example 11

## [0364] N-[2-[(cis-2,6-Dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide



[0365] Using the N-[2-(hydroxymethyl)-6-quinolinyl]acetamide obtained in Reference Example 5, the same procedures as those of 1) of Example 6 were conducted to obtain the titled compound as powders.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.02 (6H, d, J = 6.2 Hz), 1.25-1.76 (6H, m), 2.24 (3H, s), 2.50-2.70 (2H, m), 4.01 (2H, s), 7.52 (1H, dd, J = 2.2 and 8.8 Hz), 7.73 (1H, br), 7.83 (1H, d, J = 8.8 Hz), 7.94 (1H, d, J = 8.4 Hz), 8.04 (1H, d, J = 8.4 Hz), 8.29 (1H, d, J = 2.2 Hz).




Elemental analysis for $C_{19}H_{25}N_3O \cdot 0.5H_2O$			
Calcd	C, 71.22;	H, 8.18;	N, 13.11
Found	C, 71.01;	H, 7.81;	N, 12.90

m.p.: 120-122°C (crystallization solvent: ethyl acetate-hexane)

#### Reference Example 12


#### [0366] 2-(Diethylaminoethyl)-6-quinolineamine



[0367] Using the N-[2-(hydroxymethyl)-6-quinoliny]acetamide obtained in Reference Example 5, the same procedures as those of 1) of Example 6 and Reference Example 9 were conducted to obtain the titled compound as a powder.  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 1.06 (6H, t, J = 7.0 Hz), 2.60 (4H, q, J = 7.0 Hz), 3.82 (2H, s), 3.91 (2H, br), 6.90 (1H, d, J = 2.6 Hz), 7.12 (1H, dd, J = 2.6 and 8.8 Hz), 7.54 (1H, d, J = 8.4 Hz), 7.86 (2H, d, J = 8.8 Hz). m.p.: 93-94°C (crystallization solvent: diethyl ether-hexane)

#### Reference Example 13


#### [0368] N-[2-[(2,2,6,6-Tetramethyl-1-piperidinyl)methyl]-6-quinoliny]acetamide



[0369] Using the N-[2-(hydroxymethyl)-6-quinoliny]acetamide obtained in Reference Example 5, the same procedures as those of 1) of Example 6 were conducted to obtain the titled compound as powders.  $^1H$ -NMR ( $CDCl_3$ )  $\delta$ : 1.03 (12H, m), 1.50-1.73 (6H, m), 2.24 (3H, s), 4.07 (2H, s), 7.40-7.57 (2H, m), 7.93 (1H, d, J=8.8Hz), 8.00 (1H, d, J=8.8Hz), 8.06 (1H, d, J=8.8Hz), 8.27 (1H, d, J=2.4Hz).

#### Reference Example 14

#### [0370] 2-(4-Chlorophenyl)-5-carboxy-1,3-dioxane



1) A solution of p-chlorobenzaldehyde (3.2 g, 22.7 mmol), diethyl bis(hydroxymethyl)malonate (5.0 g, 22.7 mmol) and p-toluenesulfonic acid monohydrate (0.44 g, 2.3 mmol) in toluene (70 ml) was heated to reflux for 2 hours in a 200 ml eggplant-shaped flask equipped with the Dean-Stark dehydrating apparatus. After the reaction solution was cooled, 100 ml of ethyl acetate was added thereto, the mixture was washed successively with a 1N aqueous sodium hydroxide solution (50 ml), water (50 ml) and an aqueous saturated sodium chloride solution (50 ml), and

dried over magnesium sulfate. After the solvent was distilled off under reduced pressure, the residue was subjected to silica gel column chromatography (developing solvent: hexane-ethyl acetate=4/1) to obtain 2-(4-chlorophenyl)-5,5'-dicarboethoxy-1,3-dioxane (5.9g, 76%) as colorless crystals.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.27(3H, t, J=7.2Hz), 1.31(3H, t, J=7.2Hz), 4.13(2H, dd like), 4.20(2H, q, J=7.2Hz), 4.33(2H, q, J=7.2Hz), 4.85(2H, dd like), 5.46(1H, s), 7.32(2H, d, J=10.2Hz), 7.38(2H, d, J=10.2Hz).  
m.p.: 54-55°C

2) The 2-(4-chlorophenyl)-5,5'-dicarboethoxy-1,3-dioxane (5.8 g, 16.9 mmol) obtained in the above 1) was dissolved in 60 ml of 90% ethanol, potassium hydroxide (3.8 g, 67.7 mmol) was added thereto, and the mixture was heated to reflux for 3 hours. After the solvent was distilled off under reduced pressure, the resulting solid was suspended in diethyl ether (300 ml), and pH was adjusted to 2 with 2N hydrochloric acid under ice-cooling. The organic layer was separated, washed with an aqueous saturated sodium chloride solution (50 ml), and dried over magnesium sulfate. The solvent was distilled off under reduced pressure to obtain 2-(4-chlorophenyl)-5,5'-dicarboxy-1,3-dioxane (4.3 g, 82%) as a yellow solid.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.99(2H, d, J=12.2Hz), 4.50(2H, dd, J=4.6Hz, 12.2Hz), 5.42(1H, s), 7.35(2H, d, J=9.0Hz), 7.42(2H, d, J=9.0Hz).  
m.p.: 164-165°C


3) A mixture of the 2-(4-chlorophenyl)-5,5'-dicarboxy-1,3-dioxane (4.3 g, 15 mmol) obtained in the above 2) and triethylamine (20 ml) was heated at 150°C for 4 hours. After the reaction solution was concentrated under reduced pressure, the residue was dissolved in diethyl ether (200 ml), and pH was adjusted to 2 with 2N hydrochloric acid. The organic layer was washed with an aqueous saturated sodium chloride solution, dried over magnesium sulfate, and the solvent was distilled off under reduced pressure. The resulting solid was washed with a hexane-ethyl acetate solution to obtain the titled compound (3.09 g, 85%) as a pale yellow powder.

FAB(pos): 243[M+H]<sup>+</sup>

m.p.: 183-184°C

#### Reference Example 15

##### [0371] 6-(1-Piperidinylmethyl)naphthalene-2-amine




[0372] The tert-butyl 6-(1-piperidinylmethyl)-2-naphthylcarbamate (710 mg, 2.09 mmol) obtained in Reference Example 3 was dissolved in trifluoroacetic acid (10 ml), and the solution was stirred at room temperature for 1 hour. The reaction solution was concentrated under reduced pressure, ethyl acetate (50 ml) was added to the residue, the mixture was washed with an aqueous potassium carbonate solution (50 ml) and an aqueous saturated sodium chloride solution (50 ml), and concentrated under reduced pressure to obtain the titled compound (420 mg, 1.75 mmol) as pale orange crystals.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.38 (2H, m), 1.48 (4H, m), 2.32 (4H, br s), 3.44 (2H, s), 5.30 (2H, br s), 6.79 (1H, s), 6.90 (1H, dd, J=8.5 and 2.0Hz), 7.23 (1H, d, J=8.3Hz), 7.43(1H, d, J=8.5Hz), 7.46 (1H, s), 7.54 (1H, d, J=8.5Hz).

#### Reference Example 16

##### [0373] Tert-butyl 6-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-2-naphthylcarbamate




[0374] Using the tert-butyl 6-(hydroxymethyl)-2-naphthylcarbamate obtained in 2) of Reference Example 1, the same procedures as those of Reference Example 2 were conducted to obtain the titled compound as a pale yellow oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.08 (3H, s), 1.10 (3H, s), 1.34 (3H, m), 1.55 (12H, m), 2.53 (2H, m), 3.92 (2H, s), 6.58 (1H, brs), 7.30 (1H, dd, J=2.2 and 8.8Hz), 7.45 (1H, dd, J=1.5 and 8.5Hz), 7.71 (3H, m), 7.94 (1H, brs).

## Reference Example 17

[0375] 6-[(cis-2,6-Dimethyl-1-piperidinyl)methyl]naphthalene-2-amine




[0376] Using the tert-butyl 6-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-2-naphthylcarbamate obtained in Reference Example 16, the same procedures as those of Reference Example 15 were conducted to obtain the titled compound as a pale yellow powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.11 (3H, s), 1.13 (3H, s), 1.32 (3H, m), 1.56 (3H, m), 2.52 (2H, m), 3.78 (1H, brs), 3.91 (2H, s), 6.92 (1H, dd, J=2.4 and 8.7Hz), 6.96 (1H, m), 7.38 (1H, dd, J=1.5 and 8.3Hz), 7.51 (1H, d, J=8.3Hz), 7.63 (1H, d, J=8.5Hz), 7.68 (1H, brs).

## Reference Example 18

[0377] 6-(1-Pyrrolidinylmethyl)naphthalene-2-amine




[0378] Using the tert-butyl 6-(1-pyrrolidinylmethyl)-1-naphthylcarbamate obtained in Reference Example 2, the same procedures as those of Reference Example 15 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.81 (4H, m), 2.56 (4H, m), 3.74 (2H, s), 3.80 (2H, br), 6.93 (2H, dd, J = 2.1, 8.4 Hz), 6.97 (1H, d, J = 2.1 Hz), 7.38 (1H, dd, J = 2.1, 8.4 Hz), 7.55 (1H, d, J = 8.4 Hz), 7.61-7.64 (2H, m).

## Reference Example 19

[0379] 6-[(2,2,6,6-Tetramethyl-1-piperidinyl)methyl]-2-naphthaleneamine




[0380] Using the tert-butyl 6-(hydroxymethyl)-2-naphthylcarbamate obtained in 2) of Reference Example 1, the same procedures as those of Reference Example 2 and Reference Example 15 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.02 (12H, s), 1.57 (6H, m), 3.78 (2H, br), 3.89 (2H, s), 6.89-6.98 (2H, m), 7.41-7.54 (2H, m), 7.63 (1H, d, J = 8.4 Hz), 7.76 (1H, s).

## Reference Example 20

[0381] N-[6-[(Diisopropylamino)methyl]-2-naphthyl]-2-hydroxy-2-methylpropanamide



1) To a solution of 6-(hydroxymethyl)-2-naphthol (500 mg, 2.87 mmol) in dimethylacetamide (4 ml) was added sodium hydroxide (344 mg, 8.61 mmol), and the mixture was stirred for 1 hour. 2-bromo-2-methylpropanamide (1.43 g, 8.61 mmol) and potassium iodide (476 mg, 2.87 mmol) were added thereto, and the mixture was stirred at room temperature for 16 hours. To the reaction solution was added water, and the mixture was extracted with ethyl acetate. The extract was washed with a 1N aqueous sodium hydroxide solution and an aqueous saturated sodium chloride solution, and dried over anhydrous sodium sulfate. After the solvent was concentrated under reduced pressure, 2-[(6-(hydroxymethyl)-2-naphthyl)oxy]-2-methylpropanamide (506mg) was obtained as a colorless powder from isopropyl ether.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.61 (6H, s), 1.85 (1H, t, J = 6.0 Hz), 4.84 (2H, d, J = 6.0 Hz), 5.57 (1H, br), 6.68 (1H, br), 7.16 (1H, dd, J = 2.4, 8.7 Hz), 7.28 (1H, d, J = 2.7 Hz), 7.47 (1H, dd, J = 1.8, 8.7 Hz), 7.72-7.78 (3H, m).


2) To a solution of the 2-[(6-(hydroxymethyl)-2-naphthyl)oxy]-2-methylpropanamide (200 mg, 0.771 mmol) obtained in the above 1) in dimethylformamide (2.3 ml) and 1,3-dimethyl-3,4,5,6-tetrahydro-2(1H)-pyrimidinone (0.23 ml) was added sodium hydride (68 mg, 1.70 mmol), and the mixture was stirred at 100°C for 1 hour. The reaction solution was diluted with ethyl acetate, washed with an aqueous saturated sodium chloride solution, and dried over anhydrous sodium sulfate. After the solvent was concentrated under reduced pressure, the resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate) to obtain 2-hydroxy-N-[6-(hydroxymethyl)-2-naphthyl]-2-methylpropanamide (50 mg) as a colorless powder from isopropylether:hexane=1:1.

3) To a solution of the 2-hydroxy-N-[6-(hydroxymethyl)-2-naphthyl]-2-methylpropanamide (100 mg, 0.386 mmol) obtained in the above 2) and carbon tetrabromide (192 mg, 0.578 mmol) in dichloromethane (2.3 ml) was added triphenylphosphine (121 mg, 0.463 mmol), and the mixture was stirred at room temperature for 4 hours. The reaction solution was purified by alumina column chromatography (developing solvent: dichloromethane) to obtain an oil. To the resulting oil was added diisopropylamine (3 ml), and the mixture was stirred at 80°C for 16 hours. The mixture was dissolved in 1N hydrochloric acid, washed with diethyl ether, and potassium carbonate was added to the aqueous layer to adjust to basic. This was extracted with ethyl acetate, washed with an aqueous saturated sodium chloride solution, and dried over anhydrous sodium sulfate. After the solvent was concentrated under reduced pressure, the resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and converted into a powder with hexane to obtain the titled compound (58 mg).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (12H, d, J = 6.6 Hz), 1.60 (6H, s), 2.24 (1H, s), 3.05 (2H, m), 3.76 (2H, s), 7.50 (2H, m), 7.73 (3H, m), 8.25 (1H, d, J = 2.2 Hz), 8.79 (1H, s).

#### Reference Example 21


[0382] 1-[(6-Methoxy-2-naphthyl)methyl]pyrrolidine



[0383] To a solution of 6-methoxy-2-naphthaldehyde (3.00 g, 16.1 mmol) and pyrrolidine (2.69ml, 32.2mmol) in tetrahydrofuran (32ml) and acetic acid (16ml) was added sodium triacetoxyhydroborate (6.63g, 32.2mmol) at 0°C, and the mixture was stirred at room temperature for 7 hours. The solvent was distilled off under reduced pressure, 1N hydrochloric acid was added to the resulting oil, and the mixture was washed with diethyl ether. An 8N aqueous sodium hydroxide solution was added to the aqueous layer, followed by extraction with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, and dried over anhydrous sodium sulfate. The solvent was concentrated under reduced pressure to obtain the titled compound (3.89 g).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.80 (4H, m), 2.54 (4H, m), 3.74 (2H, s), 3.92 (3H, s), 7.13 (2H, m), 7.45 (1H, m), 7.68-7.73 (3H, m). Reference Example 22

[0384] 6-(1-Pyrrolidinylmethyl)-2-naphthol hydrobromide




[0385] A solution of the 1-[(6-methoxy-2-naphthyl)methyl]pyrrolidine (3.63 g, 15.0 mmol) obtained in Reference Example 21 in 48% hydrobromic acid (75 ml) was stirred at 100°C for 6 hours. The reaction solution was diluted with water, the produced crystals were collected, and washed with water, tetrahydrofuran and diisopropyl ether to obtain the titled compound (2.45 g) as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>, free base) δ: 1.80 (4H, m), 2.71 (4H, m), 3.77 (2H, s), 5.20 (1H, br), 6.78 (1H, d, J = 2.6 Hz), 7.85 (1H, dd, J = 2.6, 8.8 Hz), 7.31 (2H, s-like), 8.43 (1H, d, J = 8.8 Hz), 8.56 (1H, s).

#### Reference Example 23

[0386] 2-Methyl-1,2,3,4-tetrahydrobenzo[b][1,4]naphthyridine-8-amine



1) To a solution of 2-amino-5-nitrobenzaldehyde (1.00 g, 6.02 mmol) and 1-methyl-4-piperidinone (0.89 ml, 7.22 mmol) in ethanol (108 ml) was added a 4N aqueous sodium hydroxide solution (9.0 ml) at room temperature, and the mixture was stirred at 60°C for 1 hour. The solvent was distilled off under reduced pressure, ethyl acetate was added to the resulting oil, the mixture was washed with an aqueous saturated sodium chloride solution, and dried over anhydrous sodium sulfate. After the solvent was concentrated under reduced pressure, the resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and converted into powders with diisopropyl ether to obtain 2-methyl-8-nitro-1,2,3,4-tetrahydrobenzo[b][1,6]naphthyridine (430 mg).


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.55 (3H, s), 2.92 (2H, t, J = 6.0 Hz), 3.32 (2H, t, J = 6.0 Hz), 3.82 (2H, s), 7.96 (1H, s), 8.10 (1H, d, J = 9.0 Hz), 8.40 (1H, dd, J = 2.7, 9.0 Hz), 8.71 (1H, d, J = 2.7 Hz).

2) A suspension of the 2-methyl-8-nitro-1,2,3,4-tetrahydrobenzo[b][1,6]naphthyridine (413 mg, 1.70 mmol) obtained in the above 1) and 10% palladium-carbon in methanol (9.5 ml) was stirred for 2 hours in the hydrogen atmosphere. After the catalyst was filtered, the filtrate was concentrated under reduced pressure, and converted into powders with diisopropyl ether to obtain the titled compound (315 mg).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.51 (3H, s), 2.86 (2H, t, J = 6.0 Hz), 3.20 (2H, t, J = 6.0 Hz), 3.73 (2H, s), 3.88 (2H, br), 6.83 (1H, d, J = 2.4 Hz), 7.09 (1H, dd, J = 2.4, 9.0 Hz), 7.53 (1H, s), 7.80 (1H, d, J = 9.0 Hz).

#### Reference Example 24

[0387] [5-[(4-Bromobenzyl)oxy]-1H-indol-2-yl]methanol



1) A solution of ethyl 5-hydroxy-1H-indole-2-carboxylate (508 mg, 2.48 mmol), 1-bromo-4-(bromomethyl)benzene (681 mg, 2.72 mmol) and potassium carbonate (684 mg, 4.95 mmol) in acetonitrile (12 ml) was stirred at 80°C for 3 hours. The solvent was distilled off under reduced pressure, ethyl acetate was added to the resulting oil, the mixture was washed with an aqueous saturated sodium chloride solution, and dried over anhydrous sodium sulfate. After the solvent was concentrated under reduced pressure, the resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and converted into powders with isopropyl ether to obtain ethyl 5-[(4-bromobenzyl)oxy]-1H-indole-2-carboxylate (565 mg).


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.33 (3H, t, J = 6.9 Hz), 4.32 (2H, t, J = 6.9 Hz), 5.08 (2H, s), 6.98-7.04 (2H, m), 7.18 (1H, d, J = 2.4 Hz), 7.36 (1H, d, J = 8.7 Hz), 7.43 (2H, d, J = 8.4 Hz), 7.59 (2H, d, J = 8.4 Hz), 11.77 (1H, s).

2) To a solution of the ethyl 5-[(4-bromobenzyl)oxy]-1H-indole-2-carboxylate (300 mg, 0.802 mmol) obtained in the above 1) in tetrahydrofuran (4 ml) was added lithium aluminum hydride (60.8 mg, 1.60 mmol) at 0°C, and the mixture was stirred for 1 hour. Ethyl acetate was added to the reaction solution, the mixture was washed with 1N hydrochloric acid and an aqueous saturated sodium chloride solution, and dried over anhydrous sodium sulfate. After the solvent was concentrated under reduced pressure, the resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and converted into powders with isopropyl ether and hexane to obtain the titled compound (219 mg).

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 4.55 (2H, d, J = 8.4 Hz), 5.05 (2H, s), 5.19 (1H, t, J = 8.4 Hz), 6.16 (1H, s), 6.74 (1H, dd, J = 8.8, 2.2 Hz), 7.03 (1H, d, J = 2.2 Hz), 7.20 (1H, d, J = 8.8 Hz), 7.41 (2H, d, J = 8.4 Hz), 7.57 (2H, d, J = 8.4 Hz), 10.84 (1H, s).

#### Example 1

[0388] 4'-Chloro-N-[6-[(N,N-dimethylamino)methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide



[0389] The tert-butyl 6-[(N,N-dimethylamino)methyl]-2-naphthylcarbamate (237 mg, 0.789 mmol) obtained in Reference Example 1 was dissolved in trifluoroacetic acid (4 ml), the solution was stirred at room temperature for 1 hour, and the solvent was concentrated under reduced pressure. To the residue were added ethyl acetate and an aqueous potassium carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate and concentrated under reduced pressure. The resulting residue, 4'-chlorobiphenylcarboxylic acid (184 mg, 0.789 mmol) and dimethylaminopyridine (96.4 mg, 0.789 mmol) were dissolved in N,N-dimethylformamide (4 ml), 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (151 mg, 0.789 mmol) was added thereto under ice-cooling, and the mixture was stirred at room temperature for 16 hours. To the reaction solution were added ethyl acetate and an aqueous potassium carbonate solution, the mixture was extracted, the organic layer was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and the insolubles were crystallized from a mixed solution of ethyl acetate and diisopropyl ether (1.5) to obtain the titled compound (207 mg) as a white powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.19 (6H, s), 3.53 (2H, s), 7.45 (1H, d, J = 8.4 Hz), 7.58 (2H, d, J = 8.4 Hz), 7.73 (1H, s), 7.85 (7H, m), 8.12 (2H, d, J = 8.4 Hz), 8.45 (1H, s), 10.49 (1H, s).

FAB(pos): 415.2 [M+H]<sup>+</sup>

m.p.: 230-231°C

#### Example 2

[0390] 4'-Chloro-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl][1,1'-biphenyl]-4-carboxamide



[0391] Using the tert-butyl 6-(1-pyrrolidinylmethyl)-2-naphthylcarbamate (387 mg, 1.19 mmol) obtained in Reference

Example 2 and 4'-chlorobiphenylcarboxylic acid (112 mg, 0.482 mmol), the same procedures as those of Example 1 were conducted to obtain the titled compound (212 mg) as a white powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.71 (4H, m), 2.47 (4H, m), 3.71 (2H, s), 7.46 (1H, d, J = 8.4 Hz), 7.57 (2H, d, J = 8.4 Hz), 7.75-7.89 (8H, m), 8.12 (2H, d, J = 8.7 Hz), 8.45 (1H, s), 10.48 (1H, s).

FAB(pos): 441.1[M+H]<sup>+</sup>

m.p.: 214-217°C (crystallization solvent: ethyl acetate-diisopropyl ether)

### Example 3

[0392] 4'-Chloro-N-[6-(1-piperidinylmethyl)-2-naphthyl][1,1'-biphenyl]-4-carboxamide




[0393] Using the tert-butyl 6-(1-piperidinylmethyl)-2-naphthylcarbamate (100 mg, 0.42 mmol) obtained in Reference Example 3 and 4'-chlorobiphenylcarboxylic acid (116 mg, 0.49 mmol), the same procedures as those of Example 1 were conducted to obtain the titled compound (103 mg) as a white powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.41-1.51 (6H, m), 2.37 (4H, m), 3.56 (2H, s), 7.46 (1H, d, J = 8.4 Hz), 7.58 (2H, d, J = 8.7 Hz), 7.72 (1H, s), 7.80 (7H, m), 8.12 (2H, d, J = 8.7 Hz), 8.45 (1H, s), 10.48 (1H, s).

m.p.: 220-222°C (crystallization solvent: ethyl acetate-diisopropyl ether)

### Example 4

[0394] N-(4-Bromophenyl)-6-[(dimethylamino)methyl]-2-naphthamide




[0395] The N-(4-bromophenyl)-6-(hydroxymethyl)-2-naphthamide (349 mg, 0.980 mmol) obtained in Reference Example 4 and triethylamine (0.164 ml, 1.18 mmol) were dissolved in N,N-dimethylformamide (5 ml), methanesulfonyl chloride (0.091 ml, 18 mmol) was added thereto under ice-cooling, and the mixture was stirred for 30 minutes. To the reaction solution were added dimethylamine hydrochloride (160 mg, 96 mmol) and potassium carbonate (406 mg, 2.94 mmol), and the mixture was stirred at 60°C for 16 hours. To the reaction solution were added ethyl acetate and water, and the mixture was extracted with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. To the residue was added diisopropyl ether, followed by crystallization to obtain the titled compound (135 mg) as a white powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.19 (6H, s), 3.57 (2H, s), 7.57 (3H, m), 7.81 (2H, d, J = 9.0 Hz), 7.87 (1H, s), 8.00 (3H, m), 8.54 (1H, s), 10.52 (1H, s).

### Example 5

[0396] N-(4'-Chloro[1,1'-biphenyl]-4-yl)-6-[(N,N-dimethylamino)methyl]-2-naphthamide




[0397] N-(4-Bromophenyl)-6-[(dimethylamino)methyl]-2-naphthamide (128 mg, 0.334 mmol) obtained in Example 4, 4-chlorophenylboronic acid (62.7 mg, 401 mmol) and a 2N aqueous sodium carbonate solution (0.668 ml, 1.34 mmol) were dissolved in a mixed solution of dimethoxyethane (3 ml) and tetrahydrofuran (0.3 ml), tetrakis(triphenylphosphine)palladium (11.6 mg, 0.01 mol) was added thereto under the nitrogen atmosphere, and the mixture was stirred at 90°C for 4 hours. To the reaction solution was added an aqueous saturated sodium chloride solution, and the mixture was extracted with ethyl acetate. The extract was dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and the insolubles were crystallized from a mixed solution of ethyl acetate and diisopropyl ether (1:3) to obtain the titled compound (42 mg) as a white powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.21 (6H, s), 3.59 (2H, s), 7.51 (2H, d, J = 8.7 Hz), 7.59 (1H, d, J = 8.1 Hz), 7.72 (4H, m), 7.89 (1H, s), 7.95 (2H, d, J = 8.7 Hz), 8.05 (3H, m), 8.58 (1H, s), 10.53 (1H, s).  
m.p.: 240-242°C

#### Example 6

[0398] 4'-Chloro-N-[2-[(N,N-dimethylamino)methyl]-6-quinolinyl]-[1,1'-biphenyl]-4-carboxamide



1) The N-[2-(hydroxymethyl)-6-quinolinyl]acetamide (92.3 mg, 0.427 mmol) obtained in Reference Example 5 and triethylamine (0.0712 ml, 0.512 mmol) were dissolved in N,N-dimethylformamide (2 ml), methanesulfonyl chloride (0.0396 ml, 0.512 mmol) was added under ice-cooling, and the mixture was stirred for 30 minutes. To the reaction solution were added dimethylamine hydrochloride (69.6 mg, 0.854 mmol) and potassium carbonate (177 mg, 1.28 mmol), and the mixture was stirred at 60°C for 16 hours. To the reaction solution was added an aqueous potassium carbonate solution, followed by extraction with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate) to obtain N-[2-[(N,N-dimethylamino)methyl]-6-quinolinyl]acetamide (71.9 mg) as an oil.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.25 (3H, s), 2.34 (6H, s), 3.76 (2H, s), 7.45-7.58 (3H, m), 8.02 (1H, d, J = 9.0 Hz), 8.09 (1H, d, J = 8.4 Hz), 8.31 (1H, s).

2) The N-[2-[(N,N-dimethylamino)methyl]-6-quinolinyl]acetamide (1.9 mg, 0.296 mmol) obtained in the above 1) was dissolved in concentrated hydrochloric acid (1.5 ml), and the solution was stirred at 110°C for 2 hours. The solvent was distilled off under reduced pressure, an aqueous potassium carbonate solution was added to the residue, and the mixture was extracted with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue, 4'-chlorobiphenylcarboxylic acid (68.8 mg, 0.296 mmol) and dimethylaminopyridine (36.1 mg, 0.296 mmol) were dissolved in N,N-dimethylformamide (1.5 ml), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide hydrochloride (56.6 mg, 0.296 mmol) was added under ice-cooling, and the mixture was stirred at room temperature for 16 hours. To the reaction solution was added an aqueous potassium carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by



alumina column chromatography (developing solvent: ethyl acetate), and crystallized from ethyl acetate-diisopropyl ether (1:5) to obtain the titled compound (60.6mg) as a white powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.23 (6H, s), 3.67 (2H, s), 7.58 (3H, m), 7.82 (2H, d, J = 8.4 Hz), 7.89 (2H, d, J = 8.4 Hz), 7.96 (1H, d, J = 9.0 Hz), 8.04 (1H, dd, J = 9.0, 2.1 Hz), 8.13 (2H, d, J = 8.4 Hz), 8.29 (2H, d, J = 8.4 Hz), 8.52 (1H, d, J = 2.1 Hz), 10.60 (1H, s).

FAB(pos): 416.1[M+H]<sup>+</sup>

m.p.: 219-221°C

#### Example 7

[0399] 4'-Fluoro-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide



1) Using the N-[2-(hydroxymethyl)-6-quinolinyl]acetamide obtained in Reference Example 5 and pyrrolidine, the same procedures as those of 1) of Example 6 were conducted to obtain N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide as a white powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.85 (4H, m), 2.24 (3H, s), 2.70 (4H, m), 3.99 (2H, s), 7.58 (2H, m), 7.80 (1H, s), 7.98 (1H, d, J = 9.0 Hz), 8.07 (1H, d, J = 8.4 Hz), 8.29 (1H, s).

2) Using the N-[2-(hydroxymethyl)-6-quinolinyl]acetamide obtained in the above 1) and 4'-fluorobiphenylcarboxylic acid, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a white powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.76 (4H, m), 2.61 (4H, m), 3.94 (2H, s), 7.36 (2H, m), 7.59 (1H, d, J = 8.4 Hz), 7.87 (4H, m), 7.99-8.14 (4H, m), 8.30 (1H, d, J = 8.4 Hz), 8.54 (1H, d, J = 2.0 Hz), 10.61 (1H, s).

FAB(pos): 426.1[M+H]<sup>+</sup>

m.p.: 190-193°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 8

[0400] 4'-Chloro-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide



[0401] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7 and 4'-chlorobiphenylcarboxylic acid, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a white powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.77 (4H, m), 2.65 (4H, m), 3.98 (2H, s), 7.59 (3H, m), 7.59 (1H, d, J = 8.4 Hz), 7.87 (4H, m), 7.99-8.14 (4H, m), 8.30 (1H, d, J = 8.4 Hz), 8.54 (1H, d, J = 2.0 Hz), 10.61 (1H, s).

FAB(pos): 442.1[M+H]<sup>+</sup>

m.p.: 200-202°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 9

[0402] 4'-Fluoro-N-[2-(1-piperidinylmethyl)-6-quinolinyl]acetamide



1) Using the N-[2-(hydroxymethyl)-6-quinolinyl]acetamide obtained in Reference Example 5 and piperidine, the same procedures as those of 1) of Example 6 were conducted to obtain N-[2-(1-piperidinylmethyl)-6-quinolinyl]acetamide as a white powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.47 (2H, m), 1.64 (4H, m), 2.25 (3H, s), 2.55 (4H, m), 3.83 (2H, s), 7.55 (1H, dd, J = 2.4, 9.0 Hz), 7.60 (1H, s), 7.67 (1H, d, J = 8.4 Hz), 7.98 (1H, d, J = 9.0 Hz), 8.08 (1H, d, J = 8.4 Hz), 8.30 (1H, d, J = 2.4 Hz).

2) Using the N-[2-(1-piperidinylmethyl)-6-quinolinyl]acetamide obtained in the above 1) and 4'-fluorobiphenylcarboxylic acid, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a white powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.44 (2H, m), 1.56 (4H, m), 2.50 (4H, m), 3.77 (2H, s), 7.35 (2H, m), 7.60 (1H, d, J = 8.7 Hz), 7.81-8.14 (8H, m), 8.30 (1H, d, J = 8.4 Hz), 8.54 (1H, d, J = 2.0 Hz), 10.61 (1H, s).

FAB(pos): 440.2[M+H]<sup>+</sup>

m.p.: 202-204°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 10

[0403] 4'-Chloro-N-[2-(1-piperidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0404] Using the N-[2-(1-piperidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 9 and 4'-chlorobiphenylcarboxylic acid, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a white powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.47 (2H, m), 1.61 (4H, m), 2.50 (4H, m), 3.33 (2H, s), 7.61 (3H, m), 7.81-7.90 (4H, m), 7.98-8.15 (4H, m), 8.34 (1H, d, J = 8.4 Hz), 8.57 (1H, s), 10.65 (1H, s).

FAB(pos): 456.1[M+H]<sup>+</sup>

m.p.: 211-213°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 11

[0405] 4'-Chloro-N-[2-(4-morpholinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



1) Using the N-[2-(hydroxymethyl)-6-quinolinyl]acetamide obtained in Reference Example 5 and piperidine, the same procedures as those of 1) of Example 6 were conducted to obtain N-[2-(4-morpholinylmethyl)-6-quinolinyl]

acetamide as a powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 2.25 (3H, s), 2.55 (4H, t, J = 4.5 Hz), 3.75 (4H, t, J = 4.5 Hz), 3.81 (2H, s), 7.42 (1H, br), 7.51 (1H, dd, J = 2.4, 9.0 Hz), 7.60 (1H, d, J = 8.4 Hz), 8.01 (1H, d, J = 9.0 Hz), 8.09 (1H, d, J = 8.4 Hz), 8.32 (1H, d, J = 2.4 Hz).

2) Using the N-[2-(4-morpholinylmethyl)-6-quinolinyl]acetamide obtained in the above 1), the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.45 (4H, t, J = 4.5 Hz), 3.61 (4H, t, J = 4.5 Hz), 3.74 (2H, s), 7.59 (3H, m), 7.82 (2H, d, J = 8.4 Hz), 7.88 (2H, d, J = 8.4 Hz), 7.97 (1H, d, J = 9.0), 8.04 (1H, dd, J = 9.0, 2.1 Hz), 8.13 (2H, d, J = 8.4 Hz), 8.30 (1H, d, J = 8.4 Hz), 8.52 (1H, d, J = 2.1 Hz), 10.60 (1H, s).

Elemental analysis for C <sub>27</sub> H <sub>24</sub> ClN <sub>3</sub> O <sub>2</sub>			
Calcd	C, 70.81;	H, 5.28;	N, 9.18
Found	C, 70.66;	H, 5.31;	N, 8.90

m.p.: 236-238°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 12

[0406] N-[2-(4-Morpholinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0407] Using the N-[2-(4-morpholinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.47 (4H, t, J = 4.5 Hz), 3.61 (4H, t, J = 4.5 Hz), 3.74 (2H, s), 7.44 (1H, m), 7.53 (2H, m), 7.61 (1H, d, J = 8.4 Hz), 7.78 (2H, d, J = 7.5), 7.88 (2H, d, J = 8.4 Hz), 7.97 (1H, d, J = 9.0 Hz), 8.04 (1H, dd, J = 9.0, 2.1 Hz), 8.12 (1H, d, J = 8.4 Hz), 8.30 (1H, d, J = 8.4 Hz), 8.53 (1H, d, J = 2.1 Hz), 10.59 (1H, s).

Elemental analysis for C <sub>27</sub> H <sub>26</sub> N <sub>3</sub> O <sub>2</sub> ·0.5H <sub>2</sub> O			
Calcd	C, 74.98;	H, 6.06;	N, 9.72
Found	C, 75.08;	H, 6.07;	N, 9.80

m.p.: 214-215°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 13

[0408] 4'-Fluoro-N-[2-(4-morpholinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0409] Using the N-[2-(4-morpholinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.45 (4H, t, J = 4.5 Hz), 3.61 (4H, t, J = 4.5 Hz), 3.74 (2H, s), 7.35 (2H, m), 7.61 (1H, d, J = 8.4 Hz), 7.84 (4H, m), 7.96 (1H, d, J = 9.0 Hz), 8.04 (1H, dd, J = 9.0, 2.4 Hz), 8.12 (2H, d, J = 8.1 Hz), 8.30 (1H, d, J = 8.4 Hz), 8.53 (1H, d, J = 2.4 Hz), 10.59 (1H, s).


Elemental analysis for C<sub>27</sub>H<sub>24</sub>FN<sub>3</sub>O<sub>2</sub>

Calcd	C, 73.45;	H, 5.48;	N, 9.52
Found	C, 73.37;	H, 5.36;	N, 9.52

m.p.: 211-212°C (crystallization solvent: ethyl acetate-diisopropyl ether)

Example 14

[0410] 6-(4-Methylphenyl)-N-[2-(4-morpholinylmethyl)-6-quinolinyl]nicotinamide



[0411] Using the N-[2-(4-morpholinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.39 (3H, s), 2.45 (4H, t, J = 4.5 Hz), 3.61 (4H, t, J = 4.5 Hz), 3.75 (2H, s), 7.36 (2H, d, J = 8.4 Hz), 7.62 (1H, d, J = 8.4 Hz), 8.01 (2H, m), 8.12 (3H, m), 8.31 (1H, d, J = 8.4 Hz), 8.42 (1H, dd, J = 8.4, 2.4 Hz), 8.52 (1H, d, J = 2.4 Hz), 9.23 (1H, d, J = 2.4 Hz), 10.72 (1H, s).


Elemental analysis for C<sub>27</sub>H<sub>26</sub>N<sub>4</sub>O<sub>2</sub>·0.5H<sub>2</sub>O

Calcd	C, 73.95;	H, 5.98;	N, 12.78
Found	C, 73.92;	H, 5.92;	N, 13.01

m.p.: 214-216°C (crystallization solvent: ethyl acetate-diisopropyl ether)

Example 15

[0412] N-[2-[(Dimethylamino)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0413] Using the N-[2-[(N,N-dimethylamino)methyl]-6-quinolinyl]acetamide obtained in 1) of Example 6, and biphenylcarboxylic acid, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.23 (6H, s), 3.66 (2H, s), 7.52 (4H, m), 7.78 (2H, d, J = 7.8 Hz), 7.88 (2H, d, J = 8.4 Hz), 7.96 (1H, d, J = 9.0 Hz), 8.04 (1H, dd, J = 9.0, 2.1 Hz), 8.12 (2H, d, J = 8.1 Hz), 8.29 (1H, d, J = 8.4 Hz), 8.53 (1H, d, J = 2.1 Hz), 10.59 (1H, s).

Elemental analysis for $C_{25}H_{23}N_3O$			
Calcd	C, 78.71;	H, 6.08;	N, 11.02
Found	C, 78.44;	H, 6.07;	N, 11.01

m.p.: 191-194°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 16

[0414] N-[2-[(Dimethylamino)methyl]-6-quinolinyl]-4'-fluoro[1,1'-biphenyl]-4-carboxamide



[0415] Using the N-[2-[(N,N-dimethylamino)methyl]-6-quinolinyl]acetamide obtained in 1) of Example 6, and 4'-fluorobiphenylcarboxylic acid, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


$^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 2.23 (6H, s), 3.67 (2H, s), 7.35 (2H, m), 7.58 (1H, d,  $J = 8.4$  Hz), 7.84 (4H, m), 7.96 (1H, d,  $J = 9.0$  Hz), 8.04 (1H, dd,  $J = 9.0, 2.3$  Hz), 8.12 (2H, d,  $J = 8.4$  Hz), 8.28 (1H, d,  $J = 8.7$  Hz), 8.52 (1H, d,  $J = 2.3$  Hz), 10.58 (1H, s).

Elemental analysis for $C_{25}H_{22}FN_3O$			
Calcd	C, 75.17;	H, 5.55;	N, 10.52
Found	C, 74.89;	H, 5.60;	N, 10.52

m.p.: 205-208°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 17

[0416] N-[2-[(Dimethylamino)methyl]-6-quinolinyl]-6-(4-methylphenyl)nicotinamide



[0417] Using the N-[2-[(N,N-dimethylamino)methyl]-6-quinolinyl]acetamide obtained in 1) of Example 6, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

$^1\text{H-NMR}$  ( $\text{DMSO-}d_6$ )  $\delta$ : 2.23 (6H, s), 2.39 (3H, s), 3.67 (2H, s), 7.36 (2H, d,  $J = 8.1$  Hz), 7.59 (1H, d,  $J = 8.4$  Hz), 8.02 (2H, m), 8.12 (3H, m), 8.30 (1H, d,  $J = 8.7$  Hz), 8.42 (1H, dd,  $J = 8.4, 2.4$  Hz), 8.51 (1H, s), 9.22 (1H, d,  $J = 2.4$  Hz), 10.72 (1H, s).

Elemental analysis for $C_{25}H_{24}N_4O$			
Calcd	C, 75.73;	H, 6.10;	N, 14.13


(continued)

Elemental analysis for $C_{25}H_{24}N_4O$			
Found	C, 75.44;	H, 6.19;	N, 14.12

m.p.: 220-222°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 18

[0418] 4'-Chloro-N-[2-[(2,5-dimethyl-1-pyrrolidiny)methyl]-6-quinoliny]methyl]-6-quinoliny]-4-carboxamide



1) Using the N-[2-(hydroxymethyl)-6-quinoliny]acetamide obtained in Reference Example 5 and piperidine, the same procedures as those of 1) of Example 6 were conducted to obtain N-[2-[(2,5-dimethyl-1-pyrrolidiny)methyl]-6-quinoliny]acetamide as a powder.


$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 0.99 (6H, d, 6.0 Hz), 1.86 (4H, m), 2.24 (3H, s), 2.72 (2H, m), 3.98 (2H, s), 7.53 (1H, dd, J = 2.2, 9.0 Hz), 7.69 (1H, d, J = 8.4 Hz), 7.80 (1H, br), 7.97 (1H, d, J = 9.0 Hz), 8.03 (1H, d, J = 8.4 Hz), 8.29 (1H, s).

2) Using the N-[2-[(2,5-dimethyl-1-pyrrolidiny)methyl]-6-quinoliny]acetamide obtained in the above 1), the same procedures as those of 1) of Example 6 were conducted to obtain the titled compound as a colorless powder.

FAB(pos): 470.2[M+H]<sup>+</sup>

## Example 19

[0419] N-[2-[(2,5-Dimethyl-1-pyrrolidiny)methyl]-6-quinoliny]methyl]-4'-fluoro[1,1'-biphenyl]-4-carboxamide




[0420] Using the N-[2-[(2,5-dimethyl-1-pyrrolidiny)methyl]-6-quinoliny]acetamide obtained in 1) of Example 18, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

FAB(pos): 454.2[M+H]<sup>+</sup>

## Example 20


[0421] N-[2-[(2,5-Dimethyl-1-pyrrolidiny)methyl]-6-quinoliny]methyl]-4-carboxamide



[0422] Using the N-[2-((2,5-dimethyl-1-pyrrolidinyl)methyl)-6-quinolinyl]acetamide obtained in 1) of Example 18, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder. FAB(pos): 436.2[M+H]<sup>+</sup>

#### Example 21

[0423] 4-(4-Methyl-1-piperidiny)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]benzamide




[0424] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 0.98 (3H, d, J = 6.4 Hz), 1.33 (2H, m), 1.59 (1H, m), 1.74 (2H, m), 1.80 (4H, m), 2.62 (4H, m), 2.86 (2H, m), 3.85 (2H, m), 3.94 (2H, s), 6.95 (2H, d, J = 9.0 Hz), 7.57 (1H, d, J = 8.7 Hz), 7.65 (1H, dd, J = 2.4, 8.7 Hz), 7.81 (2H, d, J = 9.0 Hz), 7.90 (1H, br), 8.05 (1H, d, J = 8.7 Hz), 8.10 (1H, d, J = 8.7 Hz), 8.45 (1H, d, J = 2.4 Hz). FAB(pos): 429.3[M+H]<sup>+</sup>

m.p.: 200-202°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 22

[0425] 4-(2-Oxo-1-piperidiny)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]benzamide



[0426] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.84 (4H, m), 1.98 (4H, m), 2.62 (6H, m), 3.70 (2H, m), 3.94 (2H, s), 7.36 (2H, d, J = 8.4 Hz), 7.59 (1H, d, J = 8.4 Hz), 7.68 (1H, dd, J = 2.3, 9.0 Hz), 7.90 (2H, d, J = 8.4 Hz), 8.06 (1H, d, J = 9.0 Hz), 8.11 (1H, d, J = 8.4 Hz), 8.30 (1H, br), 8.45 (1H, d, J = 2.3 Hz).

FAB(pos): 429.2[M+H]<sup>+</sup>

m.p.: 210-212°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 23

[0427] 4'-Chloro-N-[2-[[[(2R,6S)-2,6-dimethyl-1-piperidinyl]methyl]-6-quinolinyl]]-1,1'-biphenyl]-4-carboxamide




[0428] A solution of the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl]]-1,1'-biphenyl]-4-carboxamide (100 mg, 0.246 mmol) obtained in Reference Example 8, (2R,6S)-2,6-dimethylpiperidine (0.331 ml, 2.46 mmol) and potassium carbonate (67.9 mg, 0.491 mmol) in dimethylformamide (1.5 ml) was stirred at 80°C for 3 hours. The reaction solution was concentrated under reduced pressure, and water was added thereto. The resulting precipitates were collected, and washed successively with water, ethanol and isopropyl ether to obtain the titled compound (65 mg) as a powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.01 (6H, d, J = 6.0 Hz), 1.29 (2H, m), 1.61 (4H, m), 2.52 (2H, m), 3.96 (2H, s), 7.59 (2H, d, J = 8.4 Hz), 7.75 (1H, d, J = 8.4 Hz), 7.83 (2H, d, J = 8.4 Hz), 7.90 (2H, d, J = 8.4 Hz), 8.00 (2H, m), 8.14 (2H, d, J = 8.4 Hz), 8.27 (1H, d, J = 8.4 Hz), 8.53 (1H, s), 10.61 (1H, s).

FAB(pos): 484[M+H]<sup>+</sup>

## Example 24

[0429] 4'-Chloro-N-[2-[(4-methyl-1-piperazinyl)methyl]-6-quinolinyl]]-1,1'-biphenyl]-4-carboxamide



[0430] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl]]-1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.16 (3H, s), 2.35-2.50 (8H, m), 3.73 (2H, s), 7.58 (3H, m), 7.82 (2H, d, J = 8.4 Hz), 7.89 (2H, d, J = 8.4 Hz), 7.96 (1H, d, J = 9.0 Hz), 8.04 (1H, m), 8.13 (2H, d, J = 8.4 Hz), 8.29 (1H, d, J = 8.7 Hz), 8.53 (1H, s), 10.61 (1H, s).


FAB(pos): 471.2[M+H]<sup>+</sup>

m.p.: 215°C (decomposition) (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 25

[0431] 4'-Chloro-N-[2-[(2-methyl-4,5-dihydro-1H-imidazol-1-yl)methyl]-6-quinolinyl]]-1,1'-biphenyl]-4-carboxamide






[0432] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as colorless a powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.97 (3H, s), 3.21 (2H, t, J = 9.0 Hz), 3.52 (2H, t, J = 9.0 Hz), 4.56 (2H, s), 7.46 (1H, d, J = 8.4 Hz), 7.58 (2H, d, J = 8.4 Hz), 7.82 (2H, d, J = 8.4 Hz), 7.89 (2H, d, J = 8.4 Hz), 7.97 (1H, d, J = 9.0 Hz), 8.07 (1H, dd, J = 9.0, 2.1 Hz), 8.13 (2H, d, J = 8.4 Hz), 8.34 (1H, d, J = 8.4 Hz), 8.56 (1H, d, J = 2.1 Hz), 10.63 (1H, s).  
FAB(pos): 455[M+H]<sup>+</sup>

#### Example 26

[0433] 4'-Chloro-N-[2-(1,3-thiazolidin-3-ylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0434] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.94 (2H, t, J = 6.0 Hz), 3.12 (2H, t, J = 6.0 Hz), 3.78 (2H, s), 4.09 (2H, s), 7.58 (2H, d, J = 8.6 Hz), 7.70 (1H, d, J = 8.0 Hz), 7.83 (2H, d, J = 8.4 Hz), 7.89 (2H, d, J = 8.4 Hz), 8.00 (2H, m), 8.14 (2H, d, J = 8.6 Hz), 8.34 (1H, d, J = 8.8 Hz), 8.56 (1H, s), 10.63 (1H, s).  
FAB(pos): 460 [M+H]<sup>+</sup>

#### Example 27

[0435] 4'-Chloro-N-[2-[(2,2,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0436] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.02 (12H, s), 1.56 (6H, m), 4.01 (2H, s), 7.59 (2H, d, J = 8.4 Hz), 7.90 (6H, m), 7.99 (1H, m), 8.13 (2H, d, J = 8.0 Hz), 8.27 (1H, d, J = 8.8 Hz), 8.51 (1H, s), 10.59 (1H, s).  
FAB(pos): 512 [M+H]<sup>+</sup>

# Example 28

[0437] 4'-Chloro-N-[2-[[4-(2-pyridinyl)-1-piperazinyl]methyl]-6-quinolinyl]-1,1'-biphenyl]-4-carboxamide




[0438] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.36 (4H, m), 3.31 (4H, m), 3.60 (2H, s), 6.43 (1H, m), 6.61 (1H, d, J = 8.8 Hz), 7.32-7.48 (4H, m), 7.60-7.71 (4H, m), 7.75-7.95 (5H, m), 8.11 (1H, d, J = 8.4 Hz), 8.36 (1H, s), 10.42 (1H, s).  
FAB(pos): 534 [M+H]<sup>+</sup>

# Example 29

[0439] 4'-Chloro-N-[2-[[4-(2-methylphenyl)-1-piperidinyl]methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0440] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.71 (4H, m), 2.23 (3H, m), 2.31 (3H, s), 2.97 (2H, m), 3.80 (2H, s), 7.08-7.24 (4H, m), 7.59 (2H, d, J = 8.4 Hz), 7.67 (1H, d, J = 8.8 Hz), 7.83 (2H, d, J = 8.4 Hz), 7.90 (2H, d, J = 8.4 Hz), 8.03 (2H, m), 8.14 (2H, d, J = 8.4 Hz), 8.32 (1H, d, J = 8.4 Hz), 8.55 (1H, s), 10.63 (1H, s).  
FAB(pos): 546 [M+H]<sup>+</sup>

# Example 30

[0441] 4'-Chloro-N-[2-[[4-(3-methylphenyl)-1-piperidinyl]methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0442] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.71 (4H, m), 2.20 (3H, m), 2.27 (3H, s), 2.97 (2H, m), 3.57 (2H, s), 7.06 (4H, m), 7.59 (3H, m), 7.82 (4H, m), 8.02 (2H, m), 8.12 (2H, d, J = 8.4 Hz), 8.35 (1H, m), 8.54 (1H, s), 10.61 (1H, s).  
FAB(pos): 546 [M+H]<sup>+</sup>

#### Example 31

[0443] 4'-Chloro-N-[2-[(4-(4-methylphenyl)-1-piperidinyl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0444] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.71 (4H, m), 2.20 (3H, m), 2.26 (3H, s), 2.96 (2H, m), 3.78 (2H, s), 7.12 (4H, m), 7.61 (3H, m), 7.83 (2H, d, J = 8.6 Hz), 7.90 (2H, d, J = 8.6 Hz), 8.03 (2H, m), 8.14 (2H, d, J = 8.2 Hz), 8.31 (1H, d, J = 8.4 Hz), 8.55 (1H, s), 10.63 (1H, s).  
FAB(pos): 546 [M+H]<sup>+</sup>

#### Example 32

[0445] 4'-Chloro-N-[2-[(2-phenyl-4,5-dihydro-1H-imidazol-1-yl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0446] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 3.46 (2H, m), 3.82 (2H, m), 4.51 (2H, s), 7.48 (3H, m), 7.61 (4H, s), 7.79-8.16 (9H, m), 8.36 (1H, d, J = 8.8 Hz), 8.58 (1H, d, J = 1.8 Hz), 10.65 (1H, s).  
FAB(pos): 517 [M+H]<sup>+</sup>

#### Example 33

[0447] 4'-Chloro-N-[2-(3,4-dihydro-1(2H)-quinolinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0448] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.00 (2H, m), 2.79 (2H, m), 3.55 (2H, m), 4.70 (2H, s), 6.47 (2H, m), 6.82-6.94 (2H, m), 7.38 (1H, d, J = 8.4 Hz), 7.58 (2H, d, J = 8.4 Hz), 7.83 (2H, d, J = 8.4 Hz), 7.89 (2H, d, J = 8.4 Hz), 8.01 (2H, m), 8.13 (2H, d, J = 8.4 Hz), 8.27 (1H, d, J = 8.4 Hz), 8.55 (1H, d, J = 1.8 Hz), 10.63 (1H, s).

FAB(pos): 504[M+H]<sup>+</sup>

#### Example 34

[0449] 4'-Chloro-N-[2-(3,4-dihydro-2(1H)-isoquinolinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0450] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.77-2.83 (4H, m), 3.64 (2H, s), 3.93 (2H, s), 7.03-7.11 (4H, m), 7.57 (2H, d, J = 8.8 Hz), 7.64 (1H, d, J = 8.4 Hz), 7.82 (2H, d, J = 8.8 Hz), 7.88 (2H, d, J = 8.4 Hz), 8.01 (2H, m), 8.12 (2H, d, J = 8.4 Hz), 8.30 (1H, d, J = 8.4 Hz), 8.54 (1H, s), 10.62 (1H, s).

FAB(pos): 504[M+H]<sup>+</sup>

#### Example 35

[0451] 4'-Chloro-N-[2-(2,3-dihydro-1H-indol-1-ylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0452] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.96 (2H, m), 3.43 (2H, m), 4.53 (2H, s), 6.61 (2H, m), 6.95-7.09 (2H, m), 7.57 (3H, m), 7.83 (2H, d, J = 8.8 Hz), 7.89 (2H, d, J = 8.8 Hz), 8.02 (2H, m), 8.13 (2H, d, J = 8.4 Hz), 8.32 (1H, d, J = 8.8 Hz), 8.55 (1H,

d, J = 2.2 Hz), 10.63 (1H, s).  
FAB(pos):490 [M+H]<sup>+</sup>

# Example 36

[0453] 4'-Chloro-N-[2-(1H-imidazol-1-ylmethyl)-6-quinolinyl]-6-quinolinyl]-4-carboxamide




[0454] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl]-[1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 5.49 (2H, s), 6.96 (1H, s), 7.28 (2H, m), 7.58 (3H, m), 7.82 (2H, d, J = 8.4 Hz), 7.89 (2H, d, J = 8.4 Hz), 8.01 (2H, m), 8.12 (2H, d, J = 8.6 Hz), 8.35 (1H, d, J = 8.4 Hz), 8.57 (1H, d, J = 2.2 Hz), 10.65 (1H, s).  
FAB(pos):439 [M+H]<sup>+</sup>

# Example 37

[0455] 4'-Chloro-N-[2-[(4-phenyl-1-piperazinyl)methyl]-6-quinolinyl]-[1,1'-biphenyl]-4-carboxamide




[0456] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl]-[1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.62 (4H, m), 3.17 (4H, m), 3.82 (2H, s), 6.78 (1H, m), 6.93 (2H, d, J = 7.8 Hz), 7.20 (2H, m), 7.58 (2H, d, J = 8.8 Hz), 7.69 (1H, d, J = 8.8 Hz), 7.83 (2H, d, J = 8.8 Hz), 7.89 (2H, d, J = 8.6 Hz), 8.01 (2H, m), 8.14 (2H, d, J = 8.6 Hz), 8.32 (1H, d, J = 8.6 Hz), 8.56 (1H, d, J = 1.8 Hz), 10.63 (1H, s).  
FAB(pos):533 [M+H]<sup>+</sup>

# Example 38

[0457] 4-(4-Fluorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]-1-piperidinecarboxamide



[0458] To a solution of the 2-(1-pyrrolidinylmethyl)-6-quinolinylamine (500 mg, 2.2 mmol) obtained in Reference Example 9 and pyridine (0.356 ml, 4.4 mmol) in tetrahydrofuran (11 ml) was added 4-nitrophenyl chloroformate (488 mg, 2.42 mmol) under ice-cooling. After stirred for 30 minutes, the reaction solution was concentrated, and dimethyl sulfoxide (11 ml) was added to the residue. 4-(4-Fluorophenyl)piperidine hydrochloride (569 mg, 2.64 mmol) and a 4N aqueous sodium hydroxide solution (0.66 ml) were added thereto at room temperature while stirring, and the mixture was stirred for 2 hours. Ethyl acetate and water were added, the mixture was extracted, the organic layer was washed with water, and concentrated, and the residue was purified by alumina column chromatography (developing solvent: ethyl acetate) to obtain the titled compound (612 mg) as a colorless powder from ethyl acetate-diisopropyl ether.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.57 (2H, m), 1.64 (4H, m), 1.82 (2H, m), 2.50 (4H, m), 2.79 (1H, m), 2.92 (2H, m), 3.81 (2H, s), 4.34 (2H, m), 7.12 (2H, m), 7.31 (2H, m), 7.51 (1H, d, J = 8.4 Hz), 7.82 (2H, s-like), 8.07 (1H, s), 8.14 (1H, d, J = 8.4 Hz), 8.83 (1H, s).

FAB(pos):433.2 [M+H]<sup>+</sup>

m.p.: 206-207°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 39

[0459] 4-Phenyl-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]-1-piperidinecarboxamide



[0460] Using the 2-(1-pyrrolidinylmethyl)-6-quinolinylamine obtained in Reference Example 9, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.59-1.85 (8H, m), 2.50 (4H, m), 2.73 (1H, m), 2.89 (2H, m), 3.81 (2H, s), 4.33 (2H, m), 7.20-7.34 (5H, m), 7.51 (1H, d, J = 8.4 Hz), 7.82 (2H, s-like), 8.07 (1H, s), 8.14 (1H, d, J = 8.4 Hz), 8.83 (1H, s).

FAB(pos):415.3 [M+H]<sup>+</sup>

m.p.: 187-189°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 40

[0461] 4-(4-Chlorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]-1-piperidinecarboxamide



[0462] Using the 2-(1-pyrrolidinylmethyl)-6-quinolinylamine obtained in Reference Example 9, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.58 (2H, m), 1.72 (4H, m), 1.80 (2H, m), 2.50 (4H, m), 2.79 (1H, m), 2.92 (2H, m), 3.82 (2H, s), 4.33 (2H, m), 7.34 (4H, m), 7.51 (1H, d, J = 8.4 Hz), 7.82 (2H, s-like), 8.07 (1H, s), 8.14 (1H, d, J = 8.4 Hz), 8.85 (1H, s).

FAB(pos):449.9 [M+H]<sup>+</sup>

m.p.: 205°C (decomposition) (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 41

[0463] 4-(4-Methylphenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]-1-piperidinecarboxamide



[0464] Using the 2-(1-pyrrolidinylmethyl)-6-quinolinylamine obtained in Reference Example 9, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.55 (2H, m), 1.72 (4H, m), 1.78 (2H, m), 2.27 (3H, s), 2.50 (4H, m), 2.72 (1H, m), 2.91 (2H, m), 3.81 (2H, s), 4.32 (2H, m), 7.13 (4H, m), 7.51 (1H, d, J = 8.4 Hz), 7.82 (2H, s-like), 8.08 (1H, s), 8.14 (1H, d, J = 8.4 Hz), 8.84 (1H, s).

FAB(pos): 429.3 [M+H]<sup>+</sup>

m.p.: 214-216°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 42

[0465] 6-(4-Methoxyphenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]nicotinamide



[0466] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 2.50 (4H, m), 3.85 (5H, s-like), 7.10 (2H, d, J = 9.0 Hz), 7.59 (1H, d, J = 8.7 Hz), 7.98 (2H, m), 8.11 (1H, d, J = 8.4 Hz), 8.18 (2H, d, J = 8.7 Hz), 8.29 (1H, d, J = 9.0 Hz), 8.40 (1H, d, J = 8.4 Hz), 8.51 (1H, s), 9.20 (1H, s), 10.71 (1H, s).

FAB(pos): 439.2 [M+H]<sup>+</sup>

m.p.: 210°C (decomposition) (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 43

[0467] N-[2-(1-Pyrrolidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0468] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 2.50 (4H, m), 3.85 (2H, s), 7.50 (4H, m), 7.78 (2H, d, J = 8.1 Hz), 7.88 (2H, d, J = 8.1 Hz), 8.02 (2H, m), 8.12 (2H, d, J = 8.1 Hz), 8.28 (1H, d, J = 8.7 Hz), 8.53 (1H, d, J = 2.0 Hz), 10.60 (1H, s).

FAB(pos): 408.2 [M+H]<sup>+</sup>

m.p.: 181-183°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 44

[0469] 6-(4-Chlorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]nicotinamide



[0470] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 2.50 (4H, m), 3.85 (2H, s), 7.61 (3H, m), 8.01 (2H, m), 8.28 (4H, m), 8.48 (2H, m), 9.25 (1H, d, J = 2.2 Hz), 10.76 (1H, s).

FAB(pos): 443.2 [M+H]<sup>+</sup>

m.p.: 225-227°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 45

[0471] 6-(4-Fluorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]nicotinamide



[0472] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 2.50 (4H, m), 3.86 (2H, s), 7.39 (2H, m), 7.60 (1H, d, J = 8.4 Hz), 8.01 (2H, m), 8.19 (1H, d, J = 8.4 Hz), 8.28 (3H, m), 8.45 (1H, dd, J = 8.4, 2.1 Hz), 8.51 (1H, d, J = 1.7 Hz), 9.24 (1H, d, J = 2.0 Hz), 10.75 (1H, s).

FAB(pos): 427.2 [M+H]<sup>+</sup>

m.p.: 210°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 46

[0473] 6-(4-Methylphenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]nicotinamide



[0474] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 2.39 (3H, s), 2.50 (4H, m), 3.86 (2H, s), 7.36 (2H, d, J = 8.1 Hz), 7.59 (1H, d, J =




8.4 Hz), 8.01 (2H, m), 8.13 (3H, m), 8.29 (1H, d, J = 8.4 Hz), 8.42 (1H, dd, J = 8.4, 2.2 Hz), 8.51 (1H, d, J = 2.0 Hz), 9.22 (1H, d, J = 2.2 Hz), 10.73 (1H, s).

FAB(pos): 423.2[M+H]<sup>+</sup>

m.p.: 207-209°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 47

[0475] 4'-Chloro-N-[2-[[4-(4-fluorobenzoyl)-1-piperidinyl]methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0476] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.76 (4H, m), 2.28 (2H, m), 2.89 (2H, m), 3.40 (1H, m), 3.79 (2H, s), 7.37 (2H, m), 7.62 (3H, m), 7.83 (2H, d, J = 8.4 Hz), 7.90 (2H, d, J = 8.4 Hz), 8.03-8.14 (6H, m), 8.31 (1H, d, J = 8.6 Hz), 8.55 (1H, s), 10.63 (1H, s). FAB(pos): 578[M+H]<sup>+</sup>

#### Example 48

[0477] 4'-Chloro-N-[2-[[4-(4-chlorobenzoyl)-1-piperidinyl]methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0478] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.77 (4H, m), 2.28 (2H, m), 2.90 (2H, m), 3.45 (1H, m), 3.79 (2H, s), 7.62 (5H, m), 7.83 (2H, d, J = 8.4 Hz), 7.90 (2H, d, J = 8.4 Hz), 8.03 (4H, m), 8.14 (2H, d, J = 8.2 Hz), 8.31 (1H, d, J = 8.4 Hz), 8.54 (1H, s), 10.63 (1H, s).

FAB(pos): 594[M+H]<sup>+</sup>

#### Example 49

[0479] 4'-Chloro-N-[2-[(methylanilino)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0480] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 3.17 (3H, s), 4.80 (2H, s), 6.62 (1H, m), 6.76 (2H, d, J = 8.0 Hz), 7.15 (2H, m), 8.30 (1H, d, J = 8.8 Hz), 7.59 (2H, d, J = 8.4 Hz), 7.83 (2H, d, J = 8.4 Hz), 7.90 (2H, d, J = 8.4 Hz), 7.96 (2H, m), 8.13 (2H, d, J = 8.4 Hz), 8.27 (1H, d, J = 8.6 Hz), 8.55 (1H, s), 10.63 (1H, s).

FAB(pos): 478[M+H]<sup>+</sup>

#### Example 50

[0481] 4'-Chloro-N-[2-[[3-(4-fluorobenzoyl)-1-piperidinyl]methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0482] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.39 (1H, s), 1.73 (3H, m), 2.23 (2H, m), 2.91 (2H, m), 3.63 (1H, m), 3.78 (2H, s), 7.31 (2H, m), 7.58 (3H, m), 7.86 (4H, m), 8.01 (4H, m), 8.13 (2H, d, J = 8.4 Hz), 8.25 (1H, d, J = 8.2 Hz), 8.54 (1H, s), 10.61 (1H, s).

FAB(pos): 578[M+H]<sup>+</sup>

#### Example 51

[0483] 4'-Chloro-N-[2-[(4-phenyl)-1-piperidinyl]methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0484] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.74 (4H, m), 2.20 (2H, m), 2.52 (1H, m), 2.97 (2H, m), 3.79 (2H, s), 7.27 (5H, m), 7.58 (2H, d, J = 8.4 Hz), 7.69 (1H, d, J = 8.4 Hz), 7.83 (2H, d, J = 8.8 Hz), 7.89 (2H, d, J = 8.4 Hz), 8.00 (2H, m), 8.14 (2H, d, J =

8.8 Hz), 8.31 (1H, d, J = 8.4 Hz), 8.54 (1H, d, J = 1.8 Hz), 10.62 (1H, s).  
FAB(pos): 532[M+H]<sup>+</sup>

# Example 52

[0485] 1-(4-Chlorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]-4-piperidinecarboxamide



[0486] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 1.89 (4H, m), 2.50 (5H, m), 2.75 (2H, m), 3.76 (2H, m), 3.83 (2H, s), 6.99 (2H, d, J = 9.0 Hz), 7.24 (2H, d, J = 9.0 Hz), 7.55 (1H, d, J = 8.8 Hz), 7.77 (1H, dd, J = 8.8, 2.2 Hz), 7.90 (1H, d, J = 8.8 Hz), 8.21 (1H, d, J = 8.8 Hz), 8.38 (1H, d, J = 2.2 Hz), 10.26 (1H, s).

FAB(pos): 449[M+H]<sup>+</sup>

m.p.: 204-206°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 53

[0487] N-[2-[(2-Benzyl-4,5-dihydro-1H-imidazol-1-yl)methyl]-6-quinolinyl]-4'-chloro[1,1'-biphenyl]-4-carboxamide




[0488] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 3.22 (2H, m), 3.60 (2H, m), 3.75 (2H, s), 4.51 (2H, s), 7.17-7.33 (6H, m), 7.57 (2H, d, J = 8.4 Hz), 7.82 (2H, d, J = 8.4 Hz), 7.88 (2H, d, J = 8.4 Hz), 7.99 (2H, m), 8.12 (2H, d, J = 8.4 Hz), 8.25 (1H, d, J = 9.0 Hz), 8.53 (1H, s), 10.62 (1H, s).

FAB(pos): 531[M+H]<sup>+</sup>

# Example 54

[0489] N-[2-(1-Piperidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



1) Using the N-[2-(hydroxymethyl)-6-quinolinyl]acetamide obtained in Reference Example 5 and piperidine, the same procedures as those of 1) of Example 6 were conducted to obtain N-[2-(1-piperidinylmethyl)-6-quinolinyl]acetamide.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.38-1.70 (6H, m), 2.24 (3H, s), 2.38-2.53 (4H, m), 3.76 (2H, s), 7.52 (1H, dd,  $J = 2.6, 9.2$  Hz), 7.63 (1H, d,  $J = 8.4$  Hz), 7.67 (1H, br), 7.99 (1H, d,  $J = 9.2$  Hz), 8.07 (1H, d,  $J = 8.4$  Hz), 8.31 (1H, d,  $J = 2.6$  Hz).

Elemental analysis for  $\text{C}_{17}\text{H}_{21}\text{N}_3\text{O} \cdot 0.25\text{H}_2\text{O}$

Calcd	C, 70.93;	H, 7.53;	N, 14.60
Found	C, 71.06;	H, 7.37;	N, 14.62

m.p.: 182-184°C (crystallization solvent: ethyl acetate-n-hexane)

2) The N-[2-(1-piperidinylmethyl)-6-quinolinyl]acetamide (4.5 g, 16 mmol) obtained in the above 1) and concentrated hydrochloric acid (70 ml) were stirred at 110°C for 1 hour. The solvent was distilled off under reduced pressure, an aqueous sodium hydroxide solution was added to the residue, and the mixture was extracted with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate) to obtain 6-amino-2-(1-piperidinylmethyl)quinoline (3.4 g) as an oil.

$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.37-1.68 (6H, m), 2.39-2.55 (4H, m), 3.72 (2H, s), 3.91 (2H, br), 6.89 (1H, d,  $J = 2.6$  Hz), 7.12 (1H, dd,  $J = 2.6$  and 8.8 Hz), 7.51 (1H, d,  $J = 8.4$  Hz), 7.86 (1H, d,  $J = 8.4$  Hz), 7.87 (1H, d,  $J = 8.8$  Hz).

3) The 6-amino-2-(1-piperidinylmethyl)quinoline (250 mg, 1 mmol) obtained in the above 2), biphenylcarboxylic acid (220 mg, 1.1 mmol) and dimethylaminopyridine (150 mg, 1.2 mmol) were dissolved in N,N-dimethylformamide (8 ml), WSC (230 mg, 1.2 mmol) was added thereto under ice-cooling, and the mixture was stirred at room temperature for 18 hours. To the reaction solution was added an aqueous potassium carbonate solution, and the mixture was extracted with ethyl acetate. The extract was washed with an aqueous saturated sodium chloride solution, dried over anhydrous sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and crystallized from acetic acid to obtain the titled compound (335mg) having a melting point of 186-188°C as colorless needle crystals.


$^1\text{H-NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 1.38-1.76 (6H, m), 2.43-2.56 (4H, m), 3.78 (2H, s), 7.38-7.56 (3H, m), 7.59-7.81 (6H, m), 7.96-8.18 (5H, m), 8.49 (1H, d,  $J = 2.2$  Hz).

Elemental analysis for  $\text{C}_{28}\text{H}_{27}\text{N}_3\text{O} \cdot 0.5\text{H}_2\text{O}$

Calcd	C, 78.11;	H, 6.56;	N, 9.76
Found	C, 78.48;	H, 6.31;	N, 10.00

#### Example 55

[0490] 4'-Methyl-N-[2-(1-piperidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0491] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.38-1.70 (6H, m), 2.39-2.56 (7H, m), 3.79 (2H, s), 7.30 (2H, d, J = 7.6 Hz), 7.56 (2H, d, J = 7.6 Hz), 7.63-7.78 (4H, m), 7.95-8.07 (5H, m), 8.49 (1H, d, J = 1.8 Hz).


Elemental analysis for C<sub>29</sub>H<sub>29</sub>N<sub>3</sub>O·0.25H<sub>2</sub>O

Calcd	C, 79.15;	H, 6.76;	N, 9.55
Found	C, 79.38;	H, 6.88;	N, 9.73

m.p.: 198-200°C (crystallization solvent: ethyl acetate)

#### Example 56

[0492] 4'-Methoxy-N-[2-(1-piperidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0493] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.38-1.85 (6H, m), 2.41-2.56 (4H, m), 3.78 (2H, s), 3.88 (3H, s), 7.02 (2H, d, J = 8.8 Hz), 7.54-7.66 (6H, m), 7.93-8.20 (5H, m), 8.49 (1H, d, J = 2.2 Hz).


Elemental analysis for C<sub>29</sub>H<sub>29</sub>N<sub>3</sub>O<sub>2</sub>·0.25H<sub>2</sub>O

Calcd	C, 76.38;	H, 6.52;	N, 9.21
Found	C, 76.21;	H, 6.38;	N, 9.32

m.p.: 192-194°C (crystallization solvent: ethyl acetate)

#### Example 57

[0494] 6-(4-Chlorophenyl)-N-[2-(1-piperidinylmethyl)-6-quinolinyl]nicotinamide



[0495] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.38-1.80 (6H, m), 2.38-2.58 (4H, m), 3.78 (2H, s), 7.48 (2H, d, J = 8.4 Hz), 7.62-7.75 (2H, m), 7.84 (1H, d, J = 8.0 Hz), 7.96-8.18 (4H, m), 8.24 (1H, br), 8.31 (1H, dd, J = 2.2, 8.4 Hz), 8.45 (1H, d, J = 2.2 Hz), 9.20 (1H, d, J = 1.4 Hz).


Elemental analysis for  $C_{27}H_{25}ClN_4O \cdot 0.5H_2O$ 

Calcd	C, 69.59;	H, 5.62;	N, 12.02
Found	C, 69.33;	H, 5.52;	N, 12.08

m.p.: 215-218°C (decomposition) (crystallization solvent: ethyl acetate)

## Example 58

[0496] 6-(4-Methylphenyl)-N-[2-(1-piperidinylmethyl)-6-quinolinyl]nicotinamide



[0497] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.38-1.80 (6H, m), 2.40-2.56 (7H, m), 3.79 (2H, s), 7.33 (2H, d, J = 8.4 Hz), 7.63-7.74 (2H, m), 7.86 (1H, d, J = 8.0 Hz), 7.93-8.19 (5H, m), 8.29 (1H, dd, J = 2.2, 8.4 Hz), 8.46 (1H, d, J = 2.2 Hz), 9.19 (1H, d, J = 2.2 Hz).

m.p.: 206-207°C (decomposition) (crystallization solvent: ethyl acetate)

FAB(pos): 437[M+H]<sup>+</sup>

## Example 59

[0498] 6-(4-Fluorophenyl)-N-[2-(1-piperidinylmethyl)-6-quinolinyl]nicotinamide



[0499] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.38-1.74 (6H, m), 2.42-2.56 (4H, m), 3.78 (2H, s), 7.20 (2H, dd, J = 8.4 and 8.8 Hz), 7.63-7.76 (2H, m), 7.84 (1H, d, J = 8.0 Hz), 8.02-8.18 (5H, m), 8.31 (1H, dd, J = 2.2, 8.4 Hz), 8.46 (1H, d, J = 2.2 Hz), 9.19 (1H, d, J = 1.8 Hz).


Elemental analysis for  $C_{27}H_{25}FN_4O \cdot 0.25H_2O$ 

Calcd	C, 72.87;	H, 5.78;	N, 12.59
Found	C, 72.91;	H, 5.45;	N, 12.75

m.p.: 209-210°C (decomposition) (crystallization solvent: ethyl acetate)

## Example 60

[0500] 6-(4-Methoxyphenyl)-N-[2-(1-piperidinylmethyl)-6-quinolinyl]nicotinamide



[0501] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.38-1.71 (6H, m), 2.43-2.56 (4H, m), 3.80 (2H, s), 3.90 (3H, s), 7.04 (2H, d, J = 9.0 Hz), 7.64-7.73 (2H, m), 7.83 (1H, d, J = 8.4 Hz), 8.00-8.18 (5H, m), 8.28 (1H, dd, J = 2.6, 8.4 Hz), 8.46 (1H, d, J = 2.6 Hz), 9.17 (1H, d, J = 1.8 Hz).

Elemental analysis for C <sub>26</sub> H <sub>28</sub> N <sub>4</sub> O <sub>2</sub> ·0.25H <sub>2</sub> O			
Calcd	C, 73.58;	H, 6.28;	N, 12.26
Found	C, 73.56;	H, 6.16;	N, 12.24

m.p.: 210-211°C (decomposition) (crystallization solvent: ethyl acetate)

#### Example 61

[0502] 4'-Methyl-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0503] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 2.37 (3H, s), 2.50 (4H, m), 3.85 (2H, s), 7.33 (2H, d, J = 8.1 Hz), 7.58 (1H, d, J = 8.4 Hz), 7.68 (2H, d, J = 8.1 Hz), 7.85 (2H, d, J = 8.4 Hz), 7.95 (1H, d, J = 9.0 Hz), 8.04 (1H, dd, J = 2.4, 9.0 Hz), 8.10 (2H, d, J = 8.4 Hz), 8.27 (1H, d, J = 8.4 Hz), 8.52 (1H, d, J = 2.4 Hz).

FAB(pos) 422.3[M+H]<sup>+</sup>

m.p.: 192-193°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 62

[0504] 4'-Methoxy-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0505] Using the N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]acetamide obtained in 1) of Example 7, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 2.50 (4H, m), 3.83 (3H, s), 3.85 (2H, s), 7.08 (2H, d, J = 8.7 Hz), 7.58 (1H, d, J = 8.4 Hz), 7.74 (2H, d, J = 8.7 Hz), 7.83 (2H, d, J = 8.4 Hz), 7.95 (1H, d, J = 9.0 Hz), 8.03 (1H, dd, J = 2.1, 9.0 Hz), 8.09 (2H, d, J = 8.4 Hz), 8.27 (1H, d, J = 8.4 Hz), 8.52 (1H, d, J = 2.1 Hz).

FAB(pos) 438.3[M+H]<sup>+</sup>

m.p.: 197-199°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 63

[0506] 4-(4-Chlorophenyl)-N-[2-(1-piperidinylmethyl)-6-quinolinyl]-1-piperidinecarboxamide



[0507] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.37-2.02 (10H, m), 2.38-2.54 (4H, m), 2.63-2.84 (1H, m), 2.94-3.14 (2H, s), 3.76 (2H, s), 4.20-4.36 (2H, m), 6.65 (1H, br), 7.15 (2H, d, J = 8.4 Hz), 7.30 (2H, d, J = 8.4 Hz), 7.49 (1H, dd, J = 2.2 and 9.2 Hz), 7.60 (1H, d, J = 8.4 Hz), 7.93-8.10 (3H, m).

Elemental analysis for C <sub>27</sub> H <sub>31</sub> ClN <sub>4</sub> O·1.5H <sub>2</sub> O			
Calcd	C, 66.18;	H, 6.99;	N, 11.43
Found	C, 66.32;	H, 6.75;	N, 11.74

m.p.: 214-217°C (dec.) (crystallization solvent: ethyl acetate-diethyl ether)

#### Example 64

[0508] 4-(4-Methylphenyl)-N-[2-(1-piperidinylmethyl)-6-quinolinyl]-1-piperidinecarboxamide



[0509] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.40-2.02 (10H, m), 2.34 (3H, s), 2.47-2.83 (5H, m), 2.95-3.15 (2H, m), 3.82 (2H, s), 4.20-4.36 (2H, m), 6.70 (1H, br), 7.14 (4H, br), 7.53 (1H, dd, J = 2.2 and 8.8 Hz), 7.64 (1H, d, J = 8.4 Hz), 7.93-8.10 (3H, m).


Elemental analysis for C <sub>28</sub> H <sub>34</sub> N <sub>4</sub> O·H <sub>2</sub> O			
Calcd	C, 73.01;	H, 7.88;	N, 12.16
Found	C, 72.68;	H, 7.57;	N, 12.20

m.p.: 204-205°C (decomposition) (crystallization solvent: ethyl acetate-diethyl ether)



## Example 65

[0510] 4-(4-Fluorophenyl)-N-[2-(1-piperidinylmethyl)-6-quinolyl]-1-piperidinecarboxamide



[0511] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.36-2.02 (10H, m), 2.37-2.55 (4H, m), 2.63-2.85 (1H, m), 2.94-3.14 (2H, m), 3.76 (2H, s), 4.10-4.36 (2H, m), 6.66 (1H, br), 7.01 (2H, dd, J = 8.4 and 8.8 Hz), 7.11-7.24 (2H, m), 7.50 (1H, dd, J = 2.2 and 8.8 Hz), 7.60 (1H, d, J = 8.4 Hz), 7.92-8.10 (3H, m).

Elemental analysis for C <sub>27</sub> H <sub>31</sub> FN <sub>4</sub> O·0.5H <sub>2</sub> O			
Calcd	C, 71.18;	H, 7.08;	N, 12.30
Found	C, 71.13;	H, 6.94;	N, 12.52

m.p.: 203-204°C (decomposition) (crystallization solvent: ethyl acetate-diethyl ether)

## Example 66

[0512] 4-(4-Methoxyphenyl)-N-[2-(1-piperidinylmethyl)-6-quinolyl]-1-piperidinecarboxamide




[0513] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.37-2.02 (10H, m), 2.42-2.57 (4H, m), 2.60-2.80 (1H, m), 2.94-3.14 (2H, m), 3.78 (2H, s), 3.80 (3H, s), 4.19-4.36 (2H, m), 6.66 (1H, br), 6.87 (2H, d, J = 8.4 Hz), 7.15 (2H, d, J = 8.4 Hz), 7.50 (1H, dd, J = 2.6 and 9.2 Hz), 7.61 (1H, d, J = 8.4 Hz), 7.93-8.10 (3H, m).

m.p.: 197-198°C (decomposition) (crystallization solvent: ethyl acetate-diethyl ether)

## Example 67

[0514] 2'-Fluoro-N-[2-(1-piperidinylmethyl)-6-quinolyl][1,1'-biphenyl]-4-carboxamide




[0515] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.40-1.77 (6H, m), 2.47-2.65 (4H, m), 3.86 (2H, s), 7.12-7.54 (4H, m), 7.63-7.78 (4H, m), 7.96-8.22 (5H, m), 8.48 (1H, d, J = 2.6 Hz).

m.p.: 163-164°C (crystallization solvent: ethyl acetate)

#### Example 68

[0516] 2',4'-Difluoro-N-[2-(1-piperidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0517] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.40-1.78 (6H, m), 2.50-2.67 (4H, m), 3.87 (2H, s), 6.89-7.08 (2H, m), 7.38-7.54 (1H, m), 7.59-7.80 (4H, m), 7.96-8.23 (5H, m), 8.47 (1H, br).


Elemental analysis for C<sub>26</sub>H<sub>25</sub>F<sub>2</sub>N<sub>3</sub>O·0.5H<sub>2</sub>O

Calcd	C, 72.09;	H, 5.62;	N, 9.01
Found	C, 71.79;	H, 5.59;	N, 8.75

m.p.: 181-182°C (crystallization solvent: ethyl acetate)

#### Example 69

[0518] 6-(2,4-Difluorophenyl)-N-[2-(1-piperidinylmethyl)-6-quinolinyl]nicotinamide



[0519] Using the 6-amino-2-(1-piperidinylmethyl)quinoline obtained in 2) of Example 54, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.41-1.77 (6H, m), 2.47-2.68 (4H, m), 3.86 (2H, s), 6.88-7.13 (2H, m), 7.73 (2H, d, J = 8.4 Hz), 7.90-8.00 (1H, m), 8.02-8.25 (4H, m), 8.32 (1H, dd, J = 2.2 and 8.4 Hz), 8.46 (1H, d, J = 1.8 Hz), 9.24 (1H, d, J = 1.8 Hz).


Elemental analysis for C<sub>27</sub>H<sub>24</sub>F<sub>2</sub>N<sub>4</sub>O·0.5H<sub>2</sub>O

Calcd	C, 69.37;	H, 5.39;	N, 11.98
Found	C, 69.14;	H, 5.21;	N, 12.04

m.p.: 182-183°C (crystallization solvent: ethyl acetate)

## Example 70

[0520] 2',4'-Dichloro-N-[2-[(1-piperidinylmethyl)-6-quinolinyl]]-1,1'-biphenyl]-4-carboxamide



[0521] Using the N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.40-1.76 (6H, m), 2.48-2.63 (4H, m), 3.84 (2H, s), 7.22-7.40 (2H, m), 7.52-7.78 (5H, m), 7.96-8.19 (5H, m), 8.47 (1H, d, J = 2.2 Hz).

Elemental analysis for C <sub>28</sub> H <sub>25</sub> Cl <sub>2</sub> N <sub>3</sub> O·0.5H <sub>2</sub> O			
Calcd	C, 67.34;	H, 5.25;	N, 8.41
Found	C, 67.46;	H, 5.20;	N, 8.47

m.p.: 210-212°C (decomposition) (crystallization solvent: ethyl acetate)

## Example 71

[0522] 4'-Fluoro-N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl]]-1,1'-biphenyl]-4-carboxamide




[0523] Using the N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (12H, s), 1.60 (6H, br), 4.10 (2H, s), 7.17 (2H, t, J = 8.8 Hz), 7.56-7.76 (5H, m), 7.96-8.17 (6H, m), 8.46 (1H, d, J = 2.6 Hz).

m.p.: 229-231°C (decomposition) (crystallization solvent: ethyl acetate)

## Example 72

[0524] 6-(4-Chlorophenyl)-N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl]nicotinamide



[0525] Using the N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (12H, s), 1.60 (6H, br), 4.10 (2H, s), 7.49 (2H, d, J = 8.8 Hz), 7.64-7.74 (1H, m), 7.81-7.90 (1H, m), 7.96-8.18 (6H, m), 8.27-8.38 (1H, m), 8.44 (1H, d, J = 2.2 Hz), 9.20 (1H, d, J = 1.8 Hz).

Elemental analysis for C <sub>31</sub> H <sub>33</sub> ClN <sub>4</sub> O·H <sub>2</sub> O				
Calcd	C, 70.11;	H, 6.64;	N, 10.55	
Found	C, 70.16;	H, 6.59;	N, 10.62	

m.p.: 258-259°C (crystallization solvent: ethyl acetate)

#### Example 73

[0526] 6-(4-Methylphenyl)-N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl]nicotinamide



[0527] Using the N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (12H, s), 1.60 (6H, br), 2.44 (3H, s), 4.10 (2H, s), 7.33 (2H, d, J = 8.2 Hz), 7.63-7.74 (1H, m), 7.87 (1H, d, J = 8.4 Hz), 7.94-8.16 (6H, m), 8.25-8.35 (1H, m), 8.44 (1H, d, J = 2.6 Hz), 9.19 (1H, d, J = 1.8 Hz).

Elemental analysis for C <sub>32</sub> H <sub>36</sub> N <sub>4</sub> O·H <sub>2</sub> O				
Calcd	C, 75.26;	H, 7.50;	N, 10.97	
Found	C, 74.98;	H, 7.44;	N, 11.07	

m.p.: 246-247°C (crystallization solvent: ethyl acetate)

#### Example 74

[0528] 4-(4-Chlorophenyl)-N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl]-1-piperidinecarboxamide



[0529] Using the N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 54 and Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.03 (12H, s), 1.46-2.01 (10H, m), 2.63-2.84 (1H, m), 2.95-3.15 (2H, m), 4.07 (2H, s), 4.20-4.37 (2H, m), 6.62 (1H, br), 7.15 (2H, d, J = 8.4 Hz), 7.30 (2H, d, J = 8.4 Hz), 7.49 (1H, dd, J = 2.6 and 8.8 Hz), 7.81-8.11 (4H, m).

FAB(pos): 519[MH]<sup>+</sup>


Elemental analysis for C<sub>31</sub>H<sub>39</sub>ClN<sub>4</sub>O·0.5H<sub>2</sub>O

Calcd	C, 70.50;	H, 7.63;	N, 10.61
Found	C, 70.88;	H, 7.79;	N, 11.14

m.p.: 203-204°C (crystallization solvent: ethyl acetate-diethyl ether)

#### Example 75

[0530] 4-(4-Methylphenyl)-N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl]-1-piperidinecarboxamide



[0531] Using the N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 54 and Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (12H, s), 1.50-2.02 (10H, m), 2.34 (3H, s), 2.63-2.84 (1H, m), 2.94-3.13 (2H, m), 4.07 (2H, s), 4.20-4.34 (2H, m), 6.59 (1H, br), 7.14 (4H, s), 7.48 (1H, dd, J = 2.2 and 9.2 Hz), 7.86-8.09 (4H, m).


Elemental analysis for C<sub>32</sub>H<sub>42</sub>N<sub>4</sub>O·0.5H<sub>2</sub>O

Calcd	C, 75.70;	H, 8.54;	N, 11.04
Found	C, 75.57;	H, 8.30;	N, 11.16

m.p.: 200-202°C (crystallization solvent: ethyl acetate-diethyl ether)

#### Example 76

[0532] N-[2-[(2,2,6,6-Tetramethyl-1-piperidinyl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0533] Using the N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (12H, s), 1.60 (6H, br), 4.10 (2H, s), 7.40-7.56 (3H, m), 7.60-7.80 (6H, m), 7.96-8.16 (5H, m), 8.47 (1H, d, J = 2.4 Hz).

m.p.: 200-201°C (crystallization solvent: ethyl acetate)

#### Example 77

[0534] 4'-Methyl-N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0535] Using the N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (12H, s), 1.59 (6H, br), 2.42 (3H, s), 4.09 (2H, s), 7.30 (2H, d, J = 8.2 Hz), 7.55 (2H, d, J = 8.2 Hz), 7.63-7.77 (4H, m), 7.95-8.16 (5H, m), 8.46 (1H, d, J = 2.6 Hz).


Elemental analysis for C<sub>33</sub>H<sub>37</sub>N<sub>3</sub>O·0.5H<sub>2</sub>O

Calcd	C, 79.16;	H, 7.65;	N, 8.39
Found	C, 79.21,	H, 7.66;	N, 8.41

m.p.: 242-243°C (crystallization solvent: ethyl acetate)

#### Example 78

[0536] 4'-Methoxy-N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0537] Using the N-[2-((2,2,6,6-tetramethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 13, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (12H, s), 1.59 (6H, br), 3.88 (3H, s), 4.10 (2H, s), 7.02 (2H, d, J = 8.8 Hz), 7.55-7.76 (5H, m), 7.94-8.16 (6H, m), 8.46 (1H, d, J = 2.6 Hz).  
m.p.: 210-211°C (crystallization solvent: ethyl acetate)

#### Example 79

[0538] 4'-Fluoro-N-[2-((cis-2,6-dimethyl-1-piperidinyl)methyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0539] Using the N-[2-((cis-2,6-dimethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (6H, d, J = 6.2 Hz), 1.28-1.80 (6H, m), 2.51-2.73 (2H, m), 4.04 (2H, s), 7.17 (2H, t, J = 8.8 Hz), 7.54-7.76 (5H, m), 7.86 (1H, d, J = 8.4 Hz), 7.96-8.17 (5H, m), 8.46 (1H, d, J = 2.2 Hz).


Elemental analysis for C<sub>30</sub>H<sub>30</sub>FN<sub>3</sub>O·0.5H<sub>2</sub>O

Calcd	C, 75.60;	H, 6.56;	N, 8.82
Found	C, 75.72;	H, 6.31;	N, 8.72

m.p.: 187-188°C (crystallization solvent: ethyl acetate)

#### Example 80

[0540] N-[2-((cis-2,6-Dimethyl-1-piperidinyl)methyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0541] Using the N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (6H, d, J = 6.2 Hz), 1.28-1.77 (6H, m), 2.52-2.72 (2H, m), 4.04 (2H, s), 7.40-7.57 (3H, m), 7.60-7.80 (5H, m), 7.86 (1H, d, J = 8.4 Hz), 7.96-8.16 (5H, m), 8.47 (1H, d, J = 2.2 Hz).

Elemental analysis for C <sub>30</sub> H <sub>31</sub> N <sub>3</sub> O·0.5H <sub>2</sub> O			
Calcd	C, 78.57;	H, 7.03;	N, 9.16
Found	C, 78.36,	H, 6.64;	N, 9.12

m.p.: 180-181°C (crystallization solvent: ethyl acetate)

#### Example 81

[0542] 4'-Methyl-N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0543] Using the N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (6H, d, J = 6.2 Hz), 1.28-1.80 (6H, m), 2.43 (3H, s), 2.52-2.73 (2H, m), 4.04 (2H, s), 7.30 (2H, d, J = 8.2 Hz), 7.55 (2H, d, J = 8.2 Hz), 7.63-7.78 (3H, m), 7.85 (1H, d, J = 8.4 Hz), 7.94-8.17 (5H, m), 8.46 (1H, d, J = 2.2 Hz).

Elemental analysis for C <sub>31</sub> H <sub>33</sub> N <sub>3</sub> O·0.5H <sub>2</sub> O			
Calcd	C, 78.78;	H, 7.25;	N, 8.89
Found	C, 78.87;	H, 7.08;	N, 8.82

m.p.: 198-199°C (dec.) (crystallization solvent: ethyl acetate)

#### Example 82

[0544] 4'-Methoxy-N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0545] Using the N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.



<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (6H, d, J = 6.0 Hz), 1.23-1.85 (6H, m), 2.50-2.72 (2H, m), 3.87 (3H, s), 4.03 (2H, s), 7.01 (2H, d, J = 8.8 Hz), 7.50-7.77 (5H, m), 7.85 (1H, d, J = 8.4 Hz), 7.90-8.20 (5H, m), 8.46 (1H, br).  
m.p.: 194-196°C (dec.) (crystallization solvent: ethyl acetate)

## Example 83

[0546] 6-(4-Chlorophenyl)-N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]nicotinamide



[0547] Using the N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (6H, d, J = 6.2 Hz), 1.27-1.76 (6H, m), 2.51-2.76 (2H, m), 4.03 (2H, s), 7.47 (2H, d, J = 8.4 Hz), 7.70 (1H, dd, J = 2.2 and 8.8 Hz), 7.76-7.92 (2H, m), 7.94-8.13 (4H, m), 8.23-8.36 (2H, m), 8.42 (1H, d, J = 2.2 Hz), 9.19 (1H, d, J = 2.2 Hz).


Elemental analysis for C<sub>29</sub>H<sub>29</sub>ClN<sub>4</sub>O·0.5H<sub>2</sub>O

Calcd	C, 70.50;	H, 6.12;	N, 11.34
Found	C, 70.58;	H, 6.06;	N, 11.14

m.p.: 217-219° (dec.) (crystallization solvent: ethyl acetate)

## Example 84

[0548] 6-(4-Methylphenyl)-N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]nicotinamide




[0549] Using the N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (6H, d, J = 6.2 Hz), 1.28-1.80 (6H, m), 2.43 (3H, s), 2.51-2.72 (2H, m), 4.04 (2H, s), 7.32 (2H, d, J = 8.0 Hz), 7.69 (1H, dd, J = 2.2 and 8.8 Hz), 7.80-8.20 (7H, m), 8.28 (1H, dd, J = 2.2 and 8.4 Hz), 8.43 (1H, d, J = 2.2 Hz), 9.18 (1H, d, J = 2.2 Hz).

m.p.: 225-226°C (dec.) (crystallization solvent: ethyl acetate)

## Example 85

[0550] 4-(4-Chlorophenyl)-N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]-1-piperidinecarboxamide




[0551] Using the N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 54 and Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (6H, d, J = 6.2 Hz), 1.23-2.02 (10H, m), 2.50-2.83 (3H, m), 2.94-3.15 (2H, m), 4.02 (2H, s), 4.20-4.37 (2H, m), 6.64 (1H, br), 7.15 (2H, d, J = 8.4 Hz), 7.30 (2H, d, J = 8.4 Hz), 7.50 (1H, dd, J = 2.2 and 8.8 Hz), 7.79 (1H, d, J = 8.8 Hz), 7.88-8.08 (3H, m).  
m.p.: 196-198°C (dec.) (crystallization solvent: ethyl acetate-diethyl ether)

#### Example 86

[0552] 4-(4-Methylphenyl)-N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]-1-piperidinecarboxamide



[0553] Using the N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 54 and Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (6H, d, J = 6.0 Hz), 1.20-2.03 (10H, m), 2.33 (3H, s), 2.50-2.82 (3H, m), 2.90-3.15 (2H, m), 4.02 (2H, s), 4.17-4.39 (2H, m), 6.70 (1H, br), 7.13 (4H, br), 7.51 (1H, d, J = 8.8 Hz), 7.78 (1H, d, J = 8.8 Hz), 7.95-8.15 (3H, m).


Elemental analysis for C<sub>30</sub>H<sub>38</sub>N<sub>4</sub>O·H<sub>2</sub>O

Calcd	C, 73.74;	H, 8.25;	N, 11.47
Found	C, 74.12;	H, 8.05;	N, 11.82

m.p.: 177-178°C (dec.) (crystallization solvent: ethyl acetate-diethyl ether)

#### Example 87

[0554] 4-(4-Fluorophenyl)-N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]-1-piperidinecarboxamide




[0555] Using the N-[2-((cis-2,6-dimethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 54 and Example 38 were conducted to obtain the titled compound as colorless amorphous powders.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (6H, d, J = 6.2 Hz), 1.22-2.02 (10H, m), 2.50-2.82 (3H, m), 2.94-3.14 (2H, m), 4.01 (2H, s), 4.10-4.36 (2H, m), 6.66 (1H, br), 7.01 (2H, dd, J = 8.4 and 8.8 Hz), 7.11-7.24 (2H, m), 7.50 (1H, dd, J = 2.2 and 8.8 Hz), 7.60 (1H, d, J = 8.4 Hz), 7.92-8.10 (3H, m).

FAB(pos): 475[MH]<sup>+</sup>

#### Example 88

[0556] 4-(4-Methoxy)-N-[2-((cis-2,6-dimethyl-1-piperidinyl)methyl)-6-quinolinyl]-1-piperidinecarboxamide




[0557] Using the N-[2-((cis-2,6-dimethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 54 and Example 38 were conducted to obtain the titled compound as colorless amorphous powders.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (6H, d, J = 6.2 Hz), 1.23-2.02 (10H, m), 2.50-2.83 (3H, m), 2.94-3.14 (2H, m), 3.80 (3H, s), 4.01 (2H, s), 4.19-4.36 (2H, m), 6.66 (1H, br), 6.87 (2H, d, J = 8.4 Hz), 7.15 (2H, d, J = 8.4 Hz), 7.50 (1H, dd, J = 2.6 and 9.2 Hz), 7.61 (1H, d, J = 8.4 Hz), 7.93-8.10 (3H, m).

FAB(pos): 487[MH]<sup>+</sup>

#### Example 89

[0558] 2',4-Difluoro-N-[2-((cis-2,6-dimethyl-1-piperidinyl)methyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0559] Using the N-[2-((cis-2,6-dimethyl-1-piperidinyl)methyl)-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as colorless amorphous powders.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.05 (6H, d, J = 6.2 Hz), 1.28-1.83 (6H, m), 2.51-2.82 (2H, m), 4.04 (2H, s), 6.88-7.08 (2H, m),

7.37-7.53 (1H, m), 7.60-7.74 (3H, m), 7.86 (1H, d, J = 8.4 Hz), 7.95-8.18 (5H, m), 8.46 (1H, d, J = 2.2 Hz).  
FAB(pos): 486[MH]<sup>+</sup>

# Example 90

[0560] 2',4-Dichloro-N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]-6-quinolinyl]-4-carboxamide




[0561] Using the N-[2-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-6-quinolinyl]acetamide obtained in Reference Example 11, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.04 (6H, d, J = 6.3 Hz), 1.27-1.80 (6H, m), 2.52-2.70 (2H, m), 4.03 (2H, s), 7.23-7.40 (2H, m), 7.46-7.60 (3H, m), 7.68 (1H, dd, J = 2.1 and 9.0 Hz), 7.86 (1H, d, J = 8.4 Hz), 7.93-8.05 (3H, m), 8.09 (1H, d, J = 8.4 Hz), 8.16 (1H, br), 8.46 (1H, d, J = 2.4 Hz).  
m.p.: 162-164°C (crystallization solvent: ethyl acetate)

# Example 91

[0562] 4'-Chloro-N-[2-[(diisopropylamino)methyl]-6-quinolinyl]-[1,1'-biphenyl]-4-carboxamide




[0563] Using the N-[2-[(diisopropylamino)methyl]-6-quinolinyl]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.06 (12H, d, J = 6.6 Hz), 2.96-3.19 (2H, m), 3.94 (2H, s), 7.45 (2H, d, J = 8.8 Hz), 7.56 (2H, d, J = 8.8 Hz), 7.63-7.73 (3H, m), 7.83 (1H, d, J = 8.8 Hz), 7.94-8.19 (5H, m), 8.46 (1H, d, J = 2.2 Hz).  
m.p.: 201-202°C (dec.) (crystallization solvent: ethyl acetate)

# Example 92

[0564] 4'-Chloro-N-[2-[(diethylamino)methyl]-6-quinolinyl]-[1,1'-biphenyl]-4-carboxamide




[0565] Using the 2-(diethylaminomethyl)-6-quinolinamine obtained in Reference Example 12, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.09 (6H, t, J = 7.2 Hz), 2.64 (4H, q, J = 7.2 Hz), 3.89 (2H, s), 7.45 (2H, d, J = 8.4 Hz), 7.57 (2H, d, J = 8.4 Hz), 7.62-7.78 (4H, m), 7.94-8.19 (5H, m), 8.47 (1H, d, J = 2.2 Hz).

m.p.: 196-198°C (decomposition) (crystallization solvent: ethyl acetate-diethyl ether)

#### Example 93

[0566] N-[2-[(2-Methyl-4,5-dihydro-1H-imidazol-1-yl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0567] Using the 2-(chloromethyl)-6-quinolinylamine dihydrochloride obtained in Reference Example 7, the same procedures as those of Reference Example 8 and Example 23 were conducted to obtain the titled compound as a powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.20 (3H, s), 3.36-3.96 (4H, m), 4.70 (2H, s), 7.34-7.56 (5H, m), 7.58-7.82 (3H, m), 7.91-8.26 (5H, m), 8.61 (1H, d, J = 2.2 Hz), 10.40 (1H, br).

FAB(pos): 421 [MH]<sup>+</sup>

m.p.: 212-220°C (decomposition) (crystallization solvent: ethyl acetate)

#### Example 94

[0568] N-[2-[(2-Phenyl-4,5-dihydro-1H-imidazol-1-yl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0569] Using the 2-(chloromethyl)-6-quinolinylamine dihydrochloride obtained in Reference Example 7, the same procedures as those of Reference Example 8 and Example 23 were conducted to obtain the titled compound as a powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 3.54 (2H, t, J = 9.8 Hz), 3.98 (2H, t, J = 9.8 Hz), 4.59 (2H, s), 7.30-7.80 (15H, m), 8.01 (2H, d, J = 8.0 Hz), 8.17 (1H, d, J = 8.4 Hz), 8.39 (1H, s), 8.55 (1H, d, J = 1.8 Hz).

FAB(pos): 483 [MH]<sup>+</sup>

m.p.: 212-216°C (decomposition) (crystallization solvent: ethyl acetate)

#### Example 95

[0570] 4'-Fluoro-N-[2-[(2-methyl-4,5-dihydro-1H-imidazol-1-yl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0571] Using the 2-(chloromethyl)-6-quinolinylamine dihydrochloride obtained in Reference Example 7, the same procedures as those of Reference Example 8 and Example 23 were conducted to obtain the titled compound as a pale brown powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.99(3H, s), 3.3(2H, br m), 3.5(2H, br m), 4.58(2H, s), 7.3-7.5(3H, br m), 7.8-8.1(9H, br m), 8.35(1H, br d), 8.56(1H, br s).

m.p.: 240-242°C (decomposition) (crystallization solvent: ethyl acetate)

#### Example 96

[0572] 4'-Fluoro-N-[2-((2-phenyl-4,5-dihydro-1H-imidazol-1-yl)methyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide




[0573] Using the 2-(chloromethyl)-6-quinolinylamine dihydrochloride obtained in Reference Example 7, the same procedures as those of Reference Example 8 and Example 23 were conducted to obtain the titled compound as a pale brown powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 3.45(2H, t, J=10.0Hz), 3.82(2H, t, J=10.0Hz), 4.50(2H, s), 7.31-7.66(8H, m), 7.80-8.15(9H, m), 8.35(1H, d, J=8.8Hz), 8.57(1H, d, J=2.2Hz).

m.p.: 209-210°C (crystallization solvent: ethyl acetate)

#### Example 97

[0574] N-[2-((2-Benzyl-4,5-dihydro-1H-imidazol-1-yl)methyl)-6-quinolinyl]-4'-fluoro-[1,1'-biphenyl]-4-carboxamide




[0575] Using the 2-(chloromethyl)-6-quinolinylamine dihydrochloride obtained in Reference Example 7, the same procedures as those of Reference Example 8 and Example 23 were conducted to obtain the titled compound as a pale brown powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 3.22(2H, t, J=9.8Hz), 3.60(2H, t, J=9.8Hz), 3.75(2H, s), 4.52(2H, s), 7.17-7.40(8H, m), 7.80-8.14(9H, m), 8.25(1H, d, J=8.4Hz), 8.54(1H, d, J=1.8Hz).

m.p.: 200-202°C (crystallization solvent: ethyl acetate)

## Example 98

[0576] Trans-2-(4-chlorophenyl)-N-[2-(1-pyrrolidinymethyl)-6-quinolinyl]-1,3-dioxane-5-carboxamide




[0577] Using the 2-(1-pyrrolidinymethyl)-6-quinolinylamine obtained in Reference Example 9 and 2-(4-chlorophenyl)-5-carboxy-1,3-dioxane obtained in Reference Example 14, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.82(4H, m), 2.62(4H, m), 3.10 (1H, m), 3.94(2H, s), 4.27 (2H, t like), 4.46(2H, dd like), 5.57(1H, s), 7.35-7.62(7H, m), 8.06(2H, t, J=12.6Hz), 8.29(1H, d, J=2.7Hz).

m.p.: 244-246°C (crystallization solvent: ethyl acetate)

## Example 99

[0578] 4-(4-Fluorophenyl)-N-[6-(1-pyrrolidinymethyl)-2-naphthyl]-1-piperidinecarboxamide



[0579] Using the 6-(1-pyrrolidinymethyl)naphthalene-2-amine obtained in Reference Example 18, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.57 (2H, m), 1.69 (4H, m), 1.80 (2H, m), 2.45 (4H, m), 2.77 (1H, m), 2.90 (2H, m), 3.68 (2H, s), 4.32 (2H, m), 7.13 (2H, m), 7.34 (2H, m), 7.38 (1H, m), 7.60 (1H, m), 7.66-7.76 (3H, m), 8.00 (1H, d, J = 2.1 Hz), 8.71 (1H, s).

FAB(pos): 432[M+H]<sup>+</sup>

m.p.: 209-211°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 100

[0580] 4-(4-Methoxyphenyl)-N-[6-(1-pyrrolidinymethyl)-2-naphthyl]-1-piperidinecarboxamide



[0581] Using the 6-(1-pyrrolidinymethyl)naphthalene-2-amine obtained in Reference Example 18, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.70-1.95 (8H, m), 2.55 (4H, m), 2.70 (1H, m), 3.02 (2H, m), 3.74 (2H, s), 3.80 (3H, s), 4.26 (2H, m), 6.56 (1H, s), 6.87 (2H, d, J = 8.4 Hz), 7.15 (2H, d, J = 8.8 Hz), 7.38 (2H, m), 7.69-7.76 (3H, m), 7.92 (1H, m).

FAB(pos): 444[M+H]<sup>+</sup>

m.p.:194-196°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 101

[0582] 4-(4-Chlorophenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide



[0583] Using the 6-(1-pyrrolidinylmethyl)naphthalene-2-amine obtained in Reference Example 18, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.55 (2H, m), 1.68 (4H, m), 1.78 (2H, m), 2.43 (4H, m), 2.76 (1H, m), 2.88 (2H, m), 3.65 (2H, s), 4.30 (2H, m), 7.33 (5H, m), 7.57 (1H, m), 7.64-7.74 (3H, m), 7.98 (1H, d, J = 2.5 Hz), 8.69 (1H, s).

FAB(pos): 432[M+H]<sup>+</sup>

m.p.:209-211°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 102

[0584] 4-Phenyl-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide



[0585] Using the 6-(1-pyrrolidinylmethyl)naphthalene-2-amine obtained in Reference Example 18, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.60 (2H, m), 1.70 (4H, m), 1.82 (2H, m), 2.46 (4H, m), 2.76 (1H, m), 2.91 (2H, m), 3.68 (2H, s), 4.32 (2H, m), 7.18-7.34 (5H, m), 7.39 (1H, m), 7.60 (1H, m), 7.69-7.76 (3H, m), 8.00 (1H, d, J = 1.8 Hz), 8.71 (1H, s).

FAB(pos): 414[M+H]<sup>+</sup>

m.p.:193-195°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 103

[0586] 4-Methylphenyl-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide



[0587] Using the 6-(1-pyrrolidinylmethyl)naphthalene-2-amine obtained in Reference Example 18, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.55 (2H, m), 1.70 (4H, m), 1.80 (2H, m), 2.26 (3H, s), 2.46 (4H, m), 2.71 (1H, m), 2.90 (2H, s),




m), 3.68 (2H, s), 4.31 (2H, m), 7.12 (4H, m), 7.39 (1H, m), 7.60 (1H, m), 7.66-7.76 (3H, m), 8.00 (1H, d, J = 1.8 Hz), 8.70 (1H, s).

FAB(pos): 428[M+H]<sup>+</sup>

m.p.: 210-212°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 104

[0588] 6-(4-Fluorophenyl)-N-[6-(1-piperidinylmethyl)-2-naphthyl]nicotinamide




[0589] Using the 6-(1-piperidinylmethyl)naphthalene-2-amine obtained in Reference Example 15, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 1.41 (2H, m), 1.51 (4H, m), 2.36 (4H, brs), 3.56 (2H, s), 7.38 (2H, dd, J=8.3 and 8.8Hz), 7.47 (1H, d, J=8.3Hz), 7.73 (1H, s), 7.83 (2H, m), 7.89 (1H, d, J=9.0Hz), 8.18 (1H, d, J=8.3Hz), 8.27 (2H, m), 8.44 (2H, m), 9.24 (1H, m), 10.64 (1H, brs).

m.p.: 218-219°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 105

[0590] 6-(4-Methoxyphenyl)-N-[6-(1-piperidinylmethyl)-2-naphthyl]nicotinamide




[0591] Using the 6-(1-piperidinylmethyl)naphthalene-2-amine obtained in Reference Example 15, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 1.40 (2H, m), 1.51 (4H, m), 2.37 (4H, brs), 3.57 (2H, s), 3.84 (3H, s), 7.09 (2H, d, J=8.3Hz), 7.46 (1H, d, J=8.1Hz), 7.73 (1H, s), 7.81 (2H, m), 7.89 (1H, d, J=9.0Hz), 8.10 (1H, d, J=8.1Hz), 8.17 (2H, d, J=8.3Hz), 8.39 (1H, d, J=8.3Hz), 8.43 (1H, s), 9.2 (1H, brs).

m.p.: 263-264°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 106

[0592] 6-(4-Chlorophenyl)-N-[6-(1-piperidinylmethyl)-2-naphthyl]nicotinamide




[0593] Using the 6-(1-piperidinylmethyl)naphthalene-2-amine obtained in Reference Example 15, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 1.41 (2H, m), 1.51 (4H, m), 2.36 (4H, brs), 3.56 (2H, s), 7.38 (2H, dd, J=8.3 and 8.8Hz), 7.47 (1H, d, J=8.3Hz), 7.73 (1H, s), 7.83 (2H, m), 7.89 (1H, d, J=9.0Hz), 8.18 (1H, d, J=8.3Hz), 8.27 (2H, m), 8.44 (2H, m) 9.24 (1H, m), 10.64 (1H, brs).

m.p.: 228-229°C (crystallization solvent: ethyl acetate-acetone)

#### Example 107

[0594] 4-(4-Fluorophenyl)-N-[6-(1-piperidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide




[0595] Using the 6-(1-piperidinylmethyl)naphthalene-2-amine obtained in Reference Example 15, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 1.40 (2H, m), 1.48-1.61 (6H, m), 1.80 (2H, d, J=12.0Hz), 2.34 (4H, brs), 2.77 (1H, m), 2.90 (2H, dd, J=12.0 and 12.2Hz), 3.52 (2H, s), 4.32 (2H, d, J=13.2Hz), 7.12 (2H, t, J=8.5Hz), 7.30 (1H, d, J=5.9Hz), 7.32 (1H, d, J=5.6Hz), 7.38 (1H, d, J=8.3Hz), 7.62 (2H, m), 7.68 (1H, d, J=8.3Hz), 7.74 (1H, d, J=8.8Hz), 8.00 (1H, s), 8.71 (1H, brs).

m.p.: 209-211°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 108

[0596] 4-(4-Chlorophenyl)-N-[6-(1-piperidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide




[0597] Using the 6-(1-piperidinylmethyl)naphthalene-2-amine obtained in Reference Example 15, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 1.39 (2H, m), 1.48-1.61 (6H, m), 1.81 (2H, m), 2.35 (4H, brs), 2.78 (1H, m), 2.90 (2H, dd, J=11.5 and 12.7Hz), 3.52 (2H, s), 4.32 (2H, d, J=13.7Hz), 7.30-7.39 (5H, m), 7.59-7.75 (4H, m), 8.01 (1H, s), 8.71 (1H, brs). m.p.: 231-232°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 109

[0598] 4-(4-Methylphenyl)-N-[6-(1-piperidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide




[0599] Using the 6-((1-piperidinyl)methyl)naphthalene-2-amine obtained in Reference Example 15, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): 1.40 (2H, m), 1.48-1.61 (6H, m), 1.80 (2H, d, J=12.4Hz), 2.26 (3H, s), 2.34 (4H, brs), 2.71 (1H, m), 2.89 (2H, dd, J=12.4 and 11.5Hz), 3.52 (2H, s), 4.31 (2H, d, J=12.9Hz), 7.10-7.16 (4H, m), 7.38 (1H, d, J=8.0Hz), 7.61 (2H, m), 7.68 (1H, d, J=8.5Hz), 7.74 (1H, d, J=8.8Hz), 8.00 (1H, s), 8.70 (1H, brs).

m.p.: 227-228°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 110

[0600] 4-(4-Chlorophenyl)-N-[6-[(*cis*-2,6-dimethyl-1-piperidinyl)methyl]-2-naphthyl]-1-piperidinecarboxamide




[0601] Using the 6-[(*cis*-2,6-dimethyl-1-piperidinyl)methyl]naphthalene-2-amine obtained in Reference Example 17, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): 0.98 (3H, s), 1.00 (3H, s), 1.28 (3H, m), 1.58 (3H, m), 1.81 (2H, m), 2.78 (1H, m), 2.90 (2H, m), 3.82 (2H, s), 4.31 (2H, m), 7.33 (4H, m), 7.43 (1H, m), 7.65 (1H, m), 7.72 (2H, m), 7.98 (1H, brs), 8.68 (1H, brs).

m.p.: 199-200°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 111

[0602] N-[6-[(*cis*-2,6-Dimethyl-1-piperidinyl)methyl]-2-naphthyl]piperidine-4-(4-methoxyphenyl)-1-carboxamide




[0603] Using the 6-[(*cis*-2,6-dimethyl-1-piperidinyl)methyl]naphthalene-2-amine obtained in Reference Example 17, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): 0.98 (3H, s), 1.00 (3H, s), 1.27 (3H, m), 1.57 (3H, m), 1.78 (2H, m), 2.69 (1H, m), 2.89 (2H, m), 3.72 (3H, s), 3.82 (2H, s), 4.31 (2H, m), 6.86 (2H, d, J=8.5Hz), 7.18 (2H, d, J=8.5Hz), 7.43 (1H, d, J=8.5Hz), 7.58 (1H, d, J=8.8Hz), 7.65 (1H, d, J=8.5Hz), 7.72 (2H, m), 7.98 (1H, brs), 8.67 (1H, brs).

m.p.: 170-171°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 112


[0604] N-6-[(cis-2,6-Dimethyl-1-piperidinyl)methyl]-2-naphthyl]piperidine-4-(4-fluorophenyl)-1-carboxamide



[0605] Using the 6-[(cis-2,6-dimethyl-1-piperidinyl)methyl]naphthalene-2-amine obtained in Reference Example 17, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): 0.98 (3H, s), 1.00 (3H, s), 1.28 (3H, m), 1.58 (3H, m), 1.81 (2H, m), 2.78 (1H, m), 2.90 (2H, m), 3.82 (2H, s), 4.31 (2H, m), 7.33 (4H, m), 7.43 (1H, m), 7.65 (1H, m), 7.72 (2H, m), 7.98 (1H, brs), 8.68 (1H, brs). m.p.:219-222°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 113


[0606] 6-(4-Chlorophenyl)-N-6-[(cis-2,6-dimethyl-1-piperidinyl)methyl]-2-naphthyl]nicotinamide



[0607] Using the 6-[(cis-2,6-dimethyl-1-piperidinyl)methyl]naphthalene-2-amine obtained in Reference Example 17, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): 0.99 (3H, s), 1.01 (3H, s), 1.29 (3H, m), 1.58 (3H, m), 3.65 (2H, s), 7.52 (1H, d, J=8.3Hz), 7.61 (2H, d, J=8.3Hz), 7.79 (2H, d, J=8.5Hz), 7.87 (2H, m), 8.20 (1H, m), 8.24 (2H, d, J=8.5Hz), 8.42 (1H, brs), 8.46 (1H, dd, J=2.2 and 8.3Hz), 9.25 (1H, d, J=2.0Hz), 10.63 (1H, brs). m.p.:229-230°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 114

[0608] N-6-[(cis-2,6-Dimethyl-1-piperidinyl)methyl]-2-naphthyl]-4'-methoxy[1,1'-biphenyl]-4-carboxamide




[0609] Using the 6-[(cis-2,6-dimethyl-1-piperidinyl)methyl]naphthalene-2-amine obtained in Reference Example 17, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder. <sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>): 0.99 (3H, s), 1.01 (3H, s), 1.29 (3H, m), 1.58 (3H, m), 3.82 (3H, s), 3.85 (2H, s), 7.07 (2H, d,

J=8.7Hz), 7.51 (1H, d, J=8.7Hz), 7.73 (2H, d, J=8.5Hz), 7.82 (6H, m), 8.06 (2H, d, J=8.3Hz), 8.42 (1H, brs), 10.41 (1H, brs).

m.p.: 199-200°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 115

[0610] 4'-Chloro-N-[6-[(cis-2,6-dimethyl-1-piperidiny)-methyl]-2-naphthyl][1,1'-biphenyl]-4-carboxamide




[0611] Using the 6-[(cis-2,6-dimethyl-1-piperidiny)methyl]naphthalene-2-amine obtained in Reference Example 17, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 0.99 (3H, s), 1.01 (3H, s), 1.29 (3H, m), 1.58 (3H, m), 3.85 (2H, s), 7.51 (1H, d, J=8.3Hz), 7.57 (2H, d, J=8.5Hz), 7.82 (8H, m), 8.12 (2H, d, J=8.1Hz), 8.43 (1H, brs), 10.47 (1H, brs).

m.p.: 220-222°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 116

[0612] 4'-Chloro-N-[2-[[2-(3-pyridinyl)-1-piperidiny]methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0613] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>): 1.63-1.75 (6H, m), 2.16 (1H, m), 2.87 (1H, m), 3.27 (2H, m), 3.73 (2H, m), 7.40 (1H, m), 7.59 (3H, m), 7.82-8.00 (7H, m), 8.13 (2H, d, J = 8.4 Hz), 8.30 (1H, d, J = 8.4 Hz), 8.47-8.54 (2H, m), 8.69 (1H, s).

FAB(pos): 533[M+H]<sup>+</sup>

m.p.: 124-126°C (dec.) (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 117

[0614] 4'-Chloro-N-[2-[(2S)-2-(methoxymethyl)pyrrolidinyl]methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide



[0615] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.68 (3H, m), 1.95 (1H, m), 2.33 (1H, m), 2.84 (2H, m), 3.25-3.44 (5H, m), 3.64-3.71 (1H, m), 4.28-4.35 (1H, m), 7.59 (3H, m), 7.82-8.08 (6H, m), 8.14 (2H, d, J = 8.4 Hz), 8.29 (1H, d, J = 8.4 Hz), 8.53 (1H, s), 10.62 (1H, s).

FAB(pos): 486[M+H]<sup>+</sup>

m.p.: 172-174°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 118

[0616] Tert-butyl (2S)-1-[(4'-Chloro[1,1'-biphenyl]-4-yl)carbonylamino]-2-quinolinylmethyl]-2-pyrrolidinecarboxylate



[0617] Using the 4'-chloro-N-[2-(chloromethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide obtained in Reference Example 8, the same procedures as those of Example 23 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.38 (9H, s), 1.80 (3H, m), 2.05 (1H, m), 2.92 (1H, m), 3.35 (2H, m), 3.82-4.17 (2H, m), 7.62 (3H, m), 7.82-8.08 (6H, m), 8.14 (2H, d, J = 8.8 Hz), 8.30 (1H, d, J = 8.4 Hz), 8.53 (1H, s), 10.63 (1H, s).

FAB(pos): 542[M+H]<sup>+</sup>

m.p.: 163-166°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 119


[0618] 4-(4-Methoxyphenyl)-N-[6-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthyl]-1-piperidinecarboxamide



[0619] Using the 6-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthaleneamine obtained in Reference Example 19, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.  
<sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 1.00 (12H, s), 1.53-1.81 (10H, m), 2.71 (1H, m), 2.88 (2H, m), 3.72 (3H, s), 3.90 (2H, s), 4.30 (2H, m), 6.86 (2H, d, *J* = 8.8 Hz), 7.18 (2H, d, *J* = 8.8 Hz), 7.52-7.79 (5H, m), 7.97 (1H, s), 8.66 (1H, s).  
 FAB(pos): 514[M+H]<sup>+</sup>  
 m.p.: 164-166°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 120


[0620] 4'-Methoxy-N-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthyl[[1,1'-biphenyl]-4-carboxamide



[0621] Using the 6-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthaleneamine obtained in Reference Example 19, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.  
<sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 1.01 (12H, s), 1.55 (6H, m), 3.82 (3H, s), 3.93 (2H, s), 7.06 (2H, d, *J* = 9.2 Hz), 7.57 (1H, d, *J* = 9.6 Hz), 7.71-7.87 (8H, m), 8.08 (2H, d, *J* = 8.4 Hz), 8.41 (1H, s), 10.40 (1H, s).  
 FAB(pos): 507[M+H]<sup>+</sup>  
 m.p.: 212-214°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 121

[0622] 4'-Fluoro-N-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthyl[[1,1'-biphenyl]-4-carboxamide



[0623] Using the 6-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthaleneamine obtained in Reference Example 19, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.01 (12H, s), 1.54 (6H, m), 3.93 (2H, s), 7.35 (2H, m), 7.74-7.87 (8H, m), 8.11 (2H, d, J = 8.4 Hz), 8.41 (1H, s), 10.43 (1H, s).

FAB(pos): 495[M+H]<sup>+</sup>

m.p.: 238-241°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 122

[0624] 4'-Methyl-N-[6-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthyl]-1,1'-biphenyl]-4-carboxamide



[0625] Using the 6-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthaleneamine obtained in Reference Example 19, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.01 (12H, s), 1.55 (6H, m), 2.37 (3H, s), 3.93 (2H, s), 7.32 (2H, d, J = 8.2 Hz), 7.57 (1H, d, J = 9.6 Hz), 7.66-7.87 (8H, m), 8.10 (2H, d, J = 8.6 Hz), 8.41 (1H, s), 10.41 (1H, s).

FAB(pos): 491[M+H]<sup>+</sup>

m.p.: 235-237°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 123

[0626] 4'-Chloro-N-[6-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthyl]-1,1'-biphenyl]-4-carboxamide



[0627] Using the 6-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-2-naphthaleneamine obtained in Reference Example 19, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.01 (12H, s), 1.54 (6H, m), 3.93 (2H, s), 7.57 (3H, m), 7.74-7.89 (8H, m), 8.12 (2H, d, J = 8.6 Hz), 8.41 (1H, s), 10.45 (1H, s).


FAB(pos): 511[M+H]<sup>+</sup>

m.p.: 244-246°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 124

[0628] 4'-Chloro-N-[6-[(diisopropylamino)methyl]-2-naphthyl]-1,1'-biphenyl]-4-carboxamide





[0629] Using the N-6-[(diisopropylamino)methyl]-2-naphthyl]-2-hydroxy-2-methylpropanamide obtained in Reference Example 20, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.03 (12H, d, J = 6.6 Hz), 3.01 (2H, m), 3.75 (2H, s), 7.52 (1H, d, J = 8.7 Hz), 7.57 (2H, d, J = 8.7 Hz), 7.76-7.89 (8H, m), 8.12 (2H, d, J = 8.4 Hz), 8.43 (1H, s), 10.47 (1H, s).

FAB(pos): 471[M+H]<sup>+</sup>

m.p.: 246-250°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 125

[0630] 6-(1-Pyrrolidiny(methyl)-2-naphthyl 4-(4-chlorophenyl)-1-piperidinecarboxylate



[0631] Using the 6-(1-pyrrolidiny(methyl)-2-naphthol hydrobromide obtained in Reference Example 22, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.68-1.83 (6H, m), 1.90-1.95 (2H, m), 2.55 (4H, m), 2.74 (1H, m), 2.97-3.12 (2H, m), 3.77 (2H, s), 4.49 (2H, m), 7.17 (2H, m), 7.26-7.33 (3H, m), 7.49 (1H, m), 7.55 (1H, d, J = 2.4 Hz), 7.73-7.76 (2H, m), 7.81 (1H, d, J = 9.3 Hz).

FAB(pos): 449[M+H]<sup>+</sup>

m.p.: 121-123°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 126

[0632] 6-(1-Pyrrolidiny(methyl)-2-naphthyl 4-(4-methoxyphenyl)-1-piperidinecarboxylate



[0633] Using the 6-(1-pyrrolidiny(methyl)-2-naphthol hydrobromide obtained in Reference Example 22, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.71-1.80 (8H, m), 2.50 (4H, m), 2.75 (1H, m), 2.99-3.15 (2H, m), 3.72 (5H, s-like), 4.14-4.32 (2H, m), 6.88 (2H, m), 7.20-7.33 (3H, m), 7.50 (1H, m), 7.65 (1H, s), 7.82-7.93 (3H, m).

FAB(pos): 445[M+H]<sup>+</sup>

m.p.: 127-129°C (crystallization solvent: ethyl acetate-diisopropyl ether)

Example 127

[0634] 6-(1-Pyrrolidinylmethyl)-2-naphthyl 4-(4-methylphenyl)-1-piperidinecarboxylate




[0635] Using the 6-(1-pyrrolidinylmethyl)-2-naphthol hydrobromide obtained in Reference Example 22, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.  
<sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 1.72-2.00 (8H, m), 2.28 (3H, s), 2.51 (4H, m), 2.68-3.15 (3H, m), 3.54 (2H, s), 4.30 (2H, m), 7.15-8.01 (10H, m).

FAB(pos): 429[M+H]<sup>+</sup>

m.p.: 238-240°C (crystallization solvent: ethyl acetate-diisopropyl ether)

Example 128

[0636] 6-(1-Pyrrolidinylmethyl)-2-naphthyl 4-(4-fluorophenyl)-1-piperidinecarboxylate



[0637] Using the 6-(1-pyrrolidinylmethyl)-2-naphthol hydrobromide obtained in Reference Example 22, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-*d*<sub>6</sub>) δ: 1.73-1.91 (8H, m), 2.51 (4H, m), 2.71-3.15 (3H, m), 3.76 (2H, s), 4.23 (2H, m), 7.15 (2H, m), 7.35 (3H, m), 7.52 (1H, m), 7.66 (1H, m), 7.90 (3H, m).

FAB(pos): 433[M+H]<sup>+</sup>

m.p.: 106-108°C (crystallization solvent: ethyl acetate-diisopropyl ether)

Example 129

[0638] 6-(1-Pyrrolidinylmethyl)-2-naphthyl 4-(4-phenyl-1-piperidinecarboxylate hydrochloride



[0639] Using the 6-(1-pyrrolidinylmethyl)-2-naphthol hydrobromide obtained in Reference Example 22, the same procedures as those of Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.69 (2H, m), 1.90 (4H, m), 2.00 (2H, m), 2.82 (1H, m), 3.12 (2H, m), 3.22 (4H, m), 4.21-4.49 (4H, m), 7.16-7.44 (5H, m), 7.75 (3H, m), 7.93-8.12 (3H, m), 10.69 (1H, br).

FAB(pos): 415[M+H]<sup>+</sup>

m.p.: 213-215°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 130

[0640] N-(2-Methyl-1,2,3,4-tetrahydrobenzo[b][1,6]naphthyridin-8-yl)[1-1'-biphenyl]-4-carboxamide




[0641] Using the 2-methyl-1,2,3,4-tetrahydrobenzo[b][1,6]naphthyridine-8-amine obtained in Reference Example 23, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.42 (3H, s), 2.80 (2H, m), 3.10 (2H, m), 3.71 (2H, s), 7.53 (3H, m), 7.78 (2H, d, J = 8.4 Hz), 7.83 (3H, m), 8.00 (2H, m), 8.12 (2H, d, J = 8.4 Hz), 8.47 (1H, d, J = 2.2 Hz), 10.56 (1H, s).

m.p.: 236-238°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 131

[0642] 4'-Fluoro-N-(2-methyl-1,2,3,4-tetrahydrobenzo[b][1,6]naphthyridin-8-yl)[1-1'-biphenyl]-4-carboxamide




[0643] Using the 2-methyl-1,2,3,4-tetrahydrobenzo[b][1,6]naphthyridine-8-amine obtained in Reference Example 23, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.42 (3H, s), 2.80 (2H, m), 3.10 (2H, m), 3.71 (2H, s), 7.36 (2H, m), 7.88 (5H, m), 7.99 (2H, m), 8.11 (2H, d, J = 8.4 Hz), 8.46 (1H, d, J = 2.2 Hz), 10.56 (1H, s).

m.p.: 230-232°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 132

[0644] 4'-Chloro-N-(2-methyl-1,2,3,4-tetrahydrobenzo[b][1,6]naphthyridin-8-yl)[1-1'-biphenyl]-4-carboxamide




[0645] Using the 2-methyl-1,2,3,4-tetrahydrobenzo[b][1,6]naphthyridine-8-amine obtained in Reference Example 23, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 2.42 (3H, s), 2.80 (2H, m), 3.07 (2H, m), 3.71 (2H, s), 7.58 (2H, d, J = 8.4 Hz), 7.80-7.99 (7H, m), 8.12 (2H, d, J = 8.4 Hz), 8.46 (1H, d, J = 2.0 Hz), 10.57 (1H, s).

m.p.:238-240°C (crystallization solvent: ethyl acetate-diisopropyl ether)

Example 133


[0646] N-[2-[(Diisopropylamino)methyl]-6-quinoliny] [1-1'-biphenyl]-4-carboxamide



[0647] Using the N-[2-[(diisopropylamino)methyl]-6-quinoliny]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.03 (12H, d, J = 6.6 Hz), 3.02 (2H, m), 3.86 (2H, s), 7.43-7.56 (3H, m), 7.69 (1H, d, J = 8.4 Hz), 7.67-8.05 (6H, m), 8.12 (2H, d, J = 8.4 Hz), 8.26 (1H, d, J = 8.4 Hz), 8.51 (1H, d, J = 2.2 Hz), 10.58 (1H, s).  
 FAB(pos) 438[M+H]<sup>+</sup>  
 m.p.:208-209°C (crystallization solvent: ethyl acetate-diisopropyl ether)

Example 134


[0648] N-[2-[(Diisopropylamino)methyl]-6-quinoliny]-4'-methoxy[1-1'-biphenyl]-4-carboxamide



[0649] Using the N-[2-[(diisopropylamino)methyl]-6-quinoliny]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ:1.02 (12H, d, J = 6.6 Hz), 3.02 (2H, m), 3.82 (3H, s), 3.85 (2H, s), 7.07 (2H, d, J = 9.2 Hz), 7.72 (3H, m), 7.82 (2H, d, J = 8.4 Hz), 7.96(2H, m), 8.08 (2H, d, J = 8.8 Hz), 8.25 (1H, d, J = 8.4 Hz), 8.50 (1H, s), 10.55 (1H, s).  
 FAB(pos) 468[M+H]<sup>+</sup>  
 m.p.:223-225°C (crystallization solvent: ethyl acetate-diisopropyl ether)

Example 135

[0650] N-[2-[(Diisopropylamino)methyl]-6-quinoliny]-4'-fluoro[1-1'-biphenyl]-4-carboxamide




[0651] Using the N-[2-[(diisopropylamino)methyl]-6-quinoliny]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.02 (12H, d, J = 6.6 Hz), 3.02 (2H, m), 3.86 (2H, s), 7.35 (2H, m), 7.69 (1H, d, J = 8.4 Hz), 7.80-7.88 (4H, m), 7.96 (2H, m), 8.11 (2H, d, J = 8.4 Hz), 8.25 (1H, d, J = 8.4 Hz), 8.51 (1H, d, J = 2.2 Hz), 10.58 (1H, s).  
FAB(pos) 456[M+H]<sup>+</sup>  
m.p.: 207-209°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 136


[0652] N-[2-[(Diisopropylamino)methyl]-6-quinolinyl]-4'-methyl [1-1'-biphenyl]-4-carboxamide



[0653] Using the N-[2-[(diisopropylamino)methyl]-6-quinolinyl]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.02 (12H, d, J = 6.6 Hz), 2.37 (3H, s), 3.02 (2H, m), 3.85 (2H, s), 7.32 (2H, d, J = 8.2 Hz), 7.69 (3H, m), 7.84 (2H, d, J = 8.4 Hz), 7.96 (2H, m), 8.10 (2H, d, J = 8.2 Hz), 8.25 (1H, d, J = 8.8 Hz), 8.51 (1H, d, J = 2.2 Hz), 10.56 (1H, s).  
FAB(pos) 452[M+H]<sup>+</sup>  
m.p.: 225-227°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 137


[0654] N-[2-[(Diisopropylamino)methyl]-6-quinolinyl]-6-(4-methylphenyl)nicotinamide



[0655] Using the N-[2-[(diisopropylamino)methyl]-6-quinolinyl]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.  
<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.02 (12H, d, J = 6.6 Hz), 2.39 (3H, s), 3.02 (2H, m), 3.86 (2H, s), 7.36 (2H, d, J = 8.4 Hz), 7.70 (1H, d, J = 8.4 Hz), 7.97 (2H, m), 8.13 (3H, m), 8.27 (1H, d, J = 8.4 Hz), 8.42 (1H, m), 8.50 (1H, d, J = 2.4 Hz), 9.22 (1H, d, J = 2.4 Hz), 10.70 (1H, s).  
FAB(pos) 453[M+H]<sup>+</sup>  
m.p.: 211-213°C (crystallization solvent: ethyl acetate-diisopropyl ether)

# Example 138

[0656] 6-(4-Chlorophenyl)-N-[2-[(diisopropylamino)methyl]-6-quinolinyl]nicotinamide



[0657] Using the N-[2-((diisopropylamino)methyl)-6-quinolinyl]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 6 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.02 (12H, d, J = 6.6 Hz), 3.02 (2H, m), 3.86 (2H, s), 7.61 (2H, m), 7.70 (1H, d, J = 8.1 Hz), 7.93 (2H, m), 8.23 (4H, m), 8.48 (2H, m), 9.25 (1H, m), 10.74 (1H, s).

FAB(pos) 473[M+H]<sup>+</sup>

m.p.: 201-203°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 139

[0658] N-[2-((Diisopropylamino)methyl)-6-quinolinyl]-4-(4-methylphenyl)-1-piperidinecarboxamide



[0659] Using the N-[2-((diisopropylamino)methyl)-6-quinolinyl]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 54 and Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.02 (12H, d, J = 6.9 Hz), 1.60-1.77 (4H, m), 2.26 (3H, s), 2.85-3.03 (5H, m), 3.82 (2H, s), 4.43 (2H, m), 7.12 (4H, m), 7.62 (1H, d, J = 8.4 Hz), 7.79 (2H, s-like), 8.05 (1H, s), 8.12 (1H, d, J = 8.8 Hz), 8.81 (1H, s).

FAB(pos) 459[M+H]<sup>+</sup>

m.p.: 184-186°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 140

[0660] 4-(4-Chlorophenyl)-N-[2-((diisopropylamino)methyl)-6-quinolinyl]-1-piperidinecarboxamide



[0661] Using the N-[2-((diisopropylamino)methyl)-6-quinolinyl]acetamide obtained in Reference Example 10, the same procedures as those of 2) of Example 54 and Example 38 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.01 (12H, d, J = 6.3 Hz), 1.59 (2H, m), 1.81 (2H, s), 2.78-3.00 (5H, m), 3.82 (2H, s), 4.32 (2H, m), 7.32 (4H, m), 7.62 (1H, d, J = 8.1 Hz), 7.79 (2H, s-like), 8.05 (1H, s), 8.12 (1H, d, J = 9.0 Hz), 8.81 (1H, s).

FAB(pos) 479[M+H]<sup>+</sup>

m.p.: 201-203°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 141

[0662] 5-[(4-Bromobenzyl)oxy]-2-(1-pyrrolidinylmethyl)-1H-indole




[0663] To a solution of [5-[(4-bromobenzyl)oxy]-1H-indol-2-yl]methanol (100 mg, 0.301 mmol) obtained in Reference Example 24 and triethylamine (0.050 ml, 0.361 mmol) in tetrahydrofuran (3 ml) was added methanesulfonyl chloride (0.0256 ml, 0.331 mmol) at 0°C, and the mixture was stirred for 15 minutes. To the reaction solution was added pyrrolidine (0.3 ml), the mixture was stirred for 1 hour, and 1N hydrochloric acid was added, followed by washing with diethyl ether. Potassium carbonate was added to the aqueous layer to adjust to basic, the mixture was extracted with ethyl acetate, washed with an aqueous saturated sodium chloride solution, and dried over anhydrous sodium sulfate. The solvent was concentrated under reduced pressure, the resulting residue was purified by alumina column chromatography (developing solvent: ethyl acetate), and converted into powders by isopropyl ether to obtain the titled compound (1.8 mg)

<sup>1</sup>H-NMR (CDCl<sub>3</sub>) δ: 1.83 (4H, m), 2.62 (4H, m), 3.80 (2H, s), 5.04 (2H, s), 6.27 (1H, s), 6.85 (1H, dd, J = 2.4, 8.4 Hz), 7.05 (1H, d, J = 2.4 Hz), 7.19 (1H, d, J = 8.4 Hz), 7.33 (2H, d, J = 8.6 Hz), 7.49 (2H, d, J = 8.6 Hz), 8.95 (1H, s).

## Example 142

[0664] 6-Phenyl-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide




[0665] Using the 6-(1-pyrrolidinylmethyl)naphehalene-2-amine obtained in Reference Example 18, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.73 (4H, m), 2.50 (4H, m), 3.75 (2H, s), 7.47-7.59 (4H, m), 7.77-7.92 (4H, m), 8.20 (3H, m), 8.44 (2H, m), 9.25 (1H, m), 10.63 (1H, s).

m.p.: 212-214°C (crystallization solvent: ethyl acetate-diisopropyl ether)

## Example 143

[0666] 6-(4-Methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide



[0667] Using the 6-(1-pyrrolidinylmethyl)naphehalene-2-amine obtained in Reference Example 18, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.


<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.72 (4H, m), 2.50 (4H, m), 3.73 (2H, s), 3.85 (3H, s), 7.09 (2H, d, J = 9.0 Hz), 7.47 (1H, dd, J

= 1.5, 8.4 Hz), 7.76-7.91 (4H, m), 8.10 (1H, d, J = 8.4 Hz), 8.18 (2H, d, J = 9.0 Hz), 8.38-8.44 (2H, m), 9.20 (1H, m), 10.59 (1H, s).

m.p.: 261-263°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 144

[0668] 6-(4-Fluorophenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide




[0669] Using the 6-(1-pyrrolidinylmethyl)naphehalene-2-amine obtained in Reference Example 18, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.71 (4H, m), 2.47 (4H, m), 3.72 (2H, s), 7.41 (2H, m), 7.48 (1H, d, J = 9.6 Hz), 7.76-7.91 (4H, m), 8.18 (1H, d, J = 8.4 Hz), 8.28 (2H, m), 8.45 (2H, m), 9.24 (1H, d, J = 2.4 Hz), 10.63 (1H, s).

m.p.: 238-240°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 145

[0670] 6-(4-Methylphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide




[0671] Using the 6-(1-pyrrolidinylmethyl)naphehalene-2-amine obtained in Reference Example 18, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.71 (4H, m), 2.39 (3H, s), 2.47 (4H, m), 3.71 (2H, s), 7.35 (2H, d, J = 7.8 Hz), 7.45 (1H, dd, J = 1.8, 8.4 Hz), 7.74-7.89 (4H, m), 8.12 (3H, m), 8.44 (2H, m), 9.23 (1H, m), 10.62 (1H, br).

m.p.: 260-262°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Example 146

[0672] 6-(4-Chlorophenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide



[0673] Using the 6-(1-pyrrolidinylmethyl)naphehalene-2-amine obtained in Reference Example 18, the same procedures as those of 3) of Example 54 were conducted to obtain the titled compound as a colorless powder.

<sup>1</sup>H-NMR (DMSO-d<sub>6</sub>) δ: 1.71 (4H, m), 2.47 (4H, m), 3.71 (2H, s), 7.47 (1H, dd, J = 1.8, 8.4 Hz), 7.61 (2H, d, J = 8.4 Hz), 7.76-7.88 (4H, m), 8.23 (3H, m), 8.45 (2H, m), 9.24 (1H, d, J = 1.5 Hz), 10.68 (1H, br).



m.p.:270-274°C (crystallization solvent: ethyl acetate-diisopropyl ether)

#### Preparation Example 1

[0674]

(1) Compound obtained in Example 1	50 mg
(2) Lactose	34 mg
(3) Corn starch	10.6 mg
(4) Corn starch (paste)	5 mg
(5) Magnesium stearate	0.4 mg
(6) Carboxymethylcellulose calcium	20 mg
Total	120 mg

[0675] In accordance with a conventional manner, the above (1) to (6) are admixed and tableted using a tableting machine to give tablets.

#### Reference Example 1-1

Amplification of rat SLC-1 receptor cDNA by PCR method using rat-brain-originated cDNA

[0676] Reverse transcription reaction was carried out using random primer, with rat-brain-originated poly (A)<sup>+</sup>RNA (Clone Tech Co.) as a template. The reagent from the TaKaRa RNA PCR ver. 2 kit was used for the reverse transcription reaction. Next, using this reverse transcription product as a template, amplification was carried out by a PCR method using synthetic DNA primers with SEQ ID NOS: 1 and 2. Synthetic DNA primers were constructed to amplify genes in the domain where genes were translated into the receptor protein. At that time, individual restriction enzyme recognition sequences were also added to the 5' side and 3' side of the gene so as to add a nucleotide sequence recognizing the restriction enzyme Sal I to the 5' side of the gene, and to add a nucleotide sequence recognizing the restriction enzyme Spe I to the 3' side of the gene. The reaction mixture was composed of 5 µl of cDNA template, 0.4 µM of respective synthetic DNA primer, 0.25 mM of dNTPs, 0.5 µl of Pfu (StrataGene Co.) DNA polymerase, and buffers attached to enzymes, with setting total reaction quantity at 50 µl.

[0677] A thermal cycler (Parkin Elmer Co.) was used to produce cycles for amplification. After heating at 94°C for 60 seconds, the cycle consisting of 94°C for 60 seconds, 60°C for 30 seconds, and 72°C for 150 seconds, was repeated 35 times, and finally reaction was conducted at 72°C for 10 minutes. After 0.8% agarose gel electrophoresis, the amplified products were confirmed by ethidium bromide dyeing.

#### Reference Example 1-2

Subcloning of PCR products into plasmid vector, and confirmation of an amplified cDNA sequence by decoding of a nucleotide sequence in an inserted cDNA portion

[0678] The reaction product after PCR conducted in Reference Example 1-1 was separated using 0.8% low-melting point agarose gel. After the band section was cut out using a razor, DNA was recovered by conducting fragmentation, phenol extraction, phenol-chloroform extraction and ethanol precipitation. The recovered DNA was subcloned on plasmid vector PCR-Script Amp SK(+) in accordance with prescription of the PCR-Script™ Amp SK(+) cloning kit (Stratagene Co.). After this was introduced into *Escherichia coli* XL-1 Blue (Stratagene Co.) by transformation, the clones with fragments of inserted cDNA were selected in LB agar culture medium containing ampicillin and X-gal. Only clones showing white color were separated using a sterilized toothpick, and transformant *E. coli* XL-1 Blue/rat SLC-1 was obtained.

[0679] Each clone was cultured overnight in LB culture medium containing ampicillin, and plasmid DNA was prepared using QIA prep8 mini prep (Qiagen). A portion of the prepared DNA was digested with Sal I and Spe I, and the size of the inserted receptor cDNA fragment was confirmed. Reactions to determine nucleotide sequences were carried out using a DyeDeoxy Terminator Cycle Sequence Kit (Parkin Elmer Co.), and decoded using a fluorescent light automatic sequencer. The sequences of the 3 clones obtained were analyzed, and it was confirmed that all of them match the reported gene sequence (SEQ ID NO: 4) in which the Sal I recognition sequence was added to the 5' side and the Spe I recognition sequence was added to the 3' side of the cDNA sequence (Lakaye, B., et al., Biochim. Biophys. Acta,

Vol. 1401, pp. 216-220 (1998), accession No. AF08650) coding rat SLC-1 protein (SEQ ID NO: 3).

#### Reference Example 1-3

##### Preparation of CHO cells for rat SLC-1 expression

[0680] The full-length amino acid sequence of rat brain originated SLC-1, which was confirmed in Reference Example 1-2, was coded, and plasmid was prepared using a plasmid Midi Kit (Qiagen) from the *E. coli* transformed by the plasmid, to which the gene with Sal I recognition sequence added to the 5' side and Spe I recognition sequence added to the 3' side, had been introduced. Then, the insert section was cut out by digesting with Sal I and Spe I. The insert DNA was cut out with a razor from the agarose gel after electrophoresis.

[0681] Next, fragmentation, phenol extraction, phenol-chloroform extraction, and ethanol precipitation, were conducted and the DNA was recovered. This insert DNA was added to vector plasmid pAKKO-111H (the same vector plasmid as pAKKO1.11H described in Hinuma, S., et al., Biochim. Biophys. Acta, Vol. 1219, pp. 251-259 (1994)) for animal cell expression which was digested with Sal I and Spe I, and ligation was conducted using T4 ligase (TaKaRa Shuzo), to construct pAKKO-SLC-1 plasmid for protein expression.

[0682] After *E. coli* DH5 (TOYOBO) transformed by pAKKO-SLC-1 was cultured, pAKKO-SLC-1 plasmid DNA was prepared using a Plasmid Midi Kit (Qiagen). This was introduced into CHO dhfr cells in accordance with the attached protocol, using a CellPfect Transfection Kit (Amersham Pharmacia Biotech Co.). A coprecipitating suspension of 10 µg of DNA and calcium phosphate was prepared, and this suspension was added to 10 cm Petri dishes in which  $5 \times 10^5$  or  $1 \times 10^6$  of CHO dhfr cells had been seeded 24 hours previously. After these cells were cultured for 1 day in MEMα culture medium containing 10% fetal bovine serum, subculture was conducted, and cultivation was conducted in selective culture medium, MEMα culture medium containing no nucleic acid but containing 10% dialyzed fetal bovine serum. 56 clones of colonies of the transformed CHO cells expressing SLC-1, proliferated in the selective culture medium, were selected.

#### Reference Example 1-4

##### Selection of CHO/SLC-1 cell strain expressing a large quantity of full-length rat SLC-1 receptor protein mRNA

[0683] The quantity of expressed full-length rat SLC-1 receptor protein mRNA of 56 clones of the CHO/SLC-1 strains established in Reference Example 1-3, was measured using a Cytostar T Plate (Amersham Pharmacia Biotech Co.) as shown below according to the attached protocol. Each well of the Cytostar T Plate was seeded with each clone of the CHO/SLC-1 strain by  $2.5 \times 10^4$ , and cultured for 24 hours, then the cells were fixed using 10% formalin. After 0.25% Triton X-100 was added to each well to increase cell permeability,  $^{35}$ S-labeled riboprobes with SEQ ID NO: 5 were added and hybridized. 20 mg/ml of RNaseA was added to each well to digest free riboprobes. After the plate was thoroughly washed, the radioactivity of the hybridized riboprobes was determined using a Topcounter. Strains with high radioactivity showed large amounts of mRNA expression. In particular, mainly used was Clone number 44 among 3 clones which showed large amounts of mRNA expression.

#### Reference Example 1-5

##### Isolation of plasmid containing human SLC-1 cDNA

[0684] After nicks were inserted into the DNA of Human fetal brain originated cDNA library (SUPERScript™ cDNA Library; GIBCOBRL Co.) according to the manual of the Genetrappor cDNA positive selection system (GIBCOBRL Co.), using phage F1 endonuclease, single stranded human fetal brain originated cDNA library was prepared by digesting the above-mentioned library with *Escherichia coli* exonuclease III.

[0685] Biotin-14-dCTP was added to the 3' end of synthetic oligonucleotide (equivalent to 1434-1451 of accession No. U71092) of SEQ ID NO: 6 which was prepared according to the report by Kolakowski Jr., et al. (Kolakowski Jr., et al. (1996) FEBS Lett. Vol. 398, pp. 253-258) using Terminal Deoxynucleotidyl Transferase, and biotinlated oligonucleotide was prepared. The above manual was followed regarding composition of a reaction mixture and reaction time.

[0686] After 4 µg of single stranded human fetal brain originated cDNA library was kept at 95°C for 1 minute, the library was rapidly cooled on ice. 20 ng of Biotinated oligonucleotide was added, which was hybridized using the attached hybridization buffer at 37°C for 1 hour. Streptavidin beads were added to the mixture, then single stranded human fetal brain originated cDNA hybridized by biotinlated oligonucleotide, was isolated using a MAGNA-SEP Magnetic Particle Separator (GIBCOBRL Co.). The complementary strand was synthesized according to the manual, using as primer 50 ng of synthetic oligonucleotide (equivalent to 1011 - 1028 of accession No. U71092) of SEQ ID NO: 7,

prepared based on the report by Kolakowski Jr., et al (Kolakowski Jr., et al. (1996) FEBS Lett. Vol. 398, pp. 253-258), to give the double stranded plasmid.

#### Reference Example 1-6

Determination of nucleotide sequence of plasmid containing isolated human SLC-1 cDNA

[0687] After the plasmid obtained in Reference Example 1-5 was introduced into ELECTROMAX™ DH10B™ Cells by the electroporation method, clones with cDNA inserted fragments were selected in LB agar culture medium containing ampicillin and X-gal. Using a sterilized toothpick, only the clones showing white color were separated to give transformant *E. coli* DH10B/hSLC-1. Individual clones were cultured overnight in LB culture medium containing ampicillin, and the plasmid DNA was refined using QIA prep8 mini prep (Qiagen). The reactions to determine nucleotide sequence were conducted using a DyeDeoxy Terminator Cycle Sequence Kit (Parkin Elmer Co.), and the nucleotide sequence was decoded using a fluorescent light automatic sequencer.

[0688] As the results, obtained was the sequence shown in SEQ ID NO: 8. The amino acid sequence (SEQ ID NO: 9) coded by the nucleotide sequence obtained here, differs from the human SLC-1 amino acid sequence predicted as the sequence analogized from rat SLC-1 based on human chromosome DNA sequence (accession number: Z86090) containing human SLC-1 sequence, in the report by Lakaye, et al. (Lakaye, B., et al. (1998) Biochem. Biophys. Acta. Vol. 1401, pp. 216-220). This shows the presence of ATG, the initiation codon, on mRNA, in the 69 and 64 amino acids upstream from the estimated sequence. *Escherichia coli* DH10B/phSLC1L8, the transformant produced by the plasmid containing DNA coding this sequence was deposited at IFO and NIBH.

#### Reference Example 1-7

Amplification of human SLC-1cDN by PCR method using human fetal brain originated cDNA

[0689] Amplification by the PCR method was conducted using as the template plasmid containing human SLC-1 DNA sequence cloned by the gene trap method, and using synthetic DNA primers of SEQ ID NO: 10 and SEQ ID NO: 11, and synthetic DNA primers of SEQ ID NO: 12 and SEQ ID NO: 13, respectively. The former amplified DNA and the latter amplified DNA were named as "human SLC-1(S)" and "human SLC-1(L)", respectively. The synthetic DNA primer was constructed so that the genes in the domain translated to the receptor protein were amplified. At that time, a recognition sequence for each restriction enzyme was added to the 5' side and 3' side, so that the nucleotide sequence recognized by restriction enzyme Sal I would be added to the 5' side of the gene, and the nucleotide sequence recognized by restriction enzyme Spe I would be added to the 3' side. The composition of the reaction mixture for human SLC-1(S) amplification was: 5 µl of plasmid template containing human SLC-1 DNA sequence, 0.4 µM of respective synthetic DNA primers, 0.2 mM of dNTPs and 0.5 µl of Pfu DNA polymerase and buffers attached to the enzyme, with setting total quantity for reaction at 50 µl. A thermal cycler (Parkin Elmer Co.) was used for the cycles for amplification. After heating at 94°C for 60 seconds, the cycle consisting of 94°C for 60 seconds, 57°C for 60 seconds, and 72°C for 150 seconds, was repeated 25 times, and finally the temperature of the reactant was maintained at 72°C for 10 minutes. The composition of the reaction mixture for human SLC-1(L) amplification was 5 µl of plasmid template containing human SLC-1 DNA sequence, 0.4 µM of respective synthetic DNA primers, 0.2 mM of dNTPs, 0.5 µl of Pfu DNA polymerase and buffers attached to the enzymes, with setting total quantity for reaction at 50 µl. A thermal cycler (Parkin Elmer Co.) was used for the cycles for amplification. After heating at 94°C for 60 seconds, the cycle consisting of 94°C for 60 seconds, 60°C for 60 seconds, and 72°C for 3 minutes, was repeated 25 times, and finally the temperature of the reactant was maintained at 72°C for 10 minutes. After 0.8% agarose gel electrophoresis, confirmation of amplified products was conducted by ethidium bromide dyeing.

#### Reference Example 1-8

Subcloning of PCR product into plasmid vector and confirmation of amplified cDNA sequence by decoding of nucleotide sequence of inserted cDNA section

[0690] The reaction product after PCR in Reference Example 1-7 was separated using 0.8% low-melting point agarose gel, and the band section was cut out using a razor. After that, fragmentation, phenol extraction, phenol-chloroform extraction, and ethanol precipitation were conducted, and the DNA was recovered. The recovered DNA was subcloned into pCR-Script Amp SK(+) plasmid vector, as prescribed by the PCR-Script™ Amp SK(+) cloning kit (Stratagene Co.). After this was introduced into *Escherichia coli* DH5 a competent cells (TOYOBO) and transformed, the clones with cDNA inserted fragments were selected in LB agar culture medium containing ampicillin and X-gal. Using a sterilized

toothpick, only clones showing white color were separated to give *E. coli* DH5 $\alpha$ /hSLC-1(S), which is a transformant of human SLC-1 (S), and *E. coli* DH5 $\alpha$ /hSLC-1(L), which is a transformant of human SLC-1 (L). Each clone was cultured overnight in LB culture medium containing ampicillin, and plasmid DNA was prepared using QIA prep8 mini prep (Qiagen). Some of the prepared DNA was digested with Sal I and Spe I restriction enzymes, and the size of the receptor cDNA fragments inserted was confirmed. The reactions to determine nucleotide sequence were conducted using a DyeDeoxy Terminator Cycle Sequence Kit (Parkin Elmer Co.) and the nucleotide sequence was decoded using a fluorescent light automatic sequencer. The sequence of the obtained clones respectively matched the DNA sequence (SEQ ID NO: 14) which should be amplified by synthetic DNA primers of SEQ ID NO: 10 and SEQ ID NO: 11 using human SLC-1 gene as a template, and the DNA sequence (SEQ ID NO: 15) which should be amplified by synthetic DNA primers of SEQ ID NO: 12 and SEQ ID NO: 13 using human SLC-1 gene as a template.

#### Reference Example 1-9

Preparation of CHO cells for expression of human SLC-1(S), and CHO cells for expression of human SLC-1(L)

[0691] Plasmid was prepared from the *E. coli* clones transformed by the plasmid wherein inserted were human SLC-1(S) and human SLC-1(L) whose sequences were confirmed in Reference Example 1-8, using a Plasmid Midi Kit (Qiagen), and the insert section was cut out using Sal I and Spe I restriction enzymes. After electrophoresis was conducted, the insert DNA was cut out from agarose gel using a razor. Next, fragmentation, phenol extraction, phenol-chloroform extraction, and ethanol precipitation were conducted, and the insert DNA was recovered.

[0692] This insert DNA was added to pAKKO-111H vector plasmid for animal cell expression, digested with Sal I and Spe I (the same vector plasmid as the pAKKO1.11H described in Hinuma, S., et al., *Biochim. Biophys. Acta*, Vol. 1219, pp. 251-259 (1994)), and ligation was conducted by adding T4 ligase (TaKaRa Shuzo), to construct pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L) plasmids for protein expression.

[0693] After *E. coli* DH5 $\alpha$  (TOYOBO) transformed by pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L) was cultured, pAKKO-hSLC-1(S) and pAKKO-hSLC-1(L) plasmid DNAs were prepared using a Plasmid Midi Kit (Qiagen). These were introduced into CHO dhfr cells in accordance with the attached protocol, using a CellPlect Transfection Kit (Amersham Pharmacia Biotech Co.). A coprecipitative suspension of 10  $\mu$ g of DNA with calcium phosphate was made, which was added to 10 cm Petri dishes seeded 24 hours in advance with  $5 \times 10^5$  or  $1 \times 10^6$  CHO dhfr cells. After the above was cultured for 1 day in MEM $\alpha$  culture medium containing 10% fetal bovine serum, subculture was conducted, and then cultivation was conducted in MEM $\alpha$  culture medium containing no nucleic acid but containing 10% dialyzed fetal bovine serum, which is a selective culture medium. 56 clones of colonies of transformed cells which are human SLC-1(S) gene introduced CHO cells, and 61 clones of colonies of transformed cells which are human SLC-1(L) gene introduced CHO cells, both of which proliferated in the selective culture medium, were selected.

#### Reference Example 1-10

Selection of cell colonies into which genes with large quantities of human SLC-1(S) and human SLC-1 (L) mRNA expression have been introduced

[0694] The quantities of expressed mRNA of 56 clones of CHO/hSLC-1(S) colonies and 61 clones of CHO/hSLC-1(L) colonies, both of which were established in Reference Example 1-9, were measured in accordance with the attached protocol using a Cytostar T Plate (Amersham Pharmacia Biotech Co.) as shown below.

[0695] After each well of the Cytostar T Plate was seeded with each clone of CHO/hSLC-1(S) colonies and CHO/hSLC-1(L) colonies by  $2.5 \times 10^4$ , and cultured for 24 hours, the cells were fixed using 10% formalin.

[0696] After 0.25% Triton X-100 was added to each well to increase cell permeability,  $^{32}$ S-labeled riboprobe of SEQ ID NO: 16 was added and hybridization was conducted.

[0697] 20 mg/ml of RNaseA was added to each well to digest free riboprobe. After the plate was washed well, the radioactivity of the hybridized riboprobe was determined. Colonies showing high radioactivity expressed large quantities of mRNA. Of the 7 clones which expressed large quantities of mRNA, mainly used was Clone number 57.

#### Experimental Example 1

Determination of antagonist activity using GTP $\gamma$ S binding assay of test compound

[0698] Membrane fraction was prepared by the following method, using the human SLC-1 expressing CHO cell clone 57 obtained in Reference Example 1-10, and the rat SLC-1 expressing CHO cell clone 44 obtained in Reference Example 1-4.

[0699] The human and rat SLC-1 expressing CHO cells ( $1 \times 10^6$ ) were scraped in buffer saline phosphate (pH 7.4) to which 5 mM EDTA (ethylenediaminetetraacetic acid) had been added, and centrifuged. 10 ml of homogenized buffer (10 mM  $\text{NaHCO}_3$ , 5 mM EDTA, pH 7.5) was added to the cell pellets, and they were homogenized using a Polytron homogenizer. The supernatant obtained by centrifugation at  $400 \times g$  for 15 minutes was further centrifuged at  $100,000 \times g$  for 1 hour, to obtain the membrane fraction precipitate. This precipitate was suspended in 2 ml of assay buffer [50 mM Tris-HCl (pH 7.5), 1 mM EDTA, 0.1% BSA (bovine serum albumin), 10 mM  $\text{MgCl}_2$ , 100 mM NaCl, 1  $\mu\text{M}$  GDP (guanosine 5'-diphosphate), 0.25 mM PMSF (phenylmethylsulfonyl fluoride), 1 mg/ml pepstatin, 20 mg/ml leupeptin, 10 mg/ml phosphoramidon], which was centrifuged at  $100,000 \times g$  for 1 hour. The membrane fraction recovered as precipitate was suspended again in 2 ml of assay buffer, and after the suspension was divided, individual portions were preserved at  $-80^\circ\text{C}$  and thawed before every use.

[0700] Determination of antagonist activity of the test compound was conducted as shown below. After 171  $\mu\text{l}$  of SLC-1 expressing CHO cell membrane fractions diluted with assay buffer was poured into each well of a 96-well polypropylene plate, 2  $\mu\text{l}$  of  $3 \times 10^{-10}\text{M}$  MCH diluted with DMSO solution, 2  $\mu\text{l}$  of test compound solution diluted to various concentrations, and 25  $\mu\text{l}$  of [ $^3\text{S}$ ]-Guanosine 5'-( $\gamma$ -thio) triphosphate (produced by Daiichi Kagaku Yakuhin) were added respectively. (Final concentration of cell membrane: 20  $\mu\text{g}/\text{ml}$ , final concentration of [ $^3\text{S}$ ]-Guanosine 5'-( $\gamma$ -thio) triphosphate: 0.33 nM).

[0701] After this reaction mixture was allowed to react for 1 hour under stirring, it was filtered under vacuum using a glass filter (GF-C), then the filter was washed 3 times with 300  $\mu\text{l}$  of washing solution (50 mM Tris-HCl buffer solution pH 7.5). 50 ml of liquid scintillator was added to the glass filter, and residual radioactivity was determined using a liquid scintillation counter.

[0702] The  $\text{IC}_{50}$  value of the compound was calculated from the binding inhibition rate (%), based on the definition that the binding inhibition rate (%) = (radioactivity when compound and MCH were added - radioactivity when DMSO solution was added) / (radioactivity when MCH was added - radioactivity when DMSO solution was added)  $\times 100$ .

[0703] The results were shown below.

Compound Number	Inhibition Activity ( $\text{IC}_{50}$ value: nM)
Example 1	5

#### Industrial Applicability

[0704] Compounds (I), (I') and salts thereof possess excellent MCH receptor antagonistic activities, and are useful as an agent for preventing or treating obesity, etc.

## SEQUENCE LISTING

5

&lt;110&gt; Takeda Chemical Industries, Ltd.

10

&lt;120&gt; Melanin-Concentrating Hormone Antagonist

15

&lt;130&gt; 2723W00P

20

&lt;150&gt; JP 2000-134295

&lt;151&gt; 2000-04-28

25

&lt;150&gt; JP 2000-384897

&lt;151&gt; 2000-12-13

30

&lt;160&gt; 16

35

&lt;210&gt; 1

&lt;211&gt; 32

40

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

45

&lt;220&gt;

&lt;223&gt;

50

&lt;400&gt; 1

55

gtcgacatgg atctgcaaac ctcgltgctg tg 32

5           <210> 2  
           <211> 32  
           <212> DNA  
 10           <213> Artificial Sequence  
  
 15           <220>  
           <223>  
  
 20           <400> 2  
 25           actagttcag gtgcctttgc ttctgtctct ct   32  
  
 30           <210> 3  
           <211> 353  
           <212> PRT  
 35           <213> Rat  
  
 40           <400> 3  
           Met Asp Leu Gln Thr Ser Leu Leu Ser Thr Gly Pro Asn Ala Ser Asn  
           1                   5                   10                   15  
 45           Ile Ser Asp Gly Gln Asp Asn Leu Thr Leu Pro Gly Ser Pro Pro Arg  
                           20                   25                   30  
 50           Thr Gly Ser Val Ser Tyr Ile Asn Ile Ile Met Pro Ser Val Phe Gly  
                   35                   40                   45  
 55           Thr Ile Cys Leu Leu Gly Ile Val Gly Asn Ser Thr Val Ile Phe Ala

	50	55	60	
5	Val Val Lys Lys Ser Lys Leu His Trp Cys Ser Asn Val Pro Asp Ile			
	65	70	75	80
10	Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu Phe Leu Leu Gly Met			
	85	90	95	
15	Pro Phe Met Ile His Gln Leu Met Gly Asn Gly Val Trp His Phe Gly			
	100	105	110	
20	Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp Ala Asn Ser Gln Phe			
	115	120	125	
25	Thr Ser Thr Tyr Ile Leu Thr Ala Met Thr Ile Asp Arg Tyr Leu Ala			
	130	135	140	
30	Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg Lys Pro Ser Met Ala			
	145	150	155	160
35	Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser Phe Ile Ser Ile Thr			
	165	170	175	
40	Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe Pro Gly Gly Ala Val			
	180	185	190	
45	Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr Asp Leu Tyr Trp Phe			
	195	200	205	
50	Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu Pro Phe Val Val Ile			
	210	215	220	
55	Thr Ala Ala Tyr Val Lys Ile Leu Gln Arg Met Thr Ser Ser Val Ala			
	225	230	235	240
	Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr Lys Arg Val Thr Arg			
	245	250	255	



Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val Cys Trp Ala Pro Tyr

5                   260                   265                   270

Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser Arg Pro Thr Leu Thr

10                   275                   280                   285

Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu Gly Tyr Ala Asn Ser

15                   290                   295                   300

Cys Leu Asn Pro Phe Val Tyr Ile Val Leu Cys Glu Thr Phe Arg Lys

305                   310                   315                   320

20 Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln Gly Gln Leu Arg Thr

325                   330                   335

25 Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg Thr Glu Ser Lys Gly

340                   345                   350

30 Thr

<210> 4

35 <211> 1074

<212> DNA

40 <213> Rat

45 <400> 1

gtcgacatgg atctgcaaac ctcgttgctg tccactggcc ccaatgccag caacatctcc 60

gatggccagg ataatctcac attgccgggg tcacctctc gcacaggag tgtctccac 120

atcaacatca llatgccttc cglgtttggt accatctgtc tctgggcat cgtgggaaac 180

tccacgglica tctttgctgt ggtgaagaag tccaagctac actgglgcag caagctcccc 240

55 gacatcttca lcatcaacct cctgttggtg gatctgclet tctgtclggg catgccttc 300

atgatccacc agclcatggg gaacggcgctc tggcactttg gggaaacat gtgcacccic 360  
 5 aicacagcca tggacgccna cagtcagttc actagcacct acatcctgac tgccatgacc 420  
 attgaccgct acttggccac cgtecacccc atctcctcca ccaagttccg gaagccctcc 480  
 10 atggccaccc lggatgatc cctcctgtgg gcgctctcct tcatcagtat caccctgtg 540  
 lggctctacg ccaggctcat tcccttccca ggggggtgclg lgggctgigg catccgcctg 600  
 15 ccaaacccgg acactgacct ctactggttc actctgtacc agtltttcct ggcttttgcc 660  
 ctlccgtttg tggcattac cgccgalac gtgaaaatac tacagcgcat gacgtcttcg 720  
 gtggccccag cctcccaacg cagcatccgg ctctggacaa agagggtgac ccgcacggcc 780  
 20 attgccatct gtciggtctt ctttgtgtgc tgggcaccct actaigtgct gcagctgacc 840  
 cagclgtcca tcagccgcc gacctcacg ttgtctact tglacaacg gccatcagc 900  
 25 ttgggctatg ctaacagctg cctgaacccc ttgtgtaca tagtgtctg tgagacctt 960  
 cgaaaacgct tgggtgtgtc agtgaacct gcagcccagg ggcagctccg caggtcagc 1020  
 30 aacgtcaga cagctgatga ggagaggaca gaaagcaaa gacctgaac tagt 1074

<210> 5

<211> 262

<212> RNA

<213> Rat

<400> 5

gcgaauuggg uaccgggccc cccucgagg ucgacggguu cgauaagcuu gauaucgaau 60  
 uccugcagcc cgggggaucc gcccacuagu ucaggugccu uugcuuucug uccucuccuc 120  
 50 aucagcuguc ugagcgugc ugaccgugcg gagcugcccc uggcgucgag gcuuacuga 180  
 caacaccaag cguuuucgaa aggucucaca gacacuaug uacacaagg gguuacggca 240  
 55 gcuguuagca uagcccaagc ug 262

5 <210> 6  
<211> 18  
10 <212> DNA  
<213> Artificial Sequence  
15 <220>  
<223>  
20  
<400> 6  
25 caacagctgc ctcaaccc 18  
30 <210> 7  
<211> 18  
<212> DNA  
35 <213> Artificial Sequence  
40 <220>  
<223>  
45 <400> 7  
cctggatgac tgcctcct 18  
50  
<210> 8  
55 <211> 1275

&lt;212&gt; DNA

&lt;213&gt; Human

&lt;400&gt; 8

taggtgatgt cagtgggagc catgaagaag ggagtgggga gggcagtlgg gcttggaggc 60  
 ggcagcggct gccaggclac ggaggaagac ccccttccca actgcggggc ttgcgtccg 120  
 ggacaagggt gcaggcgctg gaggcctgcc cagcctgcgt ggggtggagg gagctcagct 180  
 cggltgtggg agcaggcgac cggcactggc tggatggacc tggaaagctc gctgctgccc 240  
 actggtccca acgccagcaa cacctctgat ggccccgata acctcacttc ggcaggatca 300  
 cctcctcgca cggggagcat ctccacatc aacatcatca tgccttcggt gtccggcacc 360  
 atctgcctcc tgggcatcat cgggaacacc acggtcatct tcgcggctgt gaagaagtcc 420  
 aagctgcact ggtgcaacaa cgtccccgac atcttcatca tcaacctctc ggtagtagat 480  
 ctctctcttc tcttgggcat gcccttcagt atccaccagc tcatgggcaa tgggggtgtg 540  
 cactltgggg agaccatgtg caccctcatc acggccatgg atgccaatag tcagttcacc 600  
 agcacctaca tctgaccgc catggccatt gaccgctacc tggccactgt ccacccacac 660  
 tcttcacaga agttccggaa gccctctgtg gccaccctgg tgatctgctt cctgtgggcc 720  
 ctctccttca tcagcatcac cctgtgtggt ctgtatgcca gactcatccc ctccccagga 780  
 ggtgcagttg gctgcggcat acgcccggcc aaccagaca ctgacctcta ctggttcacc 840  
 ctgtaccagt ttttctggc ctttgccttg cttttgtggt tcatacagc cgcatacgtg 900  
 aggalcctgc agcgcagac gtccctcagtg gcccccgcct cccagcgcag catccggctg 960  
 cggacaaga gggtagcccg cacagccatc gccatcgtc ttggtctctt tgtgtgctgt 1020  
 gcacctact aigtgtaca gctgaccagc ttgtccatca gccgccgac cctcaccttt 1080  
 gtctacttat acaatgcggc catcagcttg ggctatgcca acagctgctt caacccttt 1140  
 gtgtacatcg tgctctgtga gacgttccgc aaacgcttgg tctgtcgtt gaagcctgca 1200  
 gccaggggc agcttcgcgc tgcagcaac gtcagacgg ctgacgagga gaggacagaa 1260

agcaaaggca cctga

1275

5

&lt;210&gt; 9

10

&lt;211&gt; 422

&lt;212&gt; PRT

15

&lt;213&gt; Human

&lt;400&gt; 9

20

MeT Ser Val Gly Ala MeT Lys Lys Gly Val Gly Arg Ala Val Gly Leu

1 5 10 15

25

Gly Gly Gly Ser Gly Cys Gln Ala Thr Glu Glu Asp Pro Leu Pro Asn

20 25 30

30

Cys Gly Ala Cys Ala Pro Gly Gln Gly Gly Arg Arg Trp Arg Leu Pro

35 40 45

35

Gln Pro Ala Trp Val Glu Gly Ser Ser Ala Arg Leu Trp Glu Gln Ala

50 55 60

Thr Gly Thr Gly Trp MeT Asp Leu Glu Ala Ser Leu Leu Pro Thr Gly

65 70 75 80

40

Pro Asn Ala Ser Asn Thr Ser Asp Gly Pro Asp Asn Leu Thr Ser Ala

85 90 95

45

Gly Ser Pro Pro Arg Thr Gly Ser Ile Ser Tyr Ile Asn Ile Ile MeT

100 105 110

50

Pro Ser Val Phe Gly Thr Ile Cys Leu Leu Gly Ile Ile Gly Asn Ser

115 120 125

55

Thr Val Ile Phe Ala Val Val Lys Lys Ser Lys Leu His Trp Cys Asn

130                      135                      140  
 5      Asn Val Pro Asp Ile Phe Ile Ile Asn Leu Ser Val Val Asp Leu Leu  
 145                      150                      155                      160  
 10      Phe Leu Leu Gly Met Pro Phe Met Ile His Gln Leu Met Gly Asn Gly  
 165                      170                      175  
 15      Val Trp His Phe Gly Glu Thr Met Cys Thr Leu Ile Thr Ala Met Asp  
 180                      185                      190  
 20      Ala Asn Ser Gln Phe Thr Ser Thr Tyr Ile Leu Thr Ala Met Ala Ile  
 195                      200                      205  
 25      Asp Arg Tyr Leu Ala Thr Val His Pro Ile Ser Ser Thr Lys Phe Arg  
 210                      215                      220  
 30      Lys Pro Ser Val Ala Thr Leu Val Ile Cys Leu Leu Trp Ala Leu Ser  
 225                      230                      235                      240  
 35      Phe Ile Ser Ile Thr Pro Val Trp Leu Tyr Ala Arg Leu Ile Pro Phe  
 245                      250                      255  
 40      Pro Gly Gly Ala Val Gly Cys Gly Ile Arg Leu Pro Asn Pro Asp Thr  
 260                      265                      270  
 45      Asp Leu Tyr Trp Phe Thr Leu Tyr Gln Phe Phe Leu Ala Phe Ala Leu  
 275                      280                      285  
 50      Pro Phe Val Val Ile Thr Ala Ala Tyr Val Arg Ile Leu Gln Arg Met  
 290                      295                      300  
 55      Thr Ser Ser Val Ala Pro Ala Ser Gln Arg Ser Ile Arg Leu Arg Thr  
 305                      310                      315                      320  
 325                      330                      335  
 55      Lys Arg Val Thr Arg Thr Ala Ile Ala Ile Cys Leu Val Phe Phe Val

Cys Trp Ala Pro Tyr Tyr Val Leu Gln Leu Thr Gln Leu Ser Ile Ser

5 340 345 350

Arg Pro Thr Leu Thr Phe Val Tyr Leu Tyr Asn Ala Ala Ile Ser Leu

10 355 360 365

Gly Tyr Ala Asn Ser Cys Leu Asn Pro Phe Val Ile Val Leu Cys

15 370 375 380

Glu Thr Phe Arg Lys Arg Leu Val Leu Ser Val Lys Pro Ala Ala Gln

385 390 395 400

20 Gly Gln Leu Arg Ala Val Ser Asn Ala Gln Thr Ala Asp Glu Glu Arg

405 410 415

25 Thr Glu Ser Lys Gly Thr

420

30 <210> 10

<211> 31

35 <212> DNA

<213> Artificial Sequence

40 <220>

<223>

45 <400> 10

50 gtcgacatgg acctggaagc ctcgtgctg c 31

55 <210> 11

<211> 31

5

<212> DNA

<213> Artificial Sequence

10

<220>

<223>

15

<400> 11

20

actagttcag gtgcctttgc tttctgtcct c 31

25

<210> 12

<211> 33

<212> DNA

30

<213> Artificial Sequence

35

<220>

<223>

40

<400> 12

45

agtcgacatg tcagtgggag ccatgaagaa ggg 33

50

<210> 13

<211> 33

<212> DNA

55

<213> Artificial Sequence



<220>

<223>

<400> 13

aactagttca ggtgccttlg ctttctgtcc tct 33

<210> 14

<211> 1074

<212> DNA

<213> Human

<400> 14

gtcgacatgg acctggaagc ctgctgctg cccactggtc ccaacgccag caacacctct 60  
gatggcccg ataacctcac ttggcaggga tcacctcttc gcacggggag catctctctac 120  
atcaacatca tcatgccttc ggtgttcggc accatctgcc tcttgggcat catcgggaac 180  
tccacggta ctttcgggt cgtgaagaag tccaagctgc actggtgcaa caacgtcccc 240  
gacatcttca tcatcaacct ctcggtagta gatctctct tctccttggg catgccttcc 300  
atgatccacc agctcatggg caatggggtg tggcacttgg gggagaccat gtgcacctc 360  
atcacggcca tggatgccaa tagtcagttc accagcacct acatcctgac cgccatggcc 420  
attgaccgt acctggccac lgtccacccc atctcttcca cgaagtccg gaagccctct 480  
gtggccaccc tggtagctg cctcctgtgg gccctctct tcatcagcat caccctgtg 540  
tggctgtat ccagactcat ccccttccca ggaggtgcag tgggctgcgg catacgccg 600  
cccaaccag acactgacct ctactgttc accctgtacc agtlttctct ggcctttgcc 660  
ctgccllttg tggatcatcac agccgcatac gtgaggatcc tgcagcgcal gacgtctca 720  
gtggcccccg cctccagcg cagcatcgg ctggggacaa agagggtgac ccgcacagcc 780

atcgccalet gictggtett cttlgtgtgc tgggcaccct actatgtgct acagctgacc 840  
 5 cagtgttcca lcagccgcc gacctcacc ttgtctact tatacaatgc ggccatcagc 900  
 ttgggctaig ccaacagctg cctcaacccc ttgtgtaca tcgtgctctg tgagacgttc 960  
 10 cgcaaacgtg tggctctgtc ggtgaagccl gcagcccagg ggcagcttgc cgtgtlccagc 1020  
 aacgctcaga cggctgacga ggagaggaca gaaagcaaag gcacctgaac tagt 1074

15 <210> 15

<211> 1283

20 <212> DNA

<213> Human

25 <400> 15

agtcgacalg tcagtgggag ccatgaagaa gggagtgagg agggcagttg ggcttggagg 60  
 30 cggcagcggc tgcaggcta cggaggaaga cccccitccc aactgcgggg cttgcctccc 120  
 gggacaaggt ggcaggcgtt ggaggtgcc gcagccttgc tgggtggagg ggagctcagc 180  
 35 tcggttgttg gagcaggcga cgggcacttg ctggatggac ctggaagcct cgctgctgcc 240  
 cactgggcc aacgccagca acacctctga tggcccgat aacctcactt cggcaggatc 300  
 40 acctctctgc acggggagca tctcttacct caacatcate atgccttcgg tgttcggcac 360  
 catctgcctc ctgggcatca tcgggaactc cagggtcatc ttgcggctgc tgaagaagtc 420  
 caagctgcac tggtgcaaca aglccccga calcttcac atcaacctct cggtagtaga 480  
 45 tctctctctt ctcttgggca tgccttcat gatccaccag clcatgggca alggggtgtg 540  
 gcacttlggg gagaccatgt gcacctcat caggccatg gatccaata gtcagttcac 600  
 50 cagcacctac atcttgaccg ccatggccat tgaccgtac ctggccactg tccaccccat 660  
 ctcttcacag aagltccgga agcctctgt gggccacctg gtgatctgcc tctgtgggc 720  
 55 cctctcttic atcagcatca cccctgtgtg gclgtatgcc agactcalcc ccttcccagg 780

aggtgcagtg ggctgcggca tacgcctgcc caaccagac actgacctct aciggttcac 840  
 cctgiaccag tttttcctgg cctttgccct gccttttttg gtcalcacag ccgcatacgt 900  
 gaggatcctg cagcgcatga cgtccicagt ggccccgcc tcccagcgca gcacacggct 960  
 gcgcacaag aggggtgacc gcacagccat cgccatctgt ctggtcttct ttgtgtgctg 1020  
 ggcacctac latgtgclac agctgacca gltgccatc agccgcgcga cctcacctt 1080  
 tglctactia tacaatgcgg ccacagcctt gggctatgcc aacagctgcc tcaaccctt 1140  
 lgtgtacatc gtgctctglg agacgttccg caaacgcttg gtcctgicgg tgaagcctgc 1200  
 agcccagggg cagcttcgcg ctgtcagcaa cgcicagacg gctgacgagg agaggacaga 1260  
 aagcaaaggc acctgaacta gtt 1283

<210> 16

<211> 420

<212> RNA


<213> Human

<400> 16

caaaagcugg agcuccaccg cgguggcggc cgcucuagcc cacuaguuca ggugccuuug 60  
 cuuuucugucc ucuccucguc agccgucuga gcguugcuga cagcgcgaag cugccccugg 120  
 gcugcaggcu ucaccgacag gaccaagcgu uugcgggaac ucucacagag caccgaugac 180  
 acaaaggggu ugaggcagcu guuggcauag cccaagcuga ugccgcgaau guauaaguag 240  
 acaaagrga gggucggcgc gcugauggac aacuggguca gcugaugcac auaguagggu 300  
 gccccagaca caaagaagac cagacagaug gcgauggcug ugccgggucac ccucuuuguc 360  
 cgcagccgga ugucgcgcur ggaggcgggg gccacugagg acgucaugcg cugcaggau 420

# Claims

1. A melanin-concentrating hormone antagonist which comprises a compound of the formula:



10 wherein Ar<sup>1</sup> is a cyclic group which may be substituted;

X and Y are the same or different and are a spacer having a main chain of 1 to 6 atoms;

Ar is a condensed polycyclic aromatic ring which may be substituted;

15 R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or


R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom, Y and Ar, may form a nitrogen-containing condensed ring which may be substituted; or a salt thereof.

- 20 2. The antagonist according to claim 1, wherein R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, form a nitrogen-containing heterocyclic ring which may be substituted.

- 25 3. The antagonist according to claim 1 which is an agent for preventing or treating diseases caused by melanin-concentrating hormone.

4. The antagonist according to claim 1 which is an agent for preventing or treating obesity.

- 30 5. A compound of the formula:



40 wherein Ar<sup>1</sup> is a cyclic group which may be substituted;

X<sup>1</sup> is CONR<sup>8</sup>, NR<sup>8</sup>CO (wherein R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl or optionally halogenated C<sub>1-6</sub> alkylsulfonyl), OCO or COO;

Y is a spacer having a main chain of 1 to 6 atoms;

45 Ar is a condensed polycyclic aromatic ring which may be substituted;

R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or

R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom, Y and Ar, may form a nitrogen-containing condensed ring which may be substituted;

50 provided that, when X<sup>1</sup> is CONR (wherein R is hydrogen atom or C<sub>1-6</sub> alkyl), Ar is not indole or benzoxazole which may have one or two halogen, hydroxy, C<sub>1-6</sub> alkyl or C<sub>1-6</sub> alkoxy;

when X<sup>1</sup> is CONH, Ar is not 4-methyl-2-quinolone which may have a substituent selected from the group consisting of alkyl, alkoxy and halogen, or is not 2-benzoylamino-quinazoline; and, when X<sup>1</sup> is COO, Ar<sup>1</sup> is not an aromatic group which may be substituted; or a salt thereof.

- 55 6. The compound according to claim 5, wherein X<sup>1</sup> is CONR<sup>8</sup> or NR<sup>8</sup>CO (wherein R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl or optionally halogenated C<sub>1-6</sub> alkylsulfonyl);


and R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted.

7. The compound according to claim 5, wherein the cyclic group represented by Ar<sup>1</sup> is an aromatic group.
8. The compound according to claim 7, wherein the aromatic group is formed by removing an optional one hydrogen atom from an aromatic ring assembly formed by 2 or 3 members selected from C<sub>6-14</sub> monocyclic or condensed polycyclic aromatic hydrocarbon and 5- to 10-membered aromatic heterocyclic ring.
9. The compound according to claim 5, wherein Ar<sup>1</sup> is phenyl, biphenyl or phenyl-pyridyl, each of which may be substituted with 1 to 3 substituents selected from halogen, optionally halogenated C<sub>1-6</sub> alkyl and optionally halogenated C<sub>1-6</sub> alkoxy.
10. The compound according to claim 5, wherein Ar<sup>1</sup> is piperidinyl which may be substituted with C<sub>6-14</sub> aryl which may be substituted.
11. The compound according to claim 5, wherein X<sup>1</sup> is CONH or COO.
12. The compound according to claim 5, wherein the condensed polycyclic aromatic ring represented by Ar is a condensed polycyclic aromatic hydrocarbon having 9 to 14 carbon atoms.
13. The compound according to claim 5, wherein the condensed polycyclic aromatic ring represented by Ar is a 10-membered condensed polycyclic aromatic heterocyclic ring.
14. The compound according to claim 5, wherein the condensed polycyclic aromatic ring represented by Ar is quinoline or naphthalene.
15. The compound according to claim 5, wherein X<sup>1</sup> is CONR<sup>8</sup> or NR<sup>8</sup>CO (wherein R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl or optionally halogenated C<sub>1-6</sub> alkyl(sulfonyl), and Ar is quinoline or naphthalene).
16. The compound according to claim 5, wherein the "spacer having a main chain of 1 to 6 atoms" represented by Y is a bivalent group consisting of 1 to 3 species selected from -O-, -S-, -CO-, -SO-, -SO<sub>2</sub>-, -NR<sup>8</sup>- (R<sup>8</sup> is hydrogen atom, optionally halogenated C<sub>1-6</sub> alkyl, optionally halogenated C<sub>1-6</sub> alkyl-carbonyl, optionally halogenated C<sub>1-6</sub> alkyl(sulfonyl), and an optionally halogenated bivalent C<sub>1-6</sub> non-cyclic hydrocarbon group).
17. The compound according to claim 5, wherein Y is C<sub>1-3</sub> alkylene.
18. The compound according to claim 5, wherein R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, form a nitrogen-containing heterocyclic ring which may be substituted.
19. The compound according to claim 18, wherein the nitrogen-containing heterocyclic ring is morpholine, piperidine, piperazine, pyrrolidine, 1,3-thiazolidine, 1H-imidazole, 4,5-dihydro-1H-imidazole, 2,3-dihydroindole, 1,2,3,4-tetrahydroquinoline or 1,2,3,4-tetrahydroisoquinoline.
20. A pharmaceutical composition comprising the compound according to claim 5, or a salt thereof.
21. A prodrug of the compound according to claim 5.
22. The compound according to claim 5 which is:
  - 4'-chloro-N-6-[(N,N-dimethylamino)methyl]-2-naphthyl[[1,1'-biphenyl]-4-carboxamide;
  - 4'-chloro-N-6-[(1-pyrrolidinyl)methyl]-2-naphthyl[[1,1'-biphenyl]-4-carboxamide;
  - 4'-fluoro-N-2-(1-pyrrolidinyl)methyl-6-quinolinyl[[1,1'-biphenyl]-4-carboxamide;
  - 4'-fluoro-N-2-(1-piperidinyl)methyl-6-quinolinyl[[1,1'-biphenyl]-4-carboxamide;
  - 4'-chloro-N-2-[(2-methyl-4,5-dihydro-1H-imidazol-1-yl)methyl]-6-quinolinyl[[1,1'-biphenyl]-4-carboxamide;

4'-chloro-N-[2-[(2,2,6,6-tetramethyl-1-piperidinyl)methyl]-6-quinolinyl][1,1'-biphenyl]-4-carboxamide;  
 4-(4-chlorophenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]-1-piperidinecarboxamide;  
 N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl][1,1'-biphenyl]-4-carboxamide;  
 6-(4-methylphenyl)-N-[2-(1-pyrrolidinylmethyl)-6-quinolinyl]nicotinamide;  
 4-(4-methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]-1-piperidinecarboxamide;  
 6-(4-methoxyphenyl)-N-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide;  
 6-(4-methylphenyl)-14-[6-(1-pyrrolidinylmethyl)-2-naphthyl]nicotinamide;

or a salt thereof.


23. A process for producing a compound of the formula:



wherein each symbol is as defined in claim 5, or a salt thereof, which comprises reacting a compound of the formula:



wherein  $\text{Ar}^1$  is as defined in claim 5, or a salt thereof with a compound of the formula:




wherein L is a leaving group and the other symbols are as defined above, or a salt thereof.

24. The antagonist according to claim 1 which is an anorectic agent.

25. A pharmaceutical which comprises the melanin-concentrating hormone antagonist according to claim 1 in combination with at least one species selected from the group consisting of an agent for treating diabetes, an agent for treating hypertension and an agent for treating arteriosclerosis.

26. A method for preventing or treating diseases caused by a melanin-concentrating hormone in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of the formula:



10 wherein Ar<sup>1</sup> is a cyclic group which may be substituted;


X and Y are the same or different and are a spacer having a main chain of 1 to 6 atoms;

Ar is a condensed polycyclic aromatic ring which may be substituted;

15 R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or

R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom, Y and Ar, may form a nitrogen-containing heterocyclic ring which may be substituted; or a salt thereof.

- 20 27. A method for preventing or treating obesity in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of the formula:



30 wherein Ar<sup>1</sup> is a cyclic group which may be substituted;


X and Y are the same or different and are a spacer having a main chain of 1 to 6 atoms;

Ar is a condensed polycyclic aromatic ring which may be substituted;

35 R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or

R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom, Y and Ar, may form a nitrogen-containing condensed ring which may be substituted; or a salt thereof.

- 40 28. Use of a compound of the formula:



50 wherein Ar<sup>1</sup> is a cyclic group which may be substituted;

X and Y are the same or different and are a spacer having a main chain of 1 to 6 atoms;


Ar is a condensed polycyclic aromatic ring which may be substituted;

55 R<sup>1</sup> and R<sup>2</sup> are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or

R<sup>1</sup> and R<sup>2</sup>, together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or R<sup>2</sup>, together with the adjacent nitrogen atom, Y and Ar, may form a nitrogen-

containing condensed ring which may be substituted; or a salt thereof for the manufacture of a pharmaceutical preparation for preventing or treating diseases caused by a melanin-concentrating hormone.

29. Use of a compound of the formula:



wherein  $\text{Ar}^1$  is a cyclic group which may be substituted;

X and Y are the same or different and are a spacer having a main chain of 1 to 6 atoms;

Ar is a condensed polycyclic aromatic ring which may be substituted;

$\text{R}^1$  and  $\text{R}^2$  are the same or different and are hydrogen atom or a hydrocarbon group which may be substituted; or

$\text{R}^1$  and  $\text{R}^2$ , together with the adjacent nitrogen atom, may form a nitrogen-containing heterocyclic ring which may be substituted; or  $\text{R}^2$ , together with the adjacent nitrogen atom and Y, may form a nitrogen-containing heterocyclic ring which may be substituted; or  $\text{R}^2$ , together with the adjacent nitrogen atom, Y and Ar, may form a nitrogen-containing condensed ring which may be substituted; or a salt thereof for the manufacture of a pharmaceutical preparation for preventing or treating obesity.



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/03614

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> Int. Cl. <sup>7</sup> A61K31/167, 31/40, 31/4453, 31/4709, 31/4545, 31/454, 31/4375, 31/47, 31/4725, C07C233/80, 237/48, C07D211/16, 211/14, 211/18, 295/135, 405/12, 401/06, 417/06, 401/12, 211/22, 471/04, A61P43/00, 3/04 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) Int. Cl. <sup>7</sup> A61K31/167, 31/40, 31/4453, 31/4709, 31/4545, 31/454, 31/4375, 31/47, 31/4725, C07C233/80, 237/48, C07D211/16, 211/14, 211/18, 295/135, 405/12, 401/06, 417/06, 401/12, 211/22, 471/04, A61P43/00, 3/04 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CAPLUS (STN), MEDLINE (STN), EMBASE (STN)		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/52875 A1 (Takeda Chemical Industries, Ltd.), 21 October, 1999 (21.10.99), especially, page 83 & EP 1070054 A1 & JP 2000-226373 A	3, 4, 5-23, 24, 25, 28, 29
X	WO 95/32967 A1 (SmithKline Beecham PLC), 07 December, 1995 (07.12.95), & EP 763034 A1 & JP 10-500960 A	5-23
A	WO 98/38156 A1 (Takeda Chemical Industries, Ltd.), 03 September, 1998 (03.09.98), & EP 971878 A1 & JP 11-80098 A	5-23
P, A	WO 01/21577 A2 (Takeda Chemical Industries, Ltd.), 29 March, 2001 (29.03.01), (Family: none)	1-4, 5-23, 24, 25, 28, 29
P, A	WO 00/31021 A1 (Takeda Chemical Industries, Ltd.), 02 June, 2000 (02.06.00) & JP 2000-212076 A	1-4, 5-23, 24, 25, 28, 29
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reasons (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to underpin the principles or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search 30 July, 2001 (30.07.01)		Date of mailing of the international search report 07 August, 2001 (07.08.01)
Name and mailing address of the ISA/ Japanese Patent Office		Authorized officer
Facsimile No.		Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/03614

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
F, A	NO 00/40725 A1 (Takeda Chemical Industries, Ltd.), 13 July, 2000 (13.07.00), & JP 2001-141728 A	1-4, 5-23, 24, 25, 28, 29

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/03614

**Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 26,27  
because they relate to subject matter not required to be searched by this Authority, namely:  
Claims 26 and 27 pertain to methods for treatment of the human body by therapy, and thus relate to subject matters which this International Searching Authority is not required, under the provisions of Article 17(2)(a) (i) of the PCT and Rule 39.1(iv) of the Regulations under the PCT, to search.
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

(See extra sheet.)

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☒ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.  
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1992)

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP01/03614

Continuation of Box No. II of continuation of first sheet (1)

The inventions of claims 1 to 4, 24, 25, 28, and 29 relate to melanin concentrating hormone (MCH) antagonists containing as the active ingredient compounds represented by the general formula (I) in claims 1, 28, and 29 and preventive and therapeutic drugs for MCH-related diseases such as obesity.

The inventions of claims 5 to 23 relate to compounds represented by the general formula (I') in claim 5, processes for the preparation of the compounds, and drugs for nonspecified uses containing the compounds.

In terms of chemical substance, the compounds of the general formula (I') are included among those represented by the general formula (I) in claim 1. Further, the results of prior art search reveal that two inventive concepts, i.e., "use of novel compounds as drugs" and "novel use of publicly known compounds as drugs" intermingle in claims 1 to 4, 24, 25, 28, and 29. Such being the case, a group of inventions of claims 1-4, 24, 25, 28 and 29 and a group of inventions of claims 5 to 23 are not considered as relating to a group of inventions so linked as to form a single general inventive concept.